



Firefly (Coleoptera, Lampyridae) species from the Atlantic Forest hotspot, Brazil

Stephanie Vaz[‡], Mariana Mendes[‡], Gabriel Khattar[§], Margarete Macedo[‡], Cristina Ronquillo[|], Alejandra Zarzo-Arias^{!¶}, Joaquín Hortal^{#,□,«}, Luiz Silveira[»]

‡ Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

§ Concordia University, Montreal, Canada

| Museo Nacional de Ciencias Naturales (MNCN-CSIC), Madrid, Spain

!¶ Universidad de Oviedo, Asturias, Spain

Dept. Biogeography & Global Change, BEI-MNCN-CSIC, Madrid, Spain

□ Universidade Federal de Goiás, Goiânia, Brazil

« Faculdade de Ciências da Universidade de Lisboa, Lisboa, Portugal

» Western Carolina University, Cullowhee, United States of America

Corresponding author: Cristina Ronquillo (cristinaronquillo@mncn.csic.es)

Academic editor: Vinicius S. Ferreira

Received: 25 Jan 2023 | Accepted: 18 Mar 2023 | Published: 23 Mar 2023

Citation: Vaz S, Mendes M, Khattar G, Macedo M, Ronquillo C, Zarzo-Arias A, Hortal J, Silveira L (2023) Firefly (Coleoptera, Lampyridae) species from the Atlantic Forest hotspot, Brazil. Biodiversity Data Journal 11: e101000. <https://doi.org/10.3897/BDJ.11.e101000>

Abstract

Background

We compiled a database of firefly species records from the Atlantic Forest hotspot in Brazil and made it available at GBIF. Data were gathered from literature and from several key entomological collections, including: Coleção entomológica Prof. José Alfredo Pinheiro Dutra (DZRJ/UFRJ) and Coleção do Laboratório de Ecologia de Insetos from Universidade Federal do Rio de Janeiro (CLEI/UFRJ); Coleção Entomológica do Instituto Oswaldo Cruz (CEIOC); Museu de Zoologia da Universidade de São Paulo (MZSP); Coleção Entomológica Pe. Jesus Santiago Moure from Universidade Federal do Paraná (DZUP/UFRPR); and Coleção Entomológica from Universidade Federal Rural de Pernambuco (UFRPE). This database represents the largest contribution to a public repository of recorded occurrences from Neotropical fireflies.

New information

This dataset shows the occurrence and abundance of firefly species in the Atlantic Forest hotspot. Firefly species endemic to this biome are also present and considered in the study. These data can assist scientific and societal needs, by supporting future research projects and conservation decision-making.

Keywords

Neotropics, dataset, species occurrences, endemism, rainforest, South America

Introduction

The Atlantic Forest covers seventeen of twenty-six Brazilian States and occupies almost the whole country's coast (Oliveira and Engel 2017). It is recognised as one of the thirty-six hotspots in the world and is a global priority for biodiversity conservation (Esser et al. 2019). Unfortunately, the Atlantic Forest also represents the biome with the highest number of fauna and flora species threatened by urban sprawl and land-use change. Landscape modification and climate change are strong drivers of habitat loss and changes in correlated ecosystem services (Maldaner et al. 2021). Due to this, the Atlantic Forest is one of the biomes most threatened by climate change (Scarano and Ceotto 2015). The evident shifts in temperature and precipitation patterns (Pereira et al. 2010) have started to induce responses on several species of plants and animals (Root et al. 2003, Parmesan 2006, VanDerWal et al. 2012). Thus, the Atlantic Forest is one of the habitats highly prone to the adverse effects of climate change, jeopardising its biodiversity and endemic species richness. According to the Intergovernmental Panel on Climate Change, the vulnerability to climate change disruption in a system is extreme (I.P.C.C. 2014). In this scenario, it is important to anticipate the effects of climate change in the ecosystems, in order to decrease its damage by building up possible solutions, aligned with mitigation. These strategies can reduce the vulnerability of species and increase the resilience of natural and human systems, aiming to avoid or minimise these negative impacts (I.P.C.C. 2014).

Fireflies (Lampyridae) are amongst the many organisms threatened by anthropogenic stressors and climate change in the Atlantic Forest (Vaz et al. 2021, Khattar et al. 2022). They are a cosmopolitan family of about 2,200 species worldwide (Branham 2010), with most of its known diversity found in the Neotropics and the Asian Southeast (Lawrence and Newton 1995). This number of species is largely underestimated due to severe taxonomic impediments, in addition to a lack of targeted sampling and experts, especially in the Neotropics (McDermott 1966, Silveira and Mermudes 2013). Indeed, Neotropical fireflies need massive nomenclatural and curatorial work, which are of utmost importance to facilitate their research and conservation.

Besides their relevance, fireflies are very charismatic for society (Fallon et al. 2021, Prischmann-Voldseth 2022). They are present in music, poems, paintings and other kinds

of arts, being used as a tool to foster conservation of nature through environmental education (Lenko and Papavero 1997, Faust 2004). Furthermore, they play a significant role in the economy of many countries as flagship species in ecotourism (Lewis et al. 2020). In addition, they are sensitive to light pollution and other forms of environmental degradation (Viviani et al. 2010, Owens et al. 2022), being important bioindicators of ecosystems' health (Fallon et al. 2021). Lastly, their medical importance is relevant since they prey on intermediate hosts of waterborne diseases (Viviani 1989).

Even though anthropogenic threats have been investigated, the extent to which climate and land use changes impact fireflies has never been quantitatively assessed anywhere in South America. An important way to protect fireflies from these threats is to protect forest remnants at the Atlantic Forest hotspot, since protected areas act as a shelter for thousands of species. In this regard, mobilising the biodiversity data currently stored in natural history collections and literature can be particularly useful to conduct large-scale analyses in macroecology and conservation, provided that its coverage is adequately assessed and field inventories are conducted to fill in the gaps that are eventually identified (Hortal et al. 2007, Ronquillo et al. 2020).

Shortfalls, particularly in species identities and distribution (Linnean and Wallacean shortfalls, respectively, see Hortal et al. (2015)), hampers the elaboration of predictive models that could provide information on the needs for management of firefly populations. In this regard, one of the main concerns about fireflies is the absence of occurrence records publicly available in repositories; therefore making it difficult to obtain any assessment on their status. Thus, this work aims to compile and to make a public dataset of firefly diversity in Brazil, which would be of high relevance to advance our knowledge of this particular taxa, especially when applying predictive models which could be the base for new and expert-knowledge-based conservation and management actions. To do this, we evaluated and gathered data from collections and fieldwork, as well as from a revision of literature. In addition, this dataset will support the project described below, which aims to propose conservation policies for the Atlantic Forest, identifying threats and predicting effects of global change on firefly communities, sustaining and providing a better global perspective on potential firefly extinctions in South America.

Project description

Title: Keep the forest shiny: mapping threats to provide information for conservation planning of firefly species endemic to the Atlantic Forest hotspot

Personnel: Stephanie Vaz, Mariana Mendes, Gabriel Khattar, Margarete Macedo, Cristina Ronquillo, Alejandra Zarzo-Arias, Joaquín Hortal, Luiz Felipe Silveira

Funding: Coordenação de Aperfeiçoamento de Pessoal de Nível Superior 88887.694716/2022-00

Sampling methods

Description: Information on lampyrid specimens included in this study was obtained from scientific collections, literature and unpublished data from fieldwork research over nine years throughout the Brazilian Atlantic Forest. The samples were evaluated from different areas in the Brazilian coast, mainly from the south-eastern and south regions (Fig. 1), which are the most densely inhabited. Our field expeditions were undertaken between 2013 and 2022 and encompassed the Parque Nacional da Serra dos Órgãos, Parque Nacional da Tijuca, Parque Estadual da Ilha Grande and Parque Estadual da Pedra Branca, all placed within Rio de Janeiro State. The scientific collections digitised contained records of fireflies sampled from the 1885s to the present.




Figure 1. [doi](#)

Fireflies distribution in the Atlantic Forest hotspot.

Sampling description: Scientific collections

- Universidade Federal do Rio de Janeiro - Coleção entomológica Prof. José Alfredo Pinheiro Dutra from Universidade Federal do Rio de Janeiro (DZRJ), Rio de Janeiro, Brazil.
- Universidade Federal do Rio de Janeiro - Coleção do Laboratório de Ecologia de Insetos from Universidade Federal do Rio de Janeiro (CLEI), Rio de Janeiro, Brazil.
- Fiocruz/CEIOC - Coleção Entomológica do Instituto Oswaldo Cruz, Rio de Janeiro, Brazil.

- Universidade Federal do Parana - Coleção Entomológica Pe. Jesus Santiago Moure from Universidade Federal do Paraná (DZUP/UFPR), Paraná, Brazil.
- Museu de Zoologia da Universidade de Sao Paulo (MZUSP), São Paulo, Brazil.
- Universidade Federal Rural de Pernambuco - Coleção entomológica da Universidade Federal Rural de Pernambuco (UFRPE), Pernambuco, Brazil.

Firefly species were analysed under a stereomicroscope and we imaged characters relevant to species identification. All of them were compared to the species-type material and original descriptions, when available. For specimens whose identity could not be confirmed at the species level, we included a morphotype code (“sp1”, “sp2” and so on). This code was standardised by the first author, so all specimens with the same code in the database pertain to the same morphotype. Entomological institutions and their collections are discriminated as provided by GBIF. Museu Nacional/UFRJ institution code holds both Coleção entomológica Prof. José Alfredo Pinheiro Dutra from Universidade Federal do Rio de Janeiro (DZRJ) and Coleção do Laboratório de Ecologia de Insetos from Universidade Federal do Rio de Janeiro (CLEI) collection codes in this paper.

Fieldwork

The surveys at Parque Nacional da Serra dos Órgãos, Parque Nacional da Tijuca, Parque Estadual da Pedra Branca and Parque Estadual da Ilha Grande were conducted following the same protocol. We used Malaise traps (Fig. 2) and active search across all habitat types and seasons, to enhance the coverage and completeness of our sampling effort. This fieldwork methodology (Suppl. material 1) increased the overall success of the surveys (see Silveira et al. (2020)).



Figure 2. [doi](#)

Typical Malaise trap in the Atlantic Forest.

Literature

Most of the data incorporated into this dataset came from our team, which includes two leading lampyrid taxonomists: Stephanie Vaz and Luiz Silveira. Together, SV and LS have been studying and collecting fireflies since 2010 and helped to build one of the most comprehensive and well-curated firefly collections in the Americas. This work includes: (i) the original descriptions of many of the endemic species covered in this dataset and (ii) monographic reviews which confirmed the endemism (to the Atlantic Forest) of several other species. Data were taken from the original source (Silveira and Mermudes 2013, Silveira and Mermudes 2014a, Silveira and Mermudes 2014b, Bocakova et al. 2016, Silveira et al. 2016, Silveira and Mermudes 2017, Silveira et al. 2018, Campello-Gonçalves et al. 2019, Nunes et al. 2019, Silveira et al. 2019, Silveira et al. 2020, Vaz et al. 2020a, Vaz et al. 2020b, Silveira et al. 2021, Vaz et al. 2021a, Bocakova et al. 2022 and Campello-Gonçalves et al. 2022).

Quality control: All records were validated geographically and taxonomically. Coordinates were plotted and revised to verify the geographical location and locality. All scientific names were checked for types and matched to the species information.

Step description: Most specimens deposited in collections were preserved in entomological drawers under dry conditions. The information contained on the specimens' labels was collected to build the dataset. Specimens from our fieldwork are preserved in ethanol > 90% since the date of collection (Fig. 3), which better preserves soft-bodied insects, such as fireflies and also prevents DNA degradation.



Figure 3. [doi](#)

Ethanol > 90% recipient to preserved specimens from our fieldwork.

Geographic coverage

Description: The samples were collected from different areas throughout the Brazilian Atlantic Forest (Fig. 4). The data obtained from entomological collections was georeferenced by using the information in specimens' labels stored in appropriate drawers or jars.



Figure 4. [doi](#)

The Brazilian Atlantic Forest - (A) Cachoeira do Papel - Trilha da Pedra do Sino, Teresópolis, Rio de Janeiro - Brazil (B) Mirante - Trilha da Pedra do Sino, Teresópolis, Rio de Janeiro - Brazil.

Coordinates: -29.78 and -4.11 Latitude; -54.95 and -34.88 Longitude.

Taxonomic coverage

Description: The dataset consists in 3010 records, 36 genera and 107 different species. Additionally, there are 362 records with genus taxonomic status comprising 44 morphotypes which could not be identified at species level with 100% certainty. That means 153 potential species in this dataset representing 7.65% of the species recorded worldwide (see Branham (2010)), although it is necessary to bear in mind that the total number of Lampyridae species worldwide is still underestimated (Silveira and Mermudes 2013).

The most frequently reported species was *Photuris femoralis* Curtis, 1839, which was recovered from 377 locations. *Photinus frater* (Olivier, 1905) was the second most abundant, found at 334 locations, followed by *Lucidota flabellicornis* Fabricius, 1781 (122), *Amydetes apicalis* Germar, 1824 (104) and *Photinus luna* Curtis, 1932 (101).

The five most abundant species were from the subfamily Photurinae and Lampyrinae. The full firefly dataset was recorded from 14 ecoregions in the Atlantic Forest.

Temporal coverage

Notes: 1885–2019

Usage licence

Usage licence: Other

IP rights notes: The dataset in the current work is licensed under a Creative Commons Attribution (CC-BY) 4.0 License.

Data resources

Data package title: Firefly (Coleoptera: Lampyridae) species from the Atlantic Forest hotspot, Brazil

Resource link: <https://www.gbif.org/dataset/f2fafb08-f21e-4d6c-82f7-ed3463ffa03c>

Alternative identifiers: https://ipt.sibbr.gov.br/mnrj/resource?r=af_fireflies_br_01

Number of data sets: 1

Data set name: Firefly (Coleoptera: Lampyridae) species from the Atlantic Forest hotspot, Brazil

Download URL: <https://www.gbif.org/dataset/f2fafb08-f21e-4d6c-82f7-ed3463ffa03c>

Data format: Darwin Core Archive format

Data format version: Darwin Core Archive format

Description: The dataset contains information on fireflies (Coleoptera, Lampyridae) from the Brazilian Atlantic Forest, obtained from our own fieldwork, specimens deposited in scientific collections and data available in literature. The aims and objectives of this project are to build and compile a dataset tailored to: (i) evaluate the coverage and eventual biases of the distributional data available for Atlantic Forest fireflies; and (ii) develop models to provide further insights on how light pollution, land-use and climate changes may affect the firefly species endemic to the Atlantic Forest. The specimens have been lumped into a single dataset, forming the basis of S. Vaz 's PhD thesis. This dataset will be used to investigate priority areas for conservation in the Atlantic Forest and assess the connection amongst firefly populations and communities located in different forest fragments. This dataset is published in GBIF under the licence [CC BY 4.0](https://creativecommons.org/licenses/by/4.0/) and the metadata can be also consulted at Vaz et al. (2022). If you have any questions regarding this dataset, please do not hesitate to contact us via the contact information provided in the metadata.

Column label	Column description
datasetName	The name identifying the dataset from which the record was derived.
occurrenceID	An identifier for the Occurrence (as opposed to a particular digital record of the occurrence). In the absence of a persistent global unique identifier, construct one from a combination of identifiers in the record that will most closely make the occurrenceID globally unique.
basisOfRecord	The specific nature of the data record.
eventDate	The date-time or interval during which an Event occurred. For occurrences, this is the date-time when the event was recorded. Not suitable for a time in a geological context.
year	The four-digit year in which the Event occurred, according to the Common Era Calendar.
month	The integer month in which the Event occurred.
day	The integer day of the month on which the Event occurred.
scientificName	The full scientific name, with authorship and date information, if known. When forming part of an Identification, this should be the name in lowest level taxonomic rank that can be determined.
higherClassification	A list (concatenated and separated) of taxa names terminating at the rank immediately superior to the taxon referenced in the taxon record.
kingdom	The full scientific name of the kingdom in which the taxon is classified.
phylum	The full scientific name of the phylum or division in which the taxon is classified.
class	The full scientific name of the class in which the taxon is classified.
order	The full scientific name of the order in which the taxon is classified.
family	The full scientific name of the family in which the taxon is classified.
genus	The full scientific name of the genus in which the taxon is classified.
specificEpithet	The name of the first or species epithet of the scientificName.
taxonRank	The taxonomic rank of the most specific name in the scientificName.
identificationRemarks	Comments or notes about the Identification.
identifiedBy	A list (concatenated and separated) of names of people, groups or organisations who assigned the Taxon to the subject.
decimalLatitude	The geographic latitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location. Positive values are north of the Equator, negative values are south of it. Legal values lie between -90 and 90, inclusive.
decimalLongitude	The geographic longitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location. Positive values are east of the Greenwich Meridian, negative values are west of it. Legal values lie between -180 and 180, inclusive.

geodeticDatum	The ellipsoid, geodetic datum or spatial reference system (SRS), upon which the geographic coordinates given in decimalLatitude and decimalLongitude are based.
georeferencedBy	A list (concatenated and separated) of names of people, groups or organisations who determined the georeference (spatial representation) for the Location.
georeferenceProtocol	A description or reference to the methods used to determine the spatial footprint, coordinates and uncertainties.
continent	The name of the continent in which the Location occurs.
country	The name of the country or major administrative unit in which the Location occurs.
countryCode	The standard code for the country in which the Location occurs.
stateProvince	The name of the next smaller administrative region than country (state, province, canton, department, region etc.) in which the Location occurs.
county	The full, unabbreviated name of the next smaller administrative region than stateProvince (county, shire, department etc.) in which the Location occurs.
locality	Less specific geographic information can be provided in other geographic terms (higherGeography, continent, country, stateProvince, county, municipality, waterBody, island, islandGroup). This term may contain information modified from the original to correct perceived errors or standardise the description.
associatedReferences	A list (concatenated and separated) of identifiers (publication, bibliographic reference, global unique identifier, URI) of literature associated with the Occurrence.
institutionCode	The name (or acronym) in use by the institution having custody of the object(s) or information referred to in the record.
collectionCode	The name, acronym, coden or initialism identifying the collection or dataset from which the record was derived.
recordedBy	A list (concatenated and separated) of names of people, groups or organisations responsible for recording the original Occurrence. The primary collector or observer, especially one who applies a personal identifier (recordNumber), should be listed first.

Acknowledgements

Permission to undertake research in the protected areas mentioned in this paper was granted by the Instituto Chico Mendes de Conservação da Biodiversidade, Ministério do Meio Ambiente, Brazil. We also thank the staff at Parque Estadual da Pedra Branca and our dear friend André Luiz Diniz Ferreira (PEPB). We thank the staff at the Parque Nacional da Serra dos Órgãos and Reserva Ecológica de Guapiaçu for supporting our fieldwork and the staff at Laboratório de Ecologia de Insetos/UFRJ and Laboratório de Entomologia/UFRJ for field assistance. We thank all the curators of DZRJ, DZUP, MZSP, CLEI, UFRPE and CEIOC, for their support during our visits to their respective institutions and Programa de Pós-Graduação em Ecologia for supporting the visitation to the collections. We would like to thank SIBBr for supporting our dataset and to provide

precious technical information to provide the firefly assessments online. S.V. was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES). We acknowledge support of the publication fee by the CSIC Open Access Publication Support Initiative through its Unit of Information Resources for Research (URICI).

Author contributions

SV and MMendes collected, counted specimens, compiled data and wrote the manuscript. GK collected data, CR curation and validation of data and support for submission to GBIF. SV and LS identified and collected specimens. M Macedo supported the fieldwork in Parque Nacional da Serra dos Órgãos and provided lab equipment/technology material for the Lampyridae collection and maintenance and edited the manuscript. AZA and JH helped with data design and management. SV, GK and LS provided specimen data. SV led the writing. All authors edited and approved the manuscript.

References

- Bocakova M, Silveira LFL, Mermudes JR (2016) Systematic review of the firefly genus *Scissicauda* (Coleoptera, Lampyridae, Amydetinae) from Brazil. *ZooKeys* 558: 55-75. <https://doi.org/10.3897/zookeys.558.6040>
- Bocakova M, Campello-Gonçalves L, Da Silveira LFL (2022) Phylogeny of the new subfamily Cladodinae: neotenic fireflies from the Neotropics (Coleoptera: Lampyridae). *Zoological Journal of the Linnean Society* 195 (4): 1181-1199. <https://doi.org/10.1093/zoolinnean/zlab091>
- Branham MA (2010) Lampyridae. In: Beutel RG, Leschen RA (Eds) *Handbook of Zoology. Arthropoda: Insecta, Coleoptera: Evolution and Systematics (Polyphaga Part)*. IV. 141 - 147 pp.
- Campello-Gonçalves L, Souto P, Mermudes JM, Silveira LL (2019) *Uanauna* gen. nov., a new genus of fireflies endemic to the Brazilian Atlantic forest (Coleoptera: Lampyridae), with key to brazilian genera of Lucidotina. *Zootaxa* 4585 (1). <https://doi.org/10.11646/zootaxa.4585.1.4>
- Campello-Gonçalves L, Vaz S, Mermudes JM, Ferreira AD, Silveira LL (2022) Comparative morphology and key to Amydetinae genera, with description of three new firefly species (Coleoptera, Lampyridae). *ZooKeys* 1114: 131-166. <https://doi.org/10.3897/zookeys.1114.77692>
- Esser LF, Neves D, Jarenkow JA (2019) Habitat-specific impacts of climate change in the Mata Atlântica biodiversity hotspot. *Diversity and Distributions* 25 (12): 1846-1856. <https://doi.org/10.1111/ddi.12984>
- Fallon C, Walker A, Lewis S, Cicero J, Faust L, Heckscher C, Pérez-Hernández C, Pfeiffer B, Jepsen S (2021) Evaluating firefly extinction risk: Initial red list assessments for North America. *PLOS One* 16 (11). <https://doi.org/10.1371/journal.pone.0259379>
- Faust LF (2004) Fireflies as a catalyst for science education. *Integrative and Comparative Biology* 44 (3): 264-265. <https://doi.org/10.1093/icb/44.3.264>

- Hortal J, Lobo J, Jimenez-Valverde A (2007) Limitations of biodiversity databases: Case study on seed-plant diversity in Tenerife, Canary Islands. *Conservation Biology* 21 (3): 853-863. <https://doi.org/10.1111/j.1523-1739.2007.00686.x>
- Hortal J, de Bello F, Diniz-Filho J, Lewinsohn T, Lobo J, Ladle R (2015) Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics* 46 (1): 523-549. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>
- I.P.C.C. (2014) Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. In: I.P.C.C. (Ed.) Summary for policymakers Climate Change 2014: Impacts, adaptation, and vulnerability. Cambridge University Press, Cambridge & New York, 1-32 pp.
- Khattar G, Vaz S, Braga PHP, Macedo M, Silveira LFL (2022) Life history traits modulate the influence of environmental stressors on biodiversity: The case of fireflies, climate and artificial light at night. *Diversity and Distributions* 28 (9): 1820-1831. <https://doi.org/10.1111/ddi.13584>
- Lawrence JF, Newton AF (1995) Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names). In: Pakaluk J, Slipinski SA (Eds) In: Pakaluk J, Slipinski SA (Eds.), *Biology, phylogeny, and classification of Coleoptera: Papers celebrating the 80th birthday of Roy A. Crowson*. Muzeum I Instytut Zoologii PAN, Warszawa. Muzeum I Instytut Zoologii PAN, 849–861 pp.
- Lenko K, Papavero N (1997) *Insetos no Folclore*. Editora Piedade, FAPESP, São Paulo, 467 pp. URL: <https://repositorio.usp.br/item/000973090> [ISBN 13. 978-8586956041]
- Lewis S, Wong C, Owens A, Fallon C, Jepsen S, Thancharoen A, Wu C, De Cock R, Novak M, López-Palafox T, Khoo V, Reed JM (2020) A global perspective on firefly extinction threats. *BioScience* 70 (2): 157-167. <https://doi.org/10.1093/biosci/biz157>
- Maldaner ME, Sobral-Souza T, Prasniewski VM, Vaz-de-Mello F (2021) Effects of climate change on the distribution of key native dung beetles in South American grasslands. *Agronomy* 11 (10). <https://doi.org/10.3390/agronomy11102033>
- McDermott FA (1966) Lampyridae. In: Steel WO, ed. *Coleopterorum catalogus supplementa, Pars 9 (editio secunda)*. s'-Gravenhage [The Hague]: W. Junk. Pars 9.
- Nunes VC, Souto PM, Monteiro RF, Silveira LF (2019) A second species of *Araucariocladus* Silveira & Mermudes, with notes on the variation in antennomere numbers in this genus (Coleoptera: Lampyridae). *Zootaxa* 4571 (4). <https://doi.org/10.11646/zootaxa.4571.4.8>
- Oliveira RE, Engel VL (2017) A restauração florestal na Mata Atlântica: Três décadas em revisão. *Revista Cincia, Tecnologia & Ambiente* 5 (1): 40-48. <https://doi.org/10.4322/2359-6643.05101>
- Owens AS, Van den Broeck M, De Cock R, Lewis S (2022) Behavioral responses of bioluminescent fireflies to artificial light at night. *Frontiers in Ecology and Evolution* 10 <https://doi.org/10.3389/fevo.2022.946640>
- Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics* 37 (1): 637-669. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110100>
- Pereira HM, Leadley PW, Proença V, Alkemade R, Scharlemann JP, Fernandez-Manjarrés JF, Araújo MB, Balvanera P, Biggs R, Cheung WW, Chini L, Cooper HD, Gilman EL, Guénette S, Hurtt GC, Huntington HP, Mace GM, Oberdorff T, Revenga C, Rodrigues P, Scholes RJ, Sumaila UR, Walpole M (2010) Scenarios for global

- biodiversity in the 21st century. *Science* 330: 1496-1501. <https://doi.org/10.1126/science.1196624>
- Prischmann-Voldseth D (2022) Fireflies in Art: Emphasis on Japanese Woodblock Prints from the Edo, Meiji, and Taishō Periods. *Insects* 13 (9). <https://doi.org/10.3390/insects13090775>
 - Ronquillo C, Alves-Martins F, Mazimpaka V, Sobral-Souza T, Vilela-Silva B, G. Medina N, Hortal J (2020) Assessing spatial and temporal biases and gaps in the publicly available distributional information of Iberian mosses. *Biodiversity Data Journal* 8 <https://doi.org/10.3897/bdj.8.e53474>
 - Root T, Price J, Hall K, Schneider S, Rosenzweig C, Pounds JA (2003) Fingerprints of global warming on wild animals and plants. *Nature* 421 (6918): 57-60. <https://doi.org/10.1038/nature01333>
 - Scarano FR, Ceotto P (2015) Brazilian Atlantic forest: impact, vulnerability, and adaptation to climate change. *Biodiversity and Conservation* 24 (9): 2319-2331. <https://doi.org/10.1007/s10531-015-0972-y>
 - Silveira L, Khattar G, Souto P, Mermudes J, Takiya D, Monteiro R (2016) Integrative taxonomy of new firefly taxa from the Atlantic Rainforest. *Systematics and Biodiversity* 14 (4): 371-384. <https://doi.org/10.1080/14772000.2016.1153006>
 - Silveira LFL, Mermudes J (2013) *Memoan ciceroi* gen. et sp. nov., a remarkable new firefly genus and species from the Atlantic Rainforest (Coleoptera: Lampyridae). *Zootaxa* 3640 (1). <https://doi.org/10.11646/zootaxa.3640.1.6>
 - Silveira LFL, Mermudes J (2014a) Systematic review of the firefly genus *Amydetes* Illiger, 1807 (Coleoptera: Lampyridae), with description of 13 new species. *Zootaxa* 3765 (3). <https://doi.org/10.11646/zootaxa.3765.3.1>
 - Silveira LFL, Mermudes J (2014b) *Ybytyramoan*, a new genus of fireflies (Coleoptera: Lampyridae, Lampyrinae, Photinini) endemic to the Brazilian Atlantic Rainforest, with description of three new species. *Zootaxa* 3835 (3). <https://doi.org/10.11646/zootaxa.3835.3.2>
 - Silveira LFL, Mermudes JRM (2017) A new tropical montane firefly genus and species, active during winter and endemic to the southeastern Atlantic Rainforest (Coleoptera: Lampyridae). *Zootaxa* 4221 (2). <https://doi.org/10.11646/zootaxa.4221.2.4>
 - Silveira LFL, Souto P, Mermudes JRM (2018) Four new species of *Luciuranus* fireflies from the Brazilian Atlantic Rainforest (Coleoptera: Lampyridae). *Zootaxa* 4413 (1). <https://doi.org/10.11646/zootaxa.4413.1.7>
 - Silveira LFL, Rosa SP, Vaz S, Mermudes JM (2019) Systematic review of the firefly genus *Lucernuta* Laporte, 1833 (Coleoptera: Lampyridae). *Annales Zoologici* 69 (2). <https://doi.org/10.3161/00034541anz2019.69.2.003>
 - Silveira LFL, Khattar G, Vaz S, Wilson V, Souto P, Mermudes JM, Stanger-Hall K, Macedo M, Monteiro R (2020) Natural history of the fireflies of the Serra dos Órgãos mountain range (Brazil: Rio de Janeiro) – one of the ‘hottest’ firefly spots on Earth, with a key to genera (Coleoptera: Lampyridae). *Journal of Natural History* 54: 275-308. <https://doi.org/10.1080/00222933.2020.1749323>
 - Silveira LFL, Roza AS, Vaz S, Mermudes J (2021) Description and phylogenetic analysis of a new firefly genus from the Atlantic Rainforest, with five new species and new combinations (Coleoptera: Lampyridae: Lampyrinae). *Arthropod Systematics & Phylogeny* 79: 115-150. <https://doi.org/10.3897/asp.79.e67185>

- VanDerWal J, Murphy H, Kutt A, Perkins G, Bateman B, Perry J, Reside A (2012) Focus on poleward shifts in species' distribution underestimates the fingerprint of climate change. *Nature Climate Change* 3 (3): 239-243. <https://doi.org/10.1038/nclimate1688>
- Vaz S, Da Silveira LFL, Rosa SP (2020a) Morphology and life cycle of a new species of *Psilocladus* Blanchard, 1846 (Coleoptera, Lampyridae, Psilocladinae), the first known bromeliad-inhabiting firefly. *Papéis Avulsos de Zoologia* <https://doi.org/10.11606/1807-0205/2020.60.special-issue.24>
- Vaz S, Mermudes JRM, Paiva PC, Da Silveira LFL (2020b) Systematic review and phylogeny of the firefly genus *Dilychnia* (Lampyridae: Lampyrinae), with notes on geographical range. *Zoological Journal of the Linnean Society* 190 (3): 844-888. <https://doi.org/10.1093/zoolinnean/zlaa041>
- Vaz S, Guerrazzi MC, Rocha M, Faust L, Khattar G, Mermudes J, Lima da Silveira LF (2021a) On the intertidal firefly genus *Micronaspis* Green, 1948, with a new species and a phylogeny of Cratomorphini based on adult and larval traits (Coleoptera: Lampyridae). *Zoologischer Anzeiger* 292: 64-91. <https://doi.org/10.1016/j.jcz.2021.01.002>
- Vaz S, Manes S, Gama-Maia D, Silveira L, Mattos G, Paiva P, Figueiredo M, Lorini ML (2021b) Light pollution is the fastest growing potential threat to firefly conservation in the Atlantic Forest hotspot. *Insect Conservation and Diversity* 14 (2): 211-224. <https://doi.org/10.1111/icad.12481>
- Vaz S, Mendes M, Khattar G, Macedo M, Ronquillo C, Zarzo-Arias A, Hortal J, Silveira L, et al. (2022) Firefly (Coleoptera: Lampyridae) species from the Atlantic Forest hotspot, Brazil Version 1.7. Museu Nacional / UFRJ. In: GBIF, et al. (Ed.) Occurrence dataset. <https://doi.org/10.15468/p7qfev>
- Viviani VR (1989) Descrição dos estágios imaturos e dados biológicos de *Aspisma* sp. *Revista Brasileira de Entomologia* 33: 359-366.
- Viviani VR, Rocha MY, Hagen O (2010) Fauna de besouros bioluminescentes (Coleoptera: Elateroidea: Lampyridae; Phengodidae, Elateridae) nos municípios de Campinas, Sorocaba-Votorantim e Rio Claro-Limeira (SP, Brasil): biodiversidade e influência da urbanização. *Biota Neotropica* 10 (2): 103-116. <https://doi.org/10.1590/s1676-06032010000200013>

Supplementary material

Suppl. material 1: Malaise trap in Atlantic Forest [doi](#)

Authors: Stephanie Vaz et al.

Data type: multimedia

Brief description: How to collect fireflies by using a Malaise trap.

[Download file](#) (28.36 MB)