



Taxonomic paper

# New records for Albania based on taxa from the Prespa National Park

Lulëzim Shuka<sup>†</sup>, Kit Tan<sup>‡</sup>

<sup>†</sup> Tirana University, Tirana, Albania

<sup>‡</sup> University of Copenhagen, Copenhagen, Denmark

Corresponding author: Kit Tan ([kitt@bio.ku.dk](mailto:kitt@bio.ku.dk))

Academic editor: Dimitrios Koureas

Received: 27 Oct 2013 | Accepted: 02 Dec 2013 | Published: 13 Dec 2013

Citation: Shuka L, Tan K (2013) New records for Albania based on taxa from the Prespa National Park. Biodiversity Data Journal 1: e1014. doi: [10.3897/BDJ.1.e1014](https://doi.org/10.3897/BDJ.1.e1014)

## Abstract

Twelve taxa are enumerated as new and three taxa confirmed for the flora of Albania. They were collected between 2007 and 2012 in the Prespa National Park of Albania which is part of the Prespa International Park, a biological protected area at the borders with F.Y.R. Macedonia and Greece. Four taxa, viz., *Centaurea galicicae*, *Festuca galicicae*, *Laserpitium ochridanum* and *Micromeria cristata* subsp. *kosaninii* are restricted to Dry and Galičica Mountains. *Centaurea decora*, a recently described species, is treated as a synonym of *C. soskæ* thus extending the known localities of the latter to the southeast. Detailed information on distribution, occurrence and habitats in Albania are provided for each taxon.

## Keywords

New records, flora, endemics, Prespa National Park, Albania

## Introduction

The Prespa National Park in Albania is part of Prespa International Park (here abbreviated to PIP), a region designated for nature conservation at the borders of Albania, Greece and

F.Y.R. Macedonia (Fig. 1). PIP lies between the geographical coordinates 40°43' to 40°51'N and 20°00' to 21°10'E and has a total surface area of 2519.1 km<sup>2</sup> including lakes Megali and Mikri Prespa. The park is considered to be an integrated ecosystem of global significance unique for its habitats, rich floristic biodiversity and high number of local and Balkan plant endemics. According to Pavlides (Pavlides 1997) the Greek part of the PIP has c. 1326 plant species, the Macedonian part c. 1597 taxa (Matevski 2010) and from current investigation, the Albanian part has at least 1130 species.




Figure 1.

Map of area showing distribution of selected species

The terrestrial area of the Albanian part covers 228 km<sup>2</sup>, much of which comprises the east- and southeast-facing karstic slopes of a ridge called Mali i Thatë or Dry Mountain; this continues in Macedonia under the name Galičica Mt. The slopes of the Mikri and Megali Prespa watershed have steep inclinations of 50–85% and altitudes rising from 848 m at the lake surface to 2288 m at Pllaja e Pusit, the highest peak of Dry Mt.

*Terra rossa* overlying limestone covers more than 95% of the ground area and only a few hectares near Zagradeci village at Mikri Prespa are predominantly serpentine. The climate is mostly moderate-continental as in central Europe and there are Mediterranean-like warm and dry periods alternating with very cold and wet ones. The lakes, high altitude and west-facing slopes of Dry Mt play an important role in creating and influencing the sub-Mediterranean-continental character of the National Park (Mersinllari 1997, Shuka et al. 2008). Climate and soil composition has also favoured the development of a rich flora with Mediterranean, Balkan and Central European floristic elements. All species which are recorded in the Albanian side of the park belongs to 438 genera and 101 plant families. This constitutes nearly 33% of the total flora of Albania within such a small area of 228 km<sup>2</sup>. According to Mersinllari (Mersinllari 1997), Balkan endemics account for c. 12% of the flora of Dry Mt and most of these occur in the upper montane zone. More than 30 taxa have been described from Galičica and Dry Mts, and 10–13 of these are considered restricted to Galičica Mt (Matevski 2013, pers. comm.).

Our study was based on fieldwork in the Prespa National Park, carried out jointly with Til Dieterich from Baku State University (Azerbaijan) during the years 2007, 2011 and 2012. More than 300 specimens which could not easily be named in the field were collected, identified and deposited in the herbarium of Tirana University (TIR). Relevant literature and herbarium material from JE, SKO and TIR were checked and all the species which have their *locus classicus* in F.Y.R. Macedonia were investigated in their original locality. The families and species in the following text are listed in alphabetical order.

## Taxon treatments

### *Laserpitium ochridanum* Micevski 1981

#### Nomenclature

#### *Apiaceae*

*Laserpitium ochridanum* Micevski in Godišen Zbornik, Biološki Fakultet Univerzitet 'Kiril i Metodij', Skopje 34:26 (1981). Fig. 2



Figure 2.

*Laserpitium ochridanum*

Type:— [F.Y.R. MACEDONIA] Stara Galičica, in saxosis calcareis, 2010 m, 16 July 1968, *Micevski* (holotype SKO!).

#### Materials

- a. country: Albania; verbatimLocality: Dry Mt, north of former military post; verbatimElevation: 1640 m; verbatimLatitude: 40°55'N; verbatimLongitude: 20°51'E; eventDate: 8 July 2011; recordedBy: Dieterich & Shuka; basisOfRecord: observation

- b. country: Albania; verbatimLocality: Dry Mt, eastern slopes above Pikina Voda; verbatimElevation: 1885–1900 m; verbatimLatitude: 40°54'N; verbatimLongitude: 20°50'E; eventDate: 8 July 2011; recordedBy: Shuka; basisOfRecord: observation
- c. country: Albania; verbatimLocality: Ivani Mt; verbatimElevation: 1710 m; verbatimLatitude: 40°44'N; verbatimLongitude: 20°53'E; eventDate: 18 July 2012; recordNumber: 5620; recordedBy: Shuka; institutionCode: TIR!

## Distribution

This species has been reported only from Galičica Mt. in F.Y.R. Macedonia. It has not yet been found in NW Greece.

## Ecology

**Phenology:** Flowering in June to mid-July, fruiting mid-July to August.

**Habitat:** Calcareous northeastern slopes of Dry and Ivani Mts, in subalpine meadows, stony places or at the border of tree line and subalpine pasture. Found in association with *Achillea holosericea*, *Aster linosyris*, *Brachypodium sylvaticum*, *Dianthus cruentus*, *D. carthusianorum*, *Iberis sempervirens*, *Festuca paniculata* and *Thymus boissieri*.

## Taxon discussion

Related to *Laserpitium siler* L. particularly subsp. *zernyi* (Hayek) Tutin from N Albania and F.Y.R. Macedonia (Tutin 1967:31). The plant is easily distinguished by its aromatic, dissected leaves with segments 15–20 (–30) x 5–8 (–15) mm. *L. siler* subsp. *zernyi* has leaves with segments 60–90 x 10–30 mm. The populations of *L. ochridanum* in Dry and Ivani Mts comprise less than a hundred individuals in each locality. The discovery on Ivani Mt extends the distribution range c. 30 km south of the *locus classicus*. Plants on Ivani Mt are usually stemless by mid-July being bitten down by sheep and thus these populations are at risk. — Rare, new for Albania.

## *Centaurea galicicae* Micevski, 1985

### Nomenclature

### Asteraceae

*Centaurea galicicae* Micevski in Acta Botanica Croatica 44:83 (1985). Fig. 3



Figure 3.

*Centaurea galicicae*

Type:— [F.Y.R. MACEDONIA] Mt Galičica, in rupestribus calcareis inter pagum Stenje et Konjsko, 20 June 1980, *Micevski* (holotype SKO!).

#### Material

- a. country: Albania; verbatimLocality: Calcareous rocky cliffs of the Gollomboçi Peninsula in Lake Megali Prespa, 7–15 m above the lake shore; verbatimElevation: 855 m; verbatimLatitude: 40°51'N; verbatimLongitude: 20°57'E; eventDate: 13 July 2011; recordNumber: 5426; recordedBy: Shuka; institutionCode: TIR!

#### Distribution

Previously known only from the type locality.

#### Ecology

**Phenology:** Flowering in June to early July, fruiting from mid-July to August.

**Habitat:** The new locality in Albania is further south than the type locality in F.Y.R. Macedonia but habitat and altitude are similar. The limestone slopes of the peninsula were covered with *Cephalaria ambrosioides*, *Centaurea soskae*, *C. graeca* var. *ceccariniana*, *Ephedra fragilis*, *Euphorbia characias* subsp. *wulfenii*, *Lilium chalcedonicum*, *Micromeria juliana*, *Satureja montana*, *Sedum acre* and *Umbilicus luteus*. Less than 50 individuals of the *Centaurea* were noted but they are not threatened by grazing or human activities. — Rare, new for Albania.

***Centaurea rupestris* subsp. *kozanii* Routsis & T. Georgiadis, 1994****Nomenclature*****Asteraceae***

*Centaurea rupestris* L. subsp. *kozanii* Routsis & T. Georgiadis in Candollea 49(2):368 (1994). Fig. 4



Figure 4.  
*Centaurea rupestris* subsp. *kozanii*

Type:— [GREECE. Nomos & eparchia Kozanis] 6 km outside Kozani, on road to Ptolemais, 5 July 1989, *Georgiadis & Routsis* 7320 (holotype UPA).

**Material**

- a. country: Albania; verbatimLocality: Prespa area, Cerja Pass, between the villages of Zaroshka and Cerja; verbatimElevation: 1110 m; verbatimLatitude: 40°45'N; verbatimLongitude: 20°56'E; eventDate: 15 July 2012; recordedBy: Shuka; collectionID: 6415; institutionCode: TIR!

**Distribution**

*Centaurea rupestris* comprises several subspecies in the Balkans. *Centaurea rupestris* subsp. *kozanii* occurs mainly on limestone substrate in NC Greece. It had been misidentified as *C. rupestris* subsp. *parnonia* (Halácsy) (Gugler 1908:194) which was described from the summit area of Mt Parnon (Megali Tourla) in the Peloponnese, southern Greece (the type of *C. parnonia* Halácsy (Halácsy 1898:648) is *Orphanides* 19/31 July 1858, ATHU, WU-Hall!).

## Ecology

**Phenology:** Flowering in June and early July, fruiting mid-July to August.

**Habitat:** In clearings of open *Quercus trojana* forest or in limestone pastures with *Eryngium campestre*, *Teucrium polium* and various grasses, in a small area of less than one hectare. The population at the Cerja Pass is endangered, mainly by grazing cows. — New for Albania.

## *Centaurea soskae* Hayek ex Kosanin, 1926

### Nomenclature

### Asteraceae

*Centaurea soskae* Hayek ex Košanin in Glasnik Srpska Kraljevska Akademija 119 (54):27 (1926). Fig. 5



Figure 5.

*Centaurea soskae*

Type:— [F.Y.R. MACEDONIA] supra pagum Trpezica (=Trpejca) ad lacum Ochrida, solo calcareo, *Soska* (holotype BEOU).

### Materials

- a. country: Albania; verbatimLocality: Dry Mt, above the villages of Korita and Shengjergji, rocky cliffs; verbatimElevation: 1200 m; eventDate: 9 July 1959; recordNumber: s. n; recordedBy: *Palikuqi*; institutionCode: TIR!
- b. country: Albania; verbatimLocality: Guri i Shengjergjit (Rock of Shengjergji); verbatimElevation: 900 m; eventDate: 22 June 1971; recordNumber: s. n; recordedBy: *Vangjeli & Tartari*; institutionCode: TIR!

- c. country: Albania; verbatimLocality: western slopes of Dry Mt; verbatimElevation: 928–1060 m; verbatimLatitude: 40°46'N; verbatimLongitude: 20°49'E; eventDate: 13 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- d. country: Albania; verbatimLocality: Gollomboçi Peninsula, Lake Megali Prespa; verbatimElevation: 855 m; verbatimLatitude: 40°51'N; verbatimLongitude: 20°57'E; eventDate: 13 July 2011; recordedBy: *Shuka*; basisOfRecord: observation
- e. country: Albania; verbatimLocality: Lake Mikri Prespa; verbatimElevation: 930 m; verbatimLatitude: 40°40'N; verbatimLongitude: 20°59"E; eventDate: 14 July 2011; recordNumber: 5520; recordedBy: *Shuka*; institutionCode: TIR!

## Distribution

Previously known only from the type locality near lake Ohrid. However, it has been reported from the western slopes of Dry Mt above Shengjergji village in Albania (Vangjeli et al. 1995:84).

## Ecology

**Phenology:** Flowering and fruiting June to July.

**Habitat:** These localities are at lower altitudes (850–1200 m) and influenced by the Mediterranean and sub-Mediterranean climate, and the moderating effects of the Devolli and Drini Rivers. The occurrence on the rocky calcareous cliff faces of lakes Megali and Mikri Prespa extends the distribution range eastwards towards the lakes. *Allium flavum* subsp. *flavum*, *Campanula versicolor*, *Fumana procumbens*, *Iris germanica*, *Nepeta spruneri*, *Ptilostemon afer*, *Salvia officinalis*, *Satureja montana*, *Sempervivum ciliatum*, as well as the woody species *Buxus sempervirens*, *Fraxinus ornus*, *Pistacia terebinthus* and *Prunus webbii* were also noted on the cliff faces. Based on habitat and ecology, we believe that *Centaurea soskai* occurs and should be looked for in the Greek part of the Prespa lakes. — Confirmed for Albania and new for the Prespa National Park.

## Taxon discussion

In 2011, Meyer described *C. decora* (Meyer 2011:167) as a new species of *Centaurea* from the rocky slopes above the villages of Shengjergji and Korita (Type:— Pogradec, Südabfall des Mali i Thatë, 800–1000 m, 5 July 1959, F.K. Meyer 3486 (holotype JE, digital specimen image!). We collected plants from the same slopes on 13 July 2011 and compared them with living plants of *C. soskai* from the *locus classicus* in F.Y.R. Macedonia, and concluded *C. decora* is identical to *C. soskai*. Meyer (2011) did not mention *C. soskai* in his publication and probably had not seen any material of the latter to realize the two taxa are conspecific.

***Helichrysum luteoalbum* (L.) Rchb., 1929****Nomenclature*****Asteraceae***

*Helichrysum luteoalbum* (L.) Rchb., Handbuch der Gewächskunde, ed. 2, 2:1460 (1829). Fig. 6



Figure 6.

*Helichrysum luteoalbum*

Basionym: *Gnaphalium luteoalbum* L., Sp. Pl. 2:851 (1753). Lectotype designated by Hilliard & Burt in Botanical Journal of the Linnean Society 82:206, 244 (1981):— Herb. A. van Royen no. 900.286-294 (L, digital specimen image!).

**Material**

- a. country: Albania; verbatimLocality: Lake shore of Megali Prespa, from the old church of Zaroshka village up to near the Greek border; verbatimElevation: 850 m; verbatimLatitude: 40°46'N; verbatimLongitude: 20°56'E; eventDate: 16 July 2012; recordNumber: 6466; recordedBy: Shuka; institutionCode: TIR!

**Distribution**

Widely distributed cosmopolitan weed, naturalized in New World. Recorded in almost every country in southern Europe but not yet for Albania.

**Ecology**

**Phenology:** Flowering in June and July.

**Habitat:** Sandy and stony calcareous shore, 3–4 m above the lake. The sparse vegetation includes *Calamintha nepeta*, *Crepis* spp., *Euphorbia falcata*, *Potentilla supina* and *Sonchus* spp., and is often submerged when the water level rises. — New for Albania.

***Tephroseris integrifolia* subsp. *aucheri* (DC.) B. Nord.**

**Nomenclature**

***Asteraceae***

*Tephroseris integrifolia* (L.) Holub subsp. *aucheri* (DC.) B. Nord. in Opera Botanica 44:44 (1978).

Type:— [Turkey, NW Anatolia] Alpes Olymp. Byth. [Ulu Dag], *Aucher-Eloy* 3424 (G, MPU!).

**Material**

- a. country: Albania; verbatimLocality: Dry Mt, Pllaja e Pusit, alpine meadows near the peak; verbatimElevation: 1900–2250 m; verbatimLatitude: 40°54'N; verbatimLongitude: 20°49'E; eventDate: 7 July 2011; recordNumber: 5334; recordedBy: *Shuka*; institutionCode: TIR!

**Distribution**

This subspecies occurs in Serbia, F.Y.R. Macedonia, northern Greece, Bulgaria and NW Anatolia. It is widely distributed in northern Greece at altitudes of 1000–2400 m but has not yet been reported from the Greek or F.Y.R. Macedonia parts of the PIP. Some collections from southern Greece (Peloponnese and S Pindos: *Baden & al.* 954, ATH! *Aldén* 3447, LD!), erroneously identified as subsp. *aucheri*, refer to subsp. *integrifolia*.

**Ecology**

**Phenology:** Flowering in July, fruiting from end of July to August.

**Habitat:** Dry alpine pastures or snowbed meadows, particularly on the western slopes of the mountain ridge. It is usually found in association with *Astragalus lacteus*, *Botrychium lunaria*, *Coeloglossum viride*, *Crocus cvijicii*, *Erysimum kuemmerlei* and *Poa alpina* — New for Albania.

***Alkanna noneiformis* Griseb., 1844****Nomenclature*****Boraginaceae***

*Alkanna noneiformis* Griseb., Spicilegium Florae Rumelicae et Bithynicae 2(4):90 (1844). Fig. 7



Figure 7.

*Alkanna noneiformis*

Lectotype designated by Strid in Mountain Flora of Greece 2:41(1991):— [GREECE] sparsim in herbosis m. Nidgé [Piperitsa] pr. Vodená, 2700'–3000', (substr. marmor.), 28 June 1839, *Grisebach 720* (GOET).

**Materials**

- a. country: Albania; verbatimLocality: Dry Mt, eastern slopes below Pllaja e Pusit; verbatimElevation: 1922 m; verbatimLatitude: 40°54'N; verbatimLongitude: 20°50'E; eventDate: 30 May 2012; recordedBy: *Shuka*; basisOfRecord: observation
- b. country: Albania; verbatimLocality: above Gorica e Madhe village, north of former military post; verbatimElevation: 1600 m; verbatimLatitude: 40°54'N; verbatimLongitude: 20°51'E; eventDate: 30 May 2012; recordNumber: 5720; recordedBy: *Shuka*; institutionCode: TIR!

**Distribution**

The *locus classicus* is Mt Piperitsa which is c. 10 km south of the present Greek–F.Y.R. Macedonian border and thus still within Greek territory, and not in F.Y.R. Macedonia as attributed by Rechinger (Rechinger 1965:209). This species occurs from S and N Pindos to NC Greece and in the southern part of F.Y.R. Macedonia.

## Ecology

**Phenology:** Flowering late May to mid-June, fruiting late June and July.

**Habitat:** Alpine and subalpine meadows overlying limestone on the rocky eastern and northeastern slopes of Dry Mt, usually between 1500 and 2000 m. At lower altitudes it occurs with *Daphne oleoides*, *Genista radiata*, *Juniperus communis* subsp. *alpina* and *J. oxycedrus*. Although rare it does not seem to be under any threat as it occurs within the protected central zone of the National Park.

## Taxon discussion

It is closely related to *A. scardica* (Grisebach 1844:91) from N Albania, F.Y.R. Macedonia, Kosovo and Montenegro; the latter differs by its completely eglandular indumentum, longer calyx, subglabrous corolla and distinctly reticulate nutlets. — Rare, new to Albania.

## *Hesperis theophrasti* Borbás, 1902

### Nomenclature

### *Brassicaceae*

*Hesperis theophrasti* Borbás in Magyar Botanikai Lapok 1:267 (1902). Fig. 8



Figure 8.

*Hesperis theophrasti*

Lectotype designated by Dvořák in Preslia 38: 62 (1996):— [GREECE, S Pindos] Pindus Tymphaeus. In silva ad monaster. Witomo, 15 May 1896, *Sintenis* 1896:221 (BPU 110036; islectotypes BPU, BRNM! LD! P, PR! PRC! W!).

### Materials

- a. country: Albania; verbatimLocality: Dry Mt; verbatimElevation: 1430 m; verbatimLatitude: 40°49'N; verbatimLongitude: 20°53'E; eventDate: 22 May 2006; recordedBy: *Shuka*; basisOfRecord: observation
- b. country: Albania; verbatimLocality: south of Dry Mt, near village of Shengjergji; verbatimElevation: 1000–1300 m; eventDate: 5 July 1959; recordNumber: 3525; recordedBy: *F.K. Meyer*; institutionCode: JE, digital specimen image!
- c. country: Albania; verbatimLocality: Ivani Mt; verbatimElevation: 1680 m; verbatimLatitude: 40°44'N; verbatimLongitude: 20°54'E; eventDate: 29 May 2012; recordNumber: 5496; recordedBy: *Shuka*; institutionCode: TIR!

### Distribution

This is *Hesperis theophrasti* subsp. *theophrasti*, which occurs mainly in the central part of the Balkan Peninsula, Bulgaria and Anatolia (Parolly and Tan 2006). It was recently reported from Albania (Meyer 2011) in a locality south of Dry Mt and outside the area of the National Park.

### Ecology

**Phenology:** Flowering from mid-May to early June, fruiting June to July.

**Habitat:** At altitudes from 1000 m on Dry Mt to nearly 1700 m on Ivani Mt. The species is often found in openings of *Fagus* or *Quercus* forest, stony and rocky meadows with shallow soil or in limestone rock crevices. In *Fagus* forest it occurs together with *Cephalanthera longifolia*, *Iberis sempervirens*, *Paeonia daurica* and *Viola kitaibeliana*. In the other habitats, it is in association with *Acanthus spinosus*, *Delphinium fissum*, *Fritillaria graeca* subsp. *thessala*, *Hypericum rumeliacum*, *Prunus prostrata*, *P. webbii*, *Valeriana montana*, *Viola eximia* and *V. tricolor*. — Confirming Meyer's report from Albania but outside the National Park.

## *Edraianthus horvatii* Lakusic, 1973

### Nomenclature

### Campanulaceae

*Edraianthus horvatii* Lakušić in Godišnjak Biološkog Instituta Univerziteta u Sarajevu 26:44 (1973). Fig. 9



Figure 9.

*Edraianthus horvatii*

Type:— [F.Y.R. MACEDONIA] Galičica, inter 1600 et 2000 m.s.m., solo calcareo, *R. Lakušić* (holotype IBUS).

### Materials

- a. country: Albania; verbatimLocality: Dry Mt, Maja e Ballamaqit; verbatimElevation: 1983 m; verbatimLatitude: 40°47'N; verbatimLongitude: 20°52'E; eventDate: 16 June 2007; recordedBy: *Shuka*; basisOfRecord: observation
- b. country: Albania; verbatimLocality: Buza e Koritës peak; verbatimElevation: 1860 m; verbatimLatitude: 40°47'N; verbatimLongitude: 20°51'E; eventDate: 10 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- c. country: Albania; verbatimLocality: Pllaja e Pusit (near border); verbatimElevation: 2200 m; verbatimLatitude: 40°52'N; verbatimLongitude: 20°50'E; eventDate: 8 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- d. country: Albania; verbatimLocality: Kurrizi i Oçait, Dry Mt; verbatimElevation: 2169 m; verbatimLatitude: 40°52'N; verbatimLongitude: 20°50'E; eventDate: 8 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- e. country: Albania; verbatimLocality: Bear Cave, Dry Mt; verbatimElevation: 1950 m; verbatimLatitude: 40°54'N; verbatimLongitude: 20°50'E; eventDate: 8 July 2011; recordNumber: 5743; recordedBy: *Shuka*; institutionCode: TIR!

### Distribution

This species was previously thought to be a local endemic of Galičica Mt in F.Y.R. Macedonia. It has since been found on Mt Jablanica (a limestone massif east of Mt Shebeniku near the border with southwestern F.Y.R. Macedonia) and Mt Boutsis in northern Greece. It probably occurs on other limestone mountains, e.g., a collection from the summit of Mt Cajupi near Gjirokaster in southern Albania has yet to be verified. We now confirm its occurrence in Albania based on collections first made in

2007 from the central part of Dry Mt, and in 2011 from other localities on the mountain ridge. The localities in Albania adjoin those in F.Y.R. Macedonia.

### Ecology

**Phenology:** Flowering mid-June to mid-July, depending on altitude and exposition.

**Habitat:** Rock crevices and ledges of calcareous cliffs in subalpine and alpine zone. Often together with *Arabis bryoides*, *Asperula doerfleri*, *Coeloglossum viride*, *Oxytropis dinarica*, *O. purpurea*, *Saxifraga* spp., *Sempervivum ciliosum*, *Sideritis raeseri*, *Thlaspi bellidifolium* and *Viola eximia*. — New for Albania.

## *Astragalus mayeri* Micevski, 1970

### Nomenclature

#### *Fabaceae*

*Astragalus mayeri* Micevski in Fragmenta Botanica Musei Macedonici Scientiarum Naturalium 7(17):164 (1970). Fig. 10



Figure 10.

*Astragalus mayeri*

Type:— [F.Y.R. MACEDONIA] Galičica Planina, in rupestribus alpinis, 1200–2000 m, solo calcareo, 16 July 1968, *Micevski* (holotype SKO!).

### Materials

- a. country: Albania; verbatimLocality: Dry Mt, peak of Buza e Koritës; verbatimElevation: 1965 m; verbatimLatitude: 40°47'N; verbatimLongitude: 20°50'E; eventDate: 11 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation

- b. country: Albania; verbatimLocality: peak of Shëngjergji; verbatimElevation: 1765-2000 m; verbatimLatitude: 40°44'N; verbatimLongitude: 20°52'E; eventDate: 11 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- c. country: Albania; verbatimLocality: Ivani Mt; verbatimElevation: 1546 m; verbatimLatitude: 40°43'N; verbatimLongitude: 20°53'E; eventDate: 6 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- d. country: Albania; verbatimLocality: Dry Mt; verbatimElevation: 1893 m; verbatimLatitude: 40°45'N; verbatimLongitude: 20°51'E; eventDate: 11 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- e. country: Albania; verbatimLocality: peak of Zvezda; verbatimElevation: 1704 m; verbatimLatitude: 40°45'N; verbatimLongitude: 20°51'E; eventDate: 11 July 2011; recordNumber: 5574; recordedBy: *Shuka*; institutionCode: TIR!

### Distribution

Also in NC and EC Greece. Undoubtedly very close to *A. sericophyllus* (Grisebach 1843:52) and plants of the latter from Mt Iti in Sterea Ellas, central Greece have been reported as *A. mayeri* (Karetsos 2002:100). The legumes of *A. mayeri* are striped like a zebra as the black and white hairs are confined to patches or in separate rows instead of intermixed.

### Ecology

**Phenology:** Flowering late June to mid-July, fruiting July to August.

**Habitat:** In *Sempervivum–Jovibarba* communities on dry, stony slopes and rocky pastures of Ivani Mt and the southern part of Dry Mt, from 1546 to 2100 m. On Ivani Mt, *A. mayeri* occurs in clearings of *Acer heldreichii* and *Prunus prostrata*. On the upper slopes of Dry Mt, it occurs in rocky places and pastures together with *Astragalus angustifolius*, *Dianthus cruentus*, *D. deltoides* subsp. *degenii*, *D. haematocalyx* subsp. *pindicola*, *Helichrysum plicatum*, *Iris attica*, *Juniperus communis* subsp. *alpina* and *Stachys germanica*. *Erodium guicciardii*, *Sempervivum ciliosum* and *Sideritis raeseri* occur on both sides of the Zvezda Pass which links the southern end of Dry Mt with Ivani Mt.

### Taxon discussion

The plants of *A. mayeri* on Ivani Mt are more robust and larger than those from Galičica or Dry Mt especially from the latter where overgrazing is rampant and the herbaceous cover sparse. — New for Albania.

***Micromeria cristata* subsp. *kosaninii* (Šilić) Bräuchler & Govaerts, 2008****Nomenclature*****Lamiaceae***

*Micromeria cristata* (Hampe) Griseb. subsp. *kosaninii* (Šilić) Bräuchler & Govaerts in Willdenowia 38(2):374 (publ. 18 Dec 2008). Fig. 11



Figure 11.

*Micromeria cristata* subsp. *kosaninii*

Type:— [F.Y.R. MACEDONIA] Galičica Mt, Poljce, c. 1600 m, solo calcareo, 11 October 1970, Šilić (holotype SARA, isotype LJU).

**Material**

- a. country: Albania; verbatimLocality: Dry Mt, rocky cliffs above Korita village;  
verbatimElevation: 1000 m; verbatimLatitude: 40°46'N; verbatimLongitude: 20°51'E;  
eventDate: 13 July 2011; recordNumber: 5529; recordedBy: Shuka; institutionCode: TIR!

**Distribution**

Previously considered endemic to F.Y.R. Macedonia.

**Ecology**

**Phenology:** Flowering mid-June to early July, fruiting July to August.

**Habitat:** Rocky limestone slopes at Lake Ohrid in F.Y.R. Macedonia, and the western slopes of Dry Mt, only a few metres from the locality of *Centaurea soskae*. The population in the cliffs above Korita had fewer than 70 individuals which were found

together with *Campanula versicolor*, *Cynoglossis barrelieri* subsp. *serpentinicola*, *Iris germanica*, *Salvia officinalis* and *Xeranthemum annuum*. — Rare, new for Albania.

### ***Stachys plumosa* Griseb., 1844**

#### **Nomenclature**

#### ***Lamiaceae***

*Stachys plumosa* Griseb., Spicilegium Florae Rumelicae et Bithynicae 2(4):139 (1844).  
Fig. 12



Figure 12.

*Stachys plumosa*

Type:— [F.Y.R. MACEDONIA] inter Komanova et Strazin (substr. trachyt.), *Friedrichsthal* 447 (GOET).

#### **Materials**

- a. country: Albania; verbatimLocality: Prespa area, near the village Cerja and to the NW of Kapshtica village; verbatimElevation: 1170 m; verbatimLatitude: 40°37'N; verbatimLongitude: 21°01'E; eventDate: 6 July 2012; recordedBy: *Shuka*; basisOfRecord: observation
- b. country: Albania; verbatimLocality: near Zagradeci village, Lake Mikri Prespa; verbatimElevation: 880 m; verbatimLatitude: 40°51'N; verbatimLongitude: 20°57'E; eventDate: 13 July 2011; recordedBy: *Shuka*; basisOfRecord: observation
- c. country: Albania; verbatimLocality: northwestern slopes of Ivani Mt; verbatimElevation: 1233 m; verbatimLatitude: 40°44'N; verbatimLongitude: 20°53'E; eventDate: 6 July 2011; recordNumber: 5587; recordedBy: *Shuka*; institutionCode: TIR!

## Distribution

*S. plumosa* has not yet been reported from Albanian territory, although known from the F.Y.R. Macedonia and Greek parts of the PIP. It is a Balkan endemic with a wide distribution on mainland Greece (northern Greece to S Pindos), W Bulgaria and F.Y.R. Macedonia.

## Ecology

**Phenology:** Flowering at the end of May till mid-June, fruiting in June and July.

**Habitat:** *Stachys plumosa* was observed for the first time in Albania near a spring in a serpentine area near the village Zgradeci at Mikri Prespa. It occurs at altitudes between 880 and 1350 m, in dry pastures and rocky places in association with *Alkanna pindicola*, *Buxus sempervirens*, *Comandra umbellata* subsp. *elegans*, *Convolvulus elegantissimus*, *Dianthus haematocalyx* subsp. *pindicola*, *Erodium guicciardii*, *Fraxinus ornus*, *Haplophyllum boissierianum*, *Helichrysum plicatum*, *Hyssopus officinalis* and *Ostrya carpinifolia*. It was also recently recorded in clearings of *Carpinus* and *Buxus* on the limestone slopes of Ivani Mt and near Kapshtica village, close to the border with Greece. — New for Albania.

## *Orobanche purpurea* Jacq., 1762

### Nomenclature

### *Lamiaceae*

*Orobanche purpurea* Jacq., Enumeratio Stirpium plerarumque, quae sponte crescunt in agro Vindobonensi 108, 252 (1762). Fig. 13



Figure 13.

*Orobanche purpurea*

Type:— Described from Austria, 'in collibus siccioribus, supra Weinhaus'.

### Materials

- a. country: Albania; verbatimLocality: Dry Mt, above Gorrica e Madhe (near the border with F.Y.R. Macedonia), Gropat e Palates; verbatimElevation: 1820 m; verbatimLatitude: 40° 54'N; verbatimLongitude: 20°50'E; eventDate: 8 July 2011; recordNumber: 5518; recordedBy: *Shuka*; institutionCode: TIR!
- b. country: Albania; verbatimLocality: Shengjergji, south of Dry Mt; verbatimElevation: 1000–1300 m; eventDate: 5 July 1959; recordNumber: 3546; recordedBy: *F.K. Meyer*; institutionCode: JE, digital specimen image!
- c. country: Albania; verbatimLocality: N Albanian Alps, Shtegu i Dheneve, Thethi; verbatimElevation: 1300 m; eventDate: 23 July 1959; recordNumber: 4085; recordedBy: *F.K. Meyer*; institutionCode: JE, digital specimen image!

### Distribution

Occurring almost throughout Europe and SWAsia. It has recently been reported by Meyer (Meyer 2011) from N Albania and Dry Mt.

### Ecology

**Phenology:** Flowering June and July, fruiting July to August.

**Habitat:** Dry, subalpine calcareous pastures or rocky slopes, parasitic on *Achillea holosericea*. — Confirming occurrence in Albania; new to the PIP, including the parts belonging to F.Y.R. Macedonia and Greece.

## *Festuca galicicae* Markgr.-Dann., 1978

### Nomenclature

### *Poaceae*

*Festuca galicicae* Markgr.-Dann. in Botanical Journal of the Linnean Society 76(3):324 (1978). Fig. 14



Figure 14.  
*Festuca galicicae*

Type:— F.Y.R. Macedonia] Galičica Planina, 2210 m, 9 July 1939, *Horvat* (holotype ZA).

### Materials

- a. country: Albania; verbatimLocality: Dry Mt, N of Pllaja e Pusit; verbatimElevation: 2108 m; verbatimLatitude: 40°53'N; verbatimLongitude: 20°50'E; eventDate: 8 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- b. country: Albania; verbatimLocality: Sheshi i Rinisë; verbatimElevation: 1863 m; verbatimLatitude: 40°48'N; verbatimLongitude: 20°51'E; eventDate: 10 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- c. country: Albania; verbatimLocality: Buza e Korites; verbatimElevation: 1904 m; verbatimLatitude: 40°47'N; verbatimLongitude: 20°50'E; samplingProtocol: 10 July 2011; recordedBy: *Dieterich & Shuka*; basisOfRecord: observation
- d. country: Albania; verbatimLocality: southern part of Dry Mt; verbatimElevation: 1817–1900 m; verbatimLatitude: 40°49'N; verbatimLongitude: 20°51'E; eventDate: 11 July 2011; recordNumber: 5825; recordedBy: *Shuka*; institutionCode: TIR!

### Distribution

Previously considered endemic to F.Y.R. Macedonia.

### Ecology

**Phenology:** Flowering in the first half of July.

**Habitat:** Alpine pastures between 1820 and 2200 m together with *Astragalus lacteus*, *Helictotrichon convolutum*, *Carex kitaibeliana*, *Poa annua*, *Gnaphalium hoppeanum*, *Onobrychis viciifolia*, *Sesleria coerulea*, *Rhinanthus nigricans*, and *Viola eximia*. — New to Albania.

***Viola eximia* Formánek, 1900****Nomenclature*****Violaceae***

*Viola eximia* Formánek in Verhandlungen des Naturforschenden Vereins in Brünn 38:221 (1900). Fig. 15



Figure 15.  
*Viola eximia*

Type:— Described from Mt Kajmakčalan on the Greek–F.Y.R. Macedonia border (BRNM).

**Materials**

- a. country: Albania; verbatimLocality: Dry Mt, Maja e Zonjës; verbatimElevation: 1969 m; verbatimLatitude: 40°49'N; verbatimLongitude: 20°52'E; eventDate: 22 May 2006; recordedBy: *Shuka*; basisOfRecord: observation
- b. country: Albania; verbatimLocality: Pllaja e Pusit; verbatimElevation: 2224 m; verbatimLatitude: 40°54'N; verbatimLongitude: 20°49'E; eventDate: 7 July 2011; recordedBy: *Shuka*; basisOfRecord: observation
- c. country: Albania; verbatimLocality: Ivani Mt; verbatimElevation: 1750 m; verbatimLatitude: 40°44'N; verbatimLongitude: 20°53'E; eventDate: 9 July 2011; recordedBy: *Shuka*; basisOfRecord: observation
- d. country: Albania; verbatimLocality: Ivani Mt; verbatimElevation: 1662 m; verbatimLatitude: 40°43'N; verbatimLongitude: 20°53'E; eventDate: 29 May 2012; recordNumber: 5312; recordedBy: *Shuka*; institutionCode: TIR!

## Distribution

Occurring in the mountains of N Greece (NC, Prespa National Park), F.Y.R. Macedonia, Dry and Ivani Mts in Albania, from 1600 to 2250 m. *Viola eximia* subsp. *tringiana* Erben occurs on Mt Tringia in S Pindos.

## Ecology

**Phenology:** Flowering mid-May to early June, fruiting in July.

**Habitat:** Subalpine and alpine pastures of the Prespa watershed. *V. eximia* is an early flowering species found in the dry pastures and rocky slopes of Dry Mt and in clearings of scrub or dwarf scrub on Ivani Mt. Several interesting plants were found in full flower in both areas, viz., *Achillea abrotanoides*, *Aethionema saxatile*, *Arabis bryoides*, *Asphodeline taurica*, *Barbarea bracteosa*, *Centaurea cana*, *Crocus cvijicii*, *C. chrysanthus*, *Cytisus tommasinii*, *Fritillaria montana*, *Morina persica*, *Ornithogalum umbellatum*, *Orobanche gracilis*, *Oxytropis purpurea*, *Saxifraga scardica*, *Stachys germanica*, *Thymus boissieri*, *T. longicaulis* and *Vicia onobrychioides*. — New to Albania.

## Acknowledgements

Fieldwork was carried out with the help of Til Dieterich and supported by Kreditanstalt für Wiederaufbau (KfW Bankengruppe) and the Ministry of Environment, Forestry and Water Administration (MoEFWA, Albania) in the frame of the project 'Transboundary biosphere reserve Prespa — support to the Prespa National Park in Albania'. We are grateful to Wolfgang Fremuth (Frankfurt) and Thimaq Lako (Tirana) for co-ordinating the project. Vasil Male and other personnel of the Prespa National Park are thanked for their generous assistance. Prof. V. Matevski (Macedonian Academy of Sciences, Skopje) kindly identified our material of *Festuca galicicae*.

## References

- Grisebach A (1843) Spicilegium Florae Rumelicae et Bithynicae exhibens synopsis plantarum quas in aest. 1839 legit auctor A. Grisebach 1. F. Vieweg & Son, Brunsvigae, 407 pp.
- Grisebach A (1844) Spicilegium Florae Rumelicae et Bithynicae exhibens synopsis plantarum quas in aest. 1839 legit auctor A. Grisebach 2(4). F. Vieweg & Son, Brunsvigae, 160 pp.
- Gugler W (1908) Die Centaureen des Ungarischen National-Museums. Vorarbeiten zu einer Monographie der Gattung *Centaurea* und der ihr nächst verwandten Genera. Annales Historico-Naturales Musei Nationalis Hungarici 6: 15-297.
- Halácsy Ev (1898) Die bisher bekannten *Centaurea*-Arten Griechenlands. Bulletin Herb. Boissier 6(7-8): 565-603, 633-659.
- Karetos G (2002) Meleti tis ikologias ke tis vlastisis tou orous Iti. University of Patras, Patras, 311 pp.

- Matevski V (2010) Report on Flora and Vegetation of Galicica National Park in Management Plan of National Park Galicica, including species list. Macedonian Academy of Sciences, Skopje, 33 pp.
- Mersinllari M (1997) General data on vegetation of Ohrid and Prespa watershed areas. Proceedings of International Symposium on Towards integrated conservation and sustainable development of transboundary Makro-Mikro Prespa lakes, Korça, Albania, 24–26 October 1997: 147-155.
- Meyer FK (2011) Beiträge zur Flora von Albanien. Haussknechtia (Beih.) 15: 1-220.
- Parolly G, Tan K (2006) A new species of *Hesperis* (Brassicaceae) from SW Anatolia, Turkey. Willdenowia 36: 851-856.
- Pavlides G (1997) Aquatic and terrestrial vegetation of the Prespa area. Hydrobiologia 351: 41-60. DOI: [10.1023/A:1003091805384](https://doi.org/10.1023/A:1003091805384)
- Reehner KH (1965) Zur Kenntnis der europäischen Arten der Gattung *Alkanna*. Annalen des Naturhistorischen Museums in Wien 68: 191-220.
- Shuka L, Malltezi J, Mersinllari M, Vardhami I (2008) Dynamics of vegetation cover of Prespa Lakes and its watershed (Albanian side). Proceedings of the Third International Scientific Conference on Balkan Water Observation and Information System (BALWOIS), 27–31 May 2008, Ohrid, Macedonia. 8 pp.
- Tutin TG (1967) New combinations and new names in Umbelliferae. In: Heywood, VH (Ed.) Notulae systematicae ad Floram Europaeam spectantes, no. 6. Repertorium specierum novarum regni vegetabilis 74(1-2): 31-34.
- Vangjeli J, Ruci B, Mullaj A (1995) Red Book: Threatened and rare plant species of Albania. Academy of Science, Tirana, 169 pp.