Data Paper

A dataset of fishes in and around Inle Lake, an ancient lake of Myanmar, with DNA barcoding, photo images and CT/3D models



‡ Institute of Decision Science for Sustainable Society, Kyushu University, Fukuoka, Japan

§ Department of Fishery Biology, Faculty of Fisheries, Kasetsart University, Chatuchak, Bangkok, Thailand

| Laboratory of Ecology and Environment, Division of Southeast Asian Area Studies, Graduate School of Asian and African Area Studies, Kyoto University, Kyoto, Japan

¶ Inle Lake Wildlife Sanctuary, Nature and Wildlife Conservation Division, Forest Department, Ministry of Natural Resources and Environmental Conservation, the Republic of the Union of Myanmar, Nyaung Shwe, Myanmar

Department of Zoology, Taunggyi University, Taunggyi, Myanmar

¤ Laboratory of Animal Ecology, Graduate School of Science, Kyoto University, Kyoto, Japan

- « Osaka Museum of Natural History, Osaka, Japan
- » Yamashina Institute for Ornithology, Abiko, Japan

Corresponding author: Katsutoshi Watanabe (watanak@terra.zool.kyoto-u.ac.jp)

Academic editor: Pavel Stoev

Received: 17 Sep 2016 | Accepted: 03 Nov 2016 | Published: 09 Nov 2016

Citation: Kano Y, Musikasinthorn P, Iwata A, Tun S, Yun L, Win S, Matsui S, Tabata R, Yamasaki T, Watanabe K (2016) A dataset of fishes in and around Inle Lake, an ancient lake of Myanmar, with DNA barcoding, photo images and CT/3D models. Biodiversity Data Journal 4: e10539. <u>https://doi.org/10.3897/BDJ.4.e10539</u>

Abstract

Background

Inle (Inlay) Lake, an ancient lake of Southeast Asia, is located at the eastern part of Myanmar, surrounded by the Shan Mountains. Detailed information on fish fauna in and around the lake has long been unknown, although its outstanding endemism was reported a century ago.



New information

Based on the fish specimens collected from markets, rivers, swamps, ponds and ditches around Inle Lake as well as from the lake itself from 2014 to 2016, we recorded a total of 948 occurrence data (2120 individuals), belonging to 10 orders, 19 families, 39 genera and 49 species. Amongst them, 13 species of 12 genera are endemic or nearly endemic to the lake system and 17 species of 16 genera are suggested as non-native. The data are all accessible from the document "A dataset of Inle Lake fish fauna and its distribution (http:// ipt.pensoft.net/resource.do?r=inle fish 2014-16)", as well as DNA barcoding data (mitochondrial COI) for all species being available from the DDBJ/EMBL/GenBank (Accession numbers: LC189568–LC190411). Live photographs of almost all the individuals and CT/3D model data of several specimens are also available at the graphical fish biodiversity database (http://ffish.asia/INLE2016; http://ffish.asia/INLE2016-3D). The information can benefit the clarification, public concern and conservation of the fish biodiversity in the region.

Keywords

Myanmar; Shan State; Inle Lake; freshwater fishes; endemic species; alien; GBIF; mitochondrial DNA; COI; CT scan; 3D model

Introduction

Inle Lake is located on the southwestern part of Shan State, which is the easternmost state of Myanmar. The lake is surrounded by Shan Hills, which isolate it from the neighbouring aquatic habitats. The lake harbours several endemic fish species (Annandale 1918). However, the detailed information of fish fauna of this region has long been unknown since Annandale (1918), while several studies reported the concerns for settlement of non-native species and decline of endemic/native species (Musikasinthorn 1998, Su and Jassby 2000, Davies et al. 2004, Oo 2010). This project aimed to elucidate the current status of fish fauna of the lake, especially focusing on survival of endemic/native species and settlement of non-native species. In addition, DNA sequences (mitochondrial COI), photo images and CT/3D models were published online in the public interest of biodiversity.

Project description

Title: Current status, origin and conservation of endemic fishes in an ancient lake, Inle Lake.

Personnel: Katsutoshi Watanabe (Project director, fieldwork and DNA barcoding), Prachya Musikasinthorn (fieldwork, fish identification, photographs and specimens management), Yuichi Kano (fieldwork, photographs and database management), Akihisa Iwata (fieldwork and fish identification), Shoko Matsui (DNA barcoding), Ryoichi Tabata (DNA barcoding), Sein Tun (management and local information), LKC Yun (fieldwork and local information), Seint Seint Win (fieldwork and local information) and Taksehi Yamasaki (CT scanning).

Study area description: Inle Lake is located on the Southwestern part of Shan State, which is the easternmost state in Myanmar. The lake is surrounded by Shan Hills and harbours several endemic fish species. The surveys were carried out in the lake and its surroundings from 23 September 2014 until 2 July 2016. In addition, local markets were also investigated for fishes caught in the study area.

Design description: This study focused on fish fauna of Inle Lake, a representative ancient lake in mainland Southeast Asia. Until now, the lake has not been investigated for nearly a century (Annandale 1918). In our study, we focused on the condition of endemic species as the environments have been drastically changed (Su and Jassby 2000) and alien species recently settled in the lake (Davies et al. 2004, Oo 2010). As information about the fish species of Inle Lake is quite limited, almost all the specimens were photographed and the photo data were deposited in an online fish database (Kano et al. 2013) (http://ffish.asia/INLE2016). CT/3D models for several specimens were also made and are available from the database (http://ffish.asia/INLE2016-3D). The main bodies were retained as formalin samples for voucher specimens and deposited at Kasetsart University, Thailand. In addition, a small piece of tissue (mainly from the right pectoral or pelvic fin) was excised, preserved in 99% alcohol and stored at Kyoto University to facilitate the study of molecular biology and genetics. The DNA sequences (mitochondrial COI) information was deposited at the DDBJ/EMBL/GenBank (Accession numbers: LC189568–LC190411).

Funding: JSPS (The Japan Society for the Promotion of Science) KAKENHI Grant Number JP26304007.

Sampling methods

Sampling description: The fish samples were either collected from 43 wild habitats as well as from one fish cage in the lake or purchased at 24 local markets (Fig. 1). The fishes were collected by hand-nets, throwing nets, traps and/or purchased from fishermen. In the markets, we avoided purchasing the obvious cultured fishes from other areas (especially from Yangon and Mandalay), such as *Wallago* spp. and *Pangasianodon hypophthalmus*, by checking with the seller.




Figure 1.

Spatial coverage of the sampling points. Solid circles indicate wild environments such as rivers and reservoirs. White circles indicate local markets and a grey circle indicates a fish cage in the lake.

Quality control: All the scientific names of fish samples were validated by the updated fish checklist in the Catalogue of Fishes, California Academy of Sciences (<u>http://www.calacademy.org/scientists/projects/catalog-of-fishes</u>), Kottelat (2013) and/or Nelson et al. (2016). For order level classification of the family Ambassidae, we followed Hastings et al. (2014). Most of the specimens were photographed in a fresh state (Kano and Nakajima 2014) and then the formalin specimens and its tissue samples were catalogued and deposited at the Research Laboratory of Ichthyology, Department of Fishery Biology, Faculty of Fisheries, Kasetsart University, Bangkok, Thailand (RLIKU) and Kyoto University, Kyoto, Japan, respectively. All the samples were assigned the IDs which were associated with the records of location (latitude, longitude and region name), the dates, methods, accession numbers of DNA sequences, etc.

Step description: Step 1: Sampling locality and date were recorded.

Step 2: Specimens were given IDs, photographed, fin-clipped and roughly classified on site.

Step 3: Specimens were fixed in 10% formalin solution for two or more weeks. Subsequently, they were cleaned with water and preserved in 75% ethanol.

Step 4: Specimens were shipped back to the lab for correct species identification.

Taxonomic coverage

Description: Annandale (1918), Akihito and Meguro (1975), Roberts (1986), Kottelat (1990), Talwar and Jhingran (1991), Fang (1997), Roberts (1997), Musikasinthorn (1998), Roberts (1998), Kottelat and Witte (1999), Ng et al. (1999), Kottelat (2001), Fang (2003), Kottelat (2003), Roberts (2007), Ng and Kottelat (2008), Fang et al. (2009), Matsumoto et al. (2009), Britz (2010), Havird and Page (2010), Jiang et al. (2012), Kottelat (2012), Kottelat et al. (2012), Pethiyagoda et al. (2012), Kottelat (2013), Hastings et al. (2014), Ratmuangkhwang et al. (2014) and Nelson et al. (2016) were used as a taxonomic reference for this work. The coverage of this dataset includes Subclass Actinopterygii. The orders are Cypriniformes (26 species), Anabantiformes (5), Siluriformes (5), Synbranchiformes (5), Cyprinodontiformes (2), Perciformes (2), Beloniformes (1), Cichliformes (1), Gobiiformes (1) and Osteoglossiformes (1) (Fig. 2). The families are Cyprinidae (20), Nemacheilidae (4), Ambassidae (2), Channidae (2), Clariidae (2), Mastacembelidae (2), Osphronemidae (2), Poeciliidae (2), Sisoridae (2), Synbranchidae (2), Anabantidae (1), Adrianichthyidae (1), Balitoridae (1), Chaudhuriidae (1), Cichlidae (1), Cobitidae (1), Gobiidae (1), Heteropneustidae (1) and Notopteridae (1) (Fig. 3).





Figure 2. Taxonomic coverage (by order).



Taxonomic coverage (by family).

Taxa included:

Rank	Scientific Name	Common Name
kingdom	Animalia	Animals
phylum	Chordata	Chordates
subphylum	Craniata	Vertebrates and hagfishes
class	Osteichthyes	Bony fishes and tetrapods
subclass	Actinopterygii	Ray-finned fishes
order	Anabantiformes	Labyrinth fishes
order	Beloniformes	Needlefishes
order	Cichliformes	Cichlids and convict blennies
order	Cypriniformes	Carps, loaches, minnows and relatives
order	Cyprinodontiformes	Killifishes
order	Gobiiformes	Gobies
order	Osteoglossiformes	Bonytongues
order	Perciformes	Perches
order	Siluriformes	Catfishes
order	Synbranchiformes	Swamp eels
family	Anabantidae	Climbing gouramies
family	Adrianichthyidae	Adrianichthyids
family	Ambassidae	Asiatic glassfishes
family	Balitoridae	Hillstream loaches
family	Channidae	Snakeheads
family	Chaudhuriidae	Earthworm eels
family	Cichlidae	Cichlids
family	Clariidae	Airbreathing catfishes
family	Cobitidae	True loaches
family	Cyprinidae	Cyprinids
family	Gobiidae	Gobies
family	Heteropneustidae	Airsac catfishes
family	Mastacembelidae	Spiny eels
family	Nemacheilidae	Stone loaches

family	Notopteridae	Knifefishes
family	Osphronemidae	Gouramies and fighting fishes
family	Poeciliidae	Livebearers
family	Sisoridae	Sisorid catfishes
family	Synbranchidae	Swamp eels
species	Anabas testudineus (Bloch 1792)	Climbing gourami
species	Balitora sp.	A species of balitorid loach
species	Barbonymus gonionotus (Bleeker 1849)	Silver barb
species	Celestichthys erythromicron (Annandale 1918)	A species of Celestichthys minnow
species	Channa harcourtbutleri (Annandale 1918)	Inle snakehead
species	Channa striata (Bloch 1793)	Striped snakehead
species	Chaudhuria caudata Annandale 1918	Inle swamp eel
species	Clarias gariepinus (Burchell 1822)	African sharptooth catfish
species	Clarias cf. batrachus (Linnaeus 1758)	Walking catfish
species	Ctenopharyngodon idella (Valenciennes 1844)	Grass carp
species	Cyprinus intha Annandale 1918	Inle carp
species	Cyprinus rubrofuscus Lacepède 1803	Common carp
species	Devario kakhienensis (Anderson 1879)	A species of Devario minnow
species	Devario sp.	A species of Devario minnow
species	Esomus danrica (Hamilton 1822)	Flying barb
species	Gambusia affinis (Baird & Girard 1853)	Western mosquitofish
species	Garra gravelyi (Annandale 1919)	Burmese Garra
species	Glossogobius cf. giuris (Hamilton 1822)	A species of Glossogobius goby
species	Glyptothorax granosus Jiang, Ng, Yang & Chen 2012	A species of sisorid catfish
species	Glyptothorax rugimentum Ng & Kottelat 2008	A species of sisorid catfish
species	Gymnostomus horai (Bănărescu 1986)	A species of Gymnostomus minnow
species	Heteropneustes fossilis (Bloch 1794)	Stinging catfish
species	Inlecypris auropurpureus (Annandale 1918)	A species of Inlecypris minnow
species	Labeo rohita (Hamilton 1822)	Rohu
species	Lepidocephalichthys berdmorei (Blyth 1860)	A species of cobitid loach
species	Mastacembelus caudiocellatus (Boulenger 1893)	A species of spiny eel

species	Mastacembelus oatesii Boulenger 1893	A species of spiny eel
species	Microrasbora rubescens Annandale 1918	Red dwarf rasbora
species	Monopterus cuchia (Hamilton 1822)	Gangetic mud eel
species	Monopterus javanensis Lacepède 1800	Asian swamp eel
species	Neolissochilus nigrovittatus (Boulenger 1893)	A species of Neolissochilus barb
species	Notopterus notopterus (Pallas 1769)	Bronze featherback
species	Oreochromis niloticus (Linnaeus 1758)	Nile tilapia
species	Oryzias uwai Roberts 1998	A species of rice fish
species	Parambassis lala (Hamilton 1822)	A species of Asiatic glassfish
species	Parambassis ranga (Hamilton 1822)	A species of Asiatic glassfish
species	Pethia stoliczkana (Day 1870)	Stoliczka's barb
species	Petruichthys brevis (Boulenger 1893)	A species of nemacheilid loach
species	Physoschistura rivulicola (Hora 1929)	A species of nemacheilid loach
species	Physoschistura shanensis (Hora 1929)	A species of nemacheilid loach
species	Poecilia reticulata Peters 1859	Guppy
species	Poropuntius schanicus (Boulenger 1893)	A species of Poropuntius barb
species	Puntius sophore (Hamilton 1822)	Spotfin swamp barb
species	Puntius cf. sophore (Hamilton 1822)	A species of <i>Puntius</i> barb
species	Sawbwa resplendens Annandale 1918	Burmese rammy nose
species	Schistura sp.	A species of nemacheilid loach
species	Systomus cf. rubripinnis (Valenciennes 1842)	A species of Systomus barb
species	Trichogaster labiosa Day 1877	Thick-lipped gourami
species	Trichopodus pectoralis Regan 1910	Snakeskin gourami

Temporal coverage

Notes: 23 September 2014 – 2 July 2016.

Usage rights

Use license: Creative Commons Public Domain Waiver (CC-Zero)

Data resources

Data package title: A Dataset of Inle Lake Fish Fauna and Its Distribution

Resource link: http://ipt.pensoft.net/resource.do?r=inle_fish_2014-16

Number of data sets: 1

Additional information

Endemic, native and non-native or status uncertain

Inle Lake has an outstanding endemic fish fauna while non-native species have established in and around the lake. Thus, the species were discriminated by endemic, native (but not endemic), non-native and unknown as shown in Fig. 4. In addition, two endemic species reported in Annandale (1918) were not ascertained in this survey: the two species seemed to be very rare or already extinct from the studied area.




Figure 4.

Distribution of the endemic, native, non-native and uncertain species in the studied area.

Endemic: Celestichthys erythromicron; Channa harcourtbutleri; Cyprinus intha; Gymnostomus horai; Inlecypris auropurpureus; Mastacembelus caudiocellatus; Mastacembelus oatesii; Microrasbora rubescens; Neolissochilus nigrovittatus; Petruichthys brevis; Physoschistura shanensis; Poropuntius schanicus; Sawbwa resplendens.

Native (but not endemic): Anabas testudineus; Channa striata; Chaudhuria caudata; Clarias cf. batrachus; Devario kakhienensis; Garra gravelyi; Glyptothorax granosus; Glyptothorax rugimentum; Lepidocephalichthys berdmorei; Monopterus cuchia; Monopterus javanensis; Notopterus notopterus; Pethia stoliczkana; Physoschistura rivulicola; Systomus cf. rubripinnis.

Non-native: Barbonymus gonionotus; Clarias gariepinus; Ctenopharyngodon idella; Cyprinus rubrofuscus; Esomus danrica; Gambusia affinis; Glossogobius cf. giuris; Heteropneustes fossilis; Labeo rohita; Oreochromis niloticus; Oryzias uwai; Parambassis lala; Parambassis ranga; Poecilia reticulata; Puntius sophore; Trichogaster labiosa; Trichopodus pectoralis.

Unknown: Balitora sp.; Devario sp.; Puntius cf. sophore; Schistura sp.

Endemic species unascertained: Systomus compressiformis; Silurus burmanensis.

Acknowledgements

We deeply appreciate the kind assistance from Nyi Nyi Kyaw (Forest Department, Ministry of Natural Resources and Environmental Conservation, Myanmar [FD-MOECAF]), Win Naing Thaw (Nature and Wildlife Conservation Division, FD-MOECAF), the late Than Htay (Inle Lake Wildlife Sanctuary, FD-MOECAF), Ohn and Than Nwai (Forest Resource Environment Development and Conservation Association, Myanmar [FREDA]) and Nyi Nyi Lwin (Green Leaf, Myanmar). 2016 GBIF Asia Biodiversity Informatics Workshop, hosted by TaiBIF and JBIF kindly assisted with the GBIF data registration and submission of the paper.

References

- Akihito P, Meguro K (1975) Description of a new gobiid fish, *Glossogobius aureus*, with notes on related species of the genus. Japanese Journal of Ichthyology 22: 127-142.
- Annandale N (1918) Fish and fisheries of the Inle Lake. Records of the Indian Museum 14: 33-64. <u>https://doi.org/10.5962/bhl.part.18603</u>
- Britz R (2010) Species of the *Macrognathus aculeatus* group in Myanmar with remarks on *M. caudiocellatus* (Teleostei: Synbranchiformes: Mastacembelidae). Ichthyological Exploration of Freshwaters 20: 295-308.
- Davies J, Sebastian AC, Chan S (2004) A wetland inventory for Myanmar. Ministry of the Environment, Japan, Tokyo, 491 pp.
- Fang F (1997) Redescription of *Danio kakhienensis*, a poorly known cyprinid fish from the Irrawaddy basin. Ichthyological Exploration of Freshwaters 7: 289-298.
- Fang F (2003) Phylogenetic analysis of the Asian cyprinid genus *Danio* (Teleostei, Cyprinidae). Copeia 2003: 714-728. <u>https://doi.org/10.1643/ia03-131.1</u>
- Fang F, Noren M, Liao TY, Källersjö M, Kullander SO (2009) Molecular phylogenetic interrelationships of the south Asian cyprinid genera *Danio*, *Devario* and *Microrasbora* (Teleostei, Cyprinidae, Danioninae). Zoologica Scripta 38: 237-256. <u>https:// doi.org/10.1111/j.1463-6409.2008.00373.x</u>
- Hastings P, Walker HJ, Galland G (2014) Fishes : a guide to their diversity. University of California Press, Oakland, 311 pp.

- Havird J, Page L (2010) A revision of *Lepidocephalichthys* (Teleostei: Cobitidae) with descriptions of two new species from Thailand, Laos, Vietnam, and Myanmar. Copeia 2010: 137-159. <u>https://doi.org/10.1643/ci-08-240</u>
- Jiang W, Ng HH, Yang J, Chen X (2012) A taxonomic review of the catfish identified as *Glyptothorax zanaensis* (Teleostei: Siluriformes: Sisoridae), with the descriptions of two new species. Zoological Journal of the Linnean Society 165: 363-389. <u>https://</u> <u>doi.org/10.1111/j.1096-3642.2011.00811.x</u>
- Kano Y, Nakajima J (2014) Non-killing simple photography techniques of small–middle freshwater fishes in the field. Japanese Journal of Ichthyology 61: 123-125. [In Japanese].
- Kano Y, Adnan MS, Grudpan C, Grudpan J, Magtoon W, Musikasinthorn P, Natori Y, Ottomanski S, Praxaysonbath B, Phongsa K, Rangsiruji A, Shibukawa K, Shimatani Y, So N, Suvarnaraksha A, Thach P, Thanh PN, Tran DD, Utsugi K, Yamashita T (2013) An online database on freshwater fish diversity and distribution in Mainland Southeast Asia. Ichthyological Research 60: 293-295. https://doi.org/10.1007/s10228-013-0349-8
- Kottelat M (1990) Indochinese nemacheilines: A revision of nemacheiline loaches (Pisces: Cypriniformes) of Thailand, Burma, Laos, Cambodia and southern Viet Nam. Verlag, Dr. Friedrich Pfiel, Munchen, 262 pp.
- Kottelat M (2001) Fishes of Laos. WHT Publications, Colombo, 198 pp.
- Kottelat M (2003) Nomenclatural status of *Crossocheilus burmanicus*, *C. horai* and *C. multirastellatus* (Osteichthyes: Cyprinidae). Raffles Bulletin of Zoology 51: 399-401.
- Kottelat M (2012) Conspectus cobitidum: an inventory of the loaches of the world (Teleostei: Cypriniformes: Cobitoidei). The Raffles Bulletin of Zoology 26: 1-199.
- Kottelat M (2013) The fishes of inland waters of Southeast Asia: A catalogue and core bibliography of fishes known to occur in freshwaters, mangroves and estuaries. The Raffles Bulletin of Zoology Supplement 27: 1-663.
- Kottelat M, Witte KE (1999) Two new species of *Microrasbora* from Thailand and Myanmar, with two new generic names for small Southeast Asian cyprinid fishes (Teleostei: Cyprinidae). Journal of South Asian Natural History 4: 49-56.
- Kottelat M, Baird I, Kullander SO, Ng H, Parenti L, Rainboth W, Vidthayanon C (2012) The status and distribution of freshwater fishes of Indo-Burma. In: Allen D, Smith K, Darwall W (Eds) The status and distribution of freshwater biodiversity in Indo-Burma. IUCN, Gland.
- Matsumoto S, Kon T, Yamaguchi M, Takeshima H, Yamazaki Y, Mukai T, Kuriiwa K, Kohda M, Nishida M (2009) Cryptic diversification of the swamp eel *Monopterus albus* in East and Southeast Asia, with special reference to the Ryukyuan populations. Ichthyological Research 57: 71-77. <u>https://doi.org/10.1007/s10228-009-0125-y</u>
- Musikasinthorn P (1998) First record of *Parambassis lala* (Pisces: Ambassidae) from Inle Lake, the Salween River Basin, Myanmar. Natural History Bulletin of the Siam Society 46: 203-205.
- Nelson J, Grande T, Wilson MH (2016) Fishes of the world (5th edition). Wiley, New Jersey, 752 pp.
- Ng H, Ng PL, Britz R (1999) *Channa harcourtbutleri* (Annandale, 1918): a valid species of snakehead (Perciformes: Channidae) from Myanmar. Journal of South Asian Natural History 4: 57-63.

- Ng HH, Kottelat M (2008) The identity of *Clarias batrachus* (Linnaeus, 1758), with the designation of a neotype (Teleostei: Clariidae). Zoological Journal of the Linnean Society 153: 725-732. <u>https://doi.org/10.1111/j.1096-3642.2008.00391.x</u>
- Oo AH (2010) Inland fisheries resources enhancement and conservation practice in Myanmar. In: Weimin M, Silva SD, Davy B (Eds) Inland fisheries resource enhancement and conservation in Asia. Food and Agriculture Organization of the United Nations regional office for Asia and the Pacific, Bangkok.
- Pethiyagoda R, Meegaskumbura M, Maduwage K (2012) A synopsis of the South Asian fishes referred to *Puntius* (Pisces: Cyprinidae). Ichthyological Exploration of Freshwaters 23: 69-95.
- Ratmuangkhwang S, Musikasinthorn P, Kumazawa Y (2014) Molecular phylogeny and biogeography of air sac catfishes of the *Heteropneustes fossilis* species complex (Siluriformes: Heteropneustidae). Molecular Phylogenetics and Evolution 79: 82-91. <u>http</u> s://doi.org/10.1016/j.ympev.2014.05.009
- Roberts TR (1986) Systematic review of the Mastacembelidae or spiny eels of Burma and Thailand, with description of two new species of *Macrognathus*. Japanese Journal of Ichthyology 33: 95-109.
- Roberts TR (1997) Systematic revision of the tropical Asian labeoin cyprinid fish genus *Cirrhinus*, with descriptions of new species and biological observations on *C. lobatus*. Natural History Bulletin of the Siam Society 45: 171-203.
- Roberts TR (1998) Systematic observations on tropical Asian medakas or ricefishes of the genus *Oryzias*, with descriptions of four new species. Ichthyological Research 45: 213-224. <u>https://doi.org/10.1007/bf02673919</u>
- Roberts TR (2007) The "celestial pearl danio", a new genus and species of colourful minute cyprinid fish from Myanmar (Pisces: Cypriniformes). The Raffles Bulletin of Zoology 55: 131-140.
- Su M, Jassby A (2000) Inle: a large Myanmar lake in transition. Lakes and Reservoirs: Research and Management 5: 49-54. <u>https://doi.org/10.1046/j.1440-1770.2000.00090.x</u>
- Talwar PK, Jhingran AG (1991) Inland fishes of India and adjacent countries. Vol. 1–2.
 Oxford & IBH Publishing Co., New Delhi, 1158 pp.