



# New records of *Harpiola isodon* (Chiroptera, Vespertilionidae) from the Chinese mainland

Song Li<sup>‡,§</sup>, Xin Mou<sup>‡</sup>, Mengcheng Li<sup>l</sup>, Fengyi Li<sup>¶</sup>, Mei Li<sup>#</sup>, Biao Li<sup>#</sup>, Mengjia Li<sup>#</sup>, Xiong Luo<sup>#</sup>, Gábor Csorba<sup>□</sup>, Haochi Kuo<sup>«</sup>

‡ Kunming Natural History Museum of Zoology, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China  
§ Yunnan Key Laboratory of Biodiversity Information, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China

l State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China

¶ 4-D Genomic Dynamics in Ecology and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China

# Conservation Area Management Committee of Guanyin Shan Provincial Nature Reserve, Yuanyang, China

□ Hungarian Natural History Museum, Department of Zoology, H-1088 Budapest, Baross u. 13, Hungary

« Biodiversity Research Center, Academia Sinica, Taipei, China

Corresponding author: Song Li ([lis@mail.kiz.ac.cn](mailto:lis@mail.kiz.ac.cn))

Academic editor: Ricardo Moratelli

Received: 10 Feb 2024 | Accepted: 14 Mar 2024 | Published: 20 Mar 2024

Citation: Li S, Mou X, Li M, Li F, Li M, Li B, Li M, Luo X, Csorba G, Kuo H (2024) New records of *Harpiola isodon* (Chiroptera, Vespertilionidae) from the Chinese mainland. Biodiversity Data Journal 12: e120670.

<https://doi.org/10.3897/BDJ.12.e120670>

## Abstract

## Background

The new species, *Harpiola isodon* Kuo et al., 2006, was described from Taiwan, China. So far, no distribution of this species outside Taiwan has been reported.

## New information

During two field investigations of small mammals in Guanyin Mountains Provincial Nature Reserve, Yuanyang, Yunnan, China, in April 2022 and May 2023, five individuals of *Harpiola* were collected in the mid-montane evergreen broad-leaved forest. Our morphological and molecular results reveal that these individuals from the Chinese

mainland belong to *Harpiola isodon*, extending the occurrence of this species well beyond its known distributions in Taiwan, China and Vietnam.

## Keywords

distribution, Murinae, tube-nosed bats, Yunnan

## Introduction

Based on a single specimen from northwest India, initially described as *Murina grisea* Peters, 1872, Thomas (1915) established *Harpiola* as a distinct genus and the generic diagnostic characters included: the wing membrane is attached to the base of the first toe; the upper incisors, canine and premolars are all subequal in size; the upper canine is slightly higher than the upper anterior premolar; and the lower canine is shorter than the lower posterior premolar.

Tate (1941) accepted *Harpiola* as a valid genus, but others, such as Ellerman and Morrison-Scott (1951), Corbet and Hill (1992), Koopman (1994) and Bates and Harrison (1997), treated it as a subgenus of *Murina*. Based on a second *H. grisea* specimen collected in Mizoram, India, Bhattacharyya (2002) re-established the genus *Harpiola*, although Simmons (2005) still listed it as a subgenus of *Murina*. With the additional 11 *Harpiola* specimens collected from Taiwan, China, Kuo et al. (2006) described *H. isodon* as a new species. By carefully reviewing the full range of documented morphological variation of *Murina* (e.g. Peters (1880), Ognev (1928), Wallin (1969), Maeda (1980), Yoshiyuki (1989), Corbet and Hill (1992)), Kuo et al. (2006) clarified that some morphological characters, such as the attachment point of the plagiopatagium to the hind foot and the reduced canines in both upper and lower toothrows, cannot completely distinguish *Harpiola* and *Murina*. Instead, Kuo et al. (2006) listed valid diagnostic characters between the two genera as follows: 1, the heights of the inner and outer upper incisors are both two-thirds of the upper canine's height (versus height ratios of up to one-half in *Murina*); 2, the upper toothrow gradually decreases in height from the canine to the first premolar, then to the second premolar, while having these teeth similar in bulk (versus a clearly lower first premolar than the other two teeth in *Murina*); 3, the lower toothrow has the canine, the first premolar and the second premolar similar in both height and bulk (versus a clearly smaller first premolar than the canine in *Murina*); 4, the lower canine is strongly bifid, with the additional cusp well developed (versus a small secondary cingular cusp in *Murina*). In 2006, a *Harpiola* bat was captured in central Vietnam, referred to by Kruskop et al. (2006) as *H. cf. isodon*, representing the first confirmed record of the species outside of Taiwan, China.

During two field investigations of small mammals in Guanyin Mountains Provincial Nature Reserve, Yuanyang, Yunnan, China, in April 2022 and May 2023, five *Murina*-like bats ( $n = 2$  and  $3$ , respectively) were captured with mist nets, showing the diagnostic characters of *Harpiola* as defined by Kuo et al. (2006). As *Harpiola* has never been reported on the

Chinese mainland before, we explored their taxonomic status and carried out comparisons, based on molecular and morphological data as described below.

## Materials and methods

### Sampling

The specimens collected in this study comprise 3 adult males and 2 adult females, according to the development degree of molars and the degree of ossification of forelimb joints. No pregnant females were found. Voucher specimens are deposited in the Kunming Natural History Museum of Zoology, Kunming Institute of Zoology, the Chinese Academy of Sciences (KIZ, CAS), Kunming, China and registered under the numbers KIZ 20220058, 20220089, 20230357, 20230425 and 20230463.

### Molecular analyses

Following the manufacturer's protocol, total genomic DNA was extracted from muscle samples using the Ezup Column Animal Genomic DNA Purification Kit (Sangon Biotech, China). The complete sequence of the mitochondrial *Cyt b* gene was amplified and sequenced with the primer pair LGL765: GAAAAACCAYCGTTGTWATTCAACT and LGL766: GTTTAATAAGAATYTYAGCTTTGGG (Bickham et al. 1995). Polymerase chain reactions (PCR) were carried out in a total volume of 25  $\mu$ l containing 1  $\mu$ l of template DNA, 1  $\mu$ l of each primer at 10  $\mu$ M, 1  $\mu$ l of dNTPmix at 10 mM, 0.2  $\mu$ l of Taq Plus DNA polymerase (Sangon Biotech, China) at 5 U/ $\mu$ l, 2.5  $\mu$ l of 10x Taq Buffer and added H<sub>2</sub>O to 25  $\mu$ l. The PCR thermal profile was: 1, 95°C 5 min for initial denaturation; 2, 94°C 30 sec for denaturation; 3, 63°C (decreases by 0.5°C per cycle) 30 sec for annealing; 4, 72°C 30 sec for extension; 5, 10 cycles of steps 2 to 4; 6, 95°C 30 sec for denaturation; 7, 58°C 30 sec for annealing; 8, 72°C 30 sec for extension; 9, 30 cycles of steps 6 to 8; 10, 72°C 10 min final extension; 11, 4°C for renaturation. PCR products were detected by agarose gel electrophoresis and purified using the SanPrep Column DNA Gel Extraction Kit (Sangon Biotech, China). Finally, purified samples were sequenced by the ABI 3730XL DNA Analyzer (USA) at Sangon Biotech (Shanghai, China). Sequences were edited and assembled using SeqMan in Lasergene 7.1 (DNASTAR Inc., Madison, WI, USA).

The full-length *Cyt b* sequences (1,140 bp) were compared with those from the National Center for Biotechnology Information (NCBI). The sequences were aligned using the default parameters of the ClustalW algorithm in the software MEGA11 (Tamura et al. 2021) and the uncorrected *P*-distances were calculated between pairwise sequences. The pairwise deletion option was used to remove ambiguous positions when calculating genetic distances. The phylogeny of the subfamily Murinae was reconstructed by MEGA11 using the Maximum Likelihood method under a GTR+G+I nucleotide substitution model and the branch support was evaluated by 1,000 bootstrap replicates. ModelFinder (Kalyanamoorthy et al. 2017) on PhyloSuite v.1.2.2 (Zhang et al. 2020) was used to select the best-fit model (GTR+G+I nucleotide substitution model), based on the Bayesian Information Criterion (BIC).

## Morphological characteristics

The morphological characters of the five Yunnan specimens were compared with those described for *Harpiola grisea* and *H. isodon* (Dobson 1878, Thomas 1915, Bhattacharyya 2002, Kuo et al. 2006, Kruskop et al. 2006). We also took external, cranial and dental measurements from our specimens, as described below. Head and body length, tail length, foot length, ear length, forearm length, thumb length, metacarpal lengths and tibia length were measured in the field. The cranial and dental measurements were taken according to Kuo et al. (2006) as follows: total length of skull — from the anterior rim of alveolus of the first upper incisor to the most projecting point of the occipital region; condylobasal length — from the exoccipital condyle to the posterior rim of the alveolus of the first upper incisor; upper canine width— taken across the outer borders of upper canines; upper molar width — taken across the outer crowns of the last upper molars; zygomatic width — the greatest width of the skull across the zygomatic arches; mastoid width — the greatest distance across the mastoid region; postorbital width — the least width of the postorbital constriction; maxillary toothrow length — from the front of the upper canine to the back of the crown of the third molar; upper canine–premolar length — the largest distance from the front of the upper canine to the back of the crown of the posterior premolar; length of mandible — from the anterior rim of the alveolus of the first lower incisor to the most posterior part of the condyle; mandibular toothrow length — from the front of the lower canine to the back of the crown of the third lower molar; lower canine–premolar length — the greatest distance from the front of the lower canine to the back of the crown of the posterior premolar; height of the coronoid process — taken perpendicularly from the extremity of the coronoid process to the ramus mandibulae. Each craniodental measurement was taken three times by Xin Mou with a caliper accurate to 0.01 mm and the average value was reported.

## Ethics statement

Following the Chinese laws and regulations on the protection of wild terrestrial animals ( State Council Decree 1992), the field investigations of small mammals in Guanyin Mountains Provincial Nature Reserve and the collection of specimens were approved by the Conservation Area Management Committee of Guanyin Mountains Provincial Nature Reserve and the Ethics Committee of KIZ, CAS.

## Taxon treatment

### *Harpiola isodon* Kuo et al., 2006

#### Materials

- a. scientificName: *Harpiola isodon* (Kuo et al. 2006); taxonID: <https://www.ncbi.nlm.nih.gov/taxonomy/685777>; kingdom: Animalia; phylum: Chordata; class: Mammalia; order: Chiroptera; family: Vespertilionidae; genus: *Harpiola*; country: China; stateProvince: Yunnan; locality: Guanyinshan Nature Reserve, Mt. Guanyin; verbatimElevation: 2381 m; verbatimCoordinates: 23°1.8'N 102°57'E; decimalLatitude: 23.03; decimalLongitude:

- 102.95; georeferenceProtocol: lable; eventDate: 24-04-22; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: KIZ20220058; recordedBy: Song Li et al.; occurrenceID: C699C5B3-9056-5C72-924C-58E795FC224D
- b. scientificName: *Harpiola isodon* (Kuo et al. 2006); taxonID: <https://www.ncbi.nlm.nih.gov/taxonomy/685777>; kingdom: Animalia; phylum: Chordata; class: Mammalia; order: Chiroptera; family: Vespertilionidae; genus: *Harpiola*; country: China; stateProvince: Yunnan; locality: Guanyinshan Nature Reserve, Mt. Guanyin; verbatimElevation: 2381 m; verbatimCoordinates: 23°1.8'N 102°57'E; decimalLatitude: 23.03; decimalLongitude: 102.95; georeferenceProtocol: lable; eventDate: 24-04-22; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: KIZ20220089; recordedBy: Song Li et al.; occurrenceID: 47008C54-877D-56A6-AF51-08E0D2BF6EC6
- c. scientificName: *Harpiola isodon* (Kuo et al. 2006); taxonID: <https://www.ncbi.nlm.nih.gov/taxonomy/685777>; kingdom: Animalia; phylum: Chordata; class: Mammalia; order: Chiroptera; family: Vespertilionidae; genus: *Harpiola*; country: China; stateProvince: Yunnan; locality: Guanyinshan Nature Reserve, Mt. Guanyin; verbatimElevation: 2463 m; verbatimCoordinates: 22°59.4'N 102°59.4'E; decimalLatitude: 22.99; decimalLongitude: 102.99; georeferenceProtocol: lable; eventDate: 22-05-23; individualCount: 1; sex: female; lifeStage: adult; catalogNumber: KIZ20230357; recordedBy: Song Li et al.; occurrenceID: 4054655C-043F-504B-8EB7-0D8664F2B213
- d. scientificName: *Harpiola isodon* (Kuo et al. 2006); taxonID: <https://www.ncbi.nlm.nih.gov/taxonomy/685777>; kingdom: Animalia; phylum: Chordata; class: Mammalia; order: Chiroptera; family: Vespertilionidae; genus: *Harpiola*; country: China; stateProvince: Yunnan; locality: Guanyinshan Nature Reserve, Mt. Guanyin; verbatimElevation: 2463 m; verbatimCoordinates: 22°59.4'N 102°59.4'E; decimalLatitude: 22.99; decimalLongitude: 102.99; georeferenceProtocol: lable; eventDate: 22-05-23; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: KIZ20230425; recordedBy: Song Li et al.; occurrenceID: A0A6D386-2B57-50FC-A91E-4639AF54638B
- e. scientificName: *Harpiola isodon* (Kuo et al. 2006); taxonID: <https://www.ncbi.nlm.nih.gov/taxonomy/685777>; kingdom: Animalia; phylum: Chordata; class: Mammalia; order: Chiroptera; family: Vespertilionidae; genus: *Harpiola*; country: China; stateProvince: Yunnan; locality: Guanyinshan Nature Reserve, Mt. Guanyin; verbatimElevation: 2463 m; verbatimCoordinates: 22°59.4'N 102°59.4'E; decimalLatitude: 22.99; decimalLongitude: 102.99; georeferenceProtocol: lable; eventDate: 22-05-23; individualCount: 1; sex: female; lifeStage: adult; catalogNumber: KIZ20230463; recordedBy: Song Li et al.; occurrenceID: 5FF3C5EC-DB42-5A77-B109-788CE15590E0

## Description

The five Chinese mainland specimens show the distinguishing dental characters of *Harpiola* as defined above (Fig. 1A2, A3 and A4).

**Body:** A medium-sized bat with the forearm length ranging from 32.33–36.12 mm (Table 1). The wing membrane is attached to the base of the first toe and the interfemoral membrane is attached to the end of the tibia (Fig. 2A and C). The nostril is slightly tubular, but not very prominent (Fig. 2A and C). The ear is small, with the tip slightly blunt and rounded and the tragus is slender and slightly curved, with its length over half that of the ear. The thumb, equipped with a curved, sharp claw, has a length nearly 20% of the forearm length (Fig. 2B and C; Table 1). The third, fourth and fifth metacarpals have about the same length (Table 1). The foot length is about 80% of the

tibia length (Table 1). The tail, which is about 70% of the head and body length (Table 1), has its very end excluded from the interfemoral membrane (Fig. 2A).



Figure 1. [doi](#)

Skull of *Harpiola isodon*. **A** is from Yunnan; **B** is from Taiwan. 1 = ventral view of skull; and 2 = lateral view of skull; 3 = lateral view of mandible; 4 = details of the anterior lower dentition; 5 = occlusal view of mandible; and 6 = details of the occlusal view of lower molars.

Table 1.

External and craniodental measurements of *Harpiola* species (mm) (\*the parameter shows obvious difference between specimens of Yunnan and Taiwan).

Parameter	<i>H. isodon</i> Yunnan, China, This study		<i>H. isodon</i> Taiwan, China, (Kuo et al. 2006)	<i>H. cf. isodon</i> , Vietnam, (Kruskop et al. 2006)	<i>H. grisea</i> , Mizoram, India, (Bhattacharyya 2002)
	Range (n=5)	Mean (n=5)			
Head and body length	44.69-46.50	45.47	-	-	42.7
Tail length	30.03-36.95	31.86	-	-	27.5
Foot length	9.92-10.94	10.26	-	-	8.20
Ear length	12.42-15.17	14.08	12.50-13.00	11.80	12.10

Forearm length	32.33-36.12	34.61	31.00-35.60	31.70	32.40
Thumb length	6.42-6.97	6.65		-	-
The third metacarpal length	29.21-32.21	30.62		-	-
The fourth metacarpal length	29.01-31.95	30.34		-	-
The fifth metacarpal length	28.91-31.85	30.21		-	-
Tibia length	12.14-12.77	12.37		-	14.80
Total length of skull	15.54-16.27	15.87	14.76-16.48	15.41	16.40
Condylobasal length	14.12-15.17	14.83	13.74-14.87	14.49	-
Upper canine width	3.82-4.10	3.95	3.65-4.02	3.98	3.70
Upper molar width	4.80-5.74	5.36	4.90-5.53	5.32	5.50
Zygomatic width	8.72-9.75	9.32	8.43-9.35	8.94	9.40
Mastoid width	7.77-8.21	8.04	7.29-7.96	7.68	-
Postorbital width	4.80-5.03	4.91	4.52-4.84	4.47	4.30
Maxillary toothrow length*	<b>5.74-5.87</b>	<b>5.82</b>	<b>4.97-5.63</b>	5.39	5.30
Upper canine–pre-molar length	2.35-2.64	2.52	2.22-2.73	-	-
Length of mandible	10.57-11.27	10.92	10.15-11.32	10.80	10.60
Mandibular toothrow length	5.63-6.08	5.88	5.35-5.90	5.68	5.70
Lower canine–pre-molar length	2.31-2.47	2.40	2.02-2.44	-	-
Height of the coronoid process	3.33-3.73	3.47	3.36-3.98	-	-

**Fur:** The face is brownish-black at the snout, becoming paler at the cheeks and around the eyes. The forehead is brown and the chin is brownish-black. Ears are naked, dark brown. The fur is soft and dense, with individual hairs on both dorsal and ventral sides yellowish-brown at their upper parts and dark brown at the bases (a little grey). Both dorsal and ventral furs have golden-tipped hairs (Fig. 2) and there are more such hairs on the back than on the abdomen. On the dorsum, the golden-tipped hairs extend from the top of the head to the back of the interfemoral membrane, where they become

sparser. There are yellowish-brown hairs covering the back of the forearm, the back of the thumb and the back of the toes. The golden-tipped hairs on the ventrum are mainly concentrated on the chest. The anal area is brownish-grey. The dorsal and ventral surfaces of the interfemoral membrane are both coated, with thicker, dark brown hairs on the dorsal surface and sparser, light brownish-grey hairs (slightly yellow) on the ventral surface. The foot soles are naked, lacking any sole pad (Fig. 2C). The wing membrane is brownish-black in colour and the area near the side of the body is sparsely coated, otherwise is bare (Fig. 2).




Figure 2. [doi](#)

Individual of *Harpiola isodon* (KIZ20230058) from Yunnan.

**Skull:** In lateral view, the skull shows a gradually rising curve from the front of the rostrum to the back of the frontal area, with the centre of the forehead slightly depressed. The braincase is rounded, the sagittal crest is absent and the lambda crest is moderately developed (Fig. 1A2). The zygomatic arches are weak and slender, showing the most outward expanded points at their posterior ends. The tympanic bullae are relatively small and the foramen magnum is relatively large (Fig. 1, A1).

**Dentition:** Dental formula:  $I \frac{-23}{123} C \frac{1}{1} PM \frac{-2-4}{-2-4} M \frac{123}{123} = 34$ . Upper incisors are large, their heights are more than two-thirds that of  $C^1$  and the outer incisor ( $I^3$ ) is slightly lower than the inner incisor ( $I^2$ ) (Fig. 1, A2).  $C^1$ ,  $PM^2$  and  $PM^4$  are gradually decreasing in height and their base sizes are similar. Amongst the three upper molars, the last one ( $M^3$ ) has a crown area only half that of each anterior one (Fig. 1, A1). The lower toothrows contain three lower incisors on each side, with their heights gradually increasing from the inner ( $I_1$ ) to the outer one ( $I_3$ ). Each lower incisor has three cusps, with a marked depression between the outermost cusp and the middle cusp. The lower canine ( $C_1$ ), with two blunt cusps, is about the same height as the anterior and posterior lower premolars ( $PM_2$  and  $PM_4$ , respectively) (Fig. 1, A4). The crown areas of  $M_1$  and  $M_2$  are slightly larger than  $M_3$ ; the postcristids are well developed and the hypoconids are more prominent than the hypoconulids (Fig. 1, A6).

## Discussion

### Morphological traits

The two species of *Harpiola*, *H. isodon* and *H. grisea*, differ from each other in the following characters: the PM<sup>4</sup> is wider than long in *H. isodon* (versus as wide as long in *H. grisea*); both M<sup>1</sup> and M<sup>2</sup> have the mesostyles present in *H. isodon* (versus M<sup>2</sup> lacking mesostyle in *H. grisea*); M<sup>1</sup> of *H. isodon* has a post-cingular platform (versus no post-cingular platform in *H. grisea*) (Kuo et al. 2006). The five Chinese mainland *Harpiola* specimens showed these dental characters in line with *H. isodon*. We noted that the Chinese mainland taxon had a range of maxillary tooththrow length surpassing that of the Taiwanese *H. isodon* provided in Kuo et al. (2006) (Table 1). Nevertheless, we should warn that the measurements were taken by different people across the two studies and those taken by the same investigator are warranted in the future for a valid evaluation of the morphometric differences between the two taxa.

### Genetic distances and phylogenetic relationships

We compared the *Cyt b* sequences of our five specimens with those downloaded from NCBI for 19 Vespertilionids, including *Harpiola isodon* from Taiwan, *Harpiocephalus harpia*, 15 *Murina*, one *Kerivoula* and one *Myotis* species (Table 2). The novel sequences are deposited in the NCBI GenBank database under accession numbers [PP476123](#) (KIZ 20220058), [PP476124](#) (KIZ20220089), [PP476125](#) (KIZ20230357), [PP476126](#) (KIZ20230425) and [PP476127](#) (KIZ20230463).

Table 2.

Additional samples and their GenBank accession numbers used in the phylogenetic reconstruction.

species	GenBank	species	GenBank
<i>Harpiola isodon</i>	<a href="#">GQ168914</a>	<i>Murina florum</i>	<a href="#">GQ168902</a>
<i>Harpiola isodon</i>	<a href="#">GQ168920</a>	<i>Murina gracilis</i>	<a href="#">GQ168900</a>
<i>Harpiocephalus harpia</i>	<a href="#">GQ168923</a>	<i>Murina recondita</i>	<a href="#">KJ198270</a>
<i>Murina eleryi</i>	<a href="#">GQ168908</a>	<i>Murina bicolor</i>	<a href="#">JQ044696</a>
<i>Murina cyclotis</i>	<a href="#">MK747248</a>	<i>Murina fanjingshanensis</i>	<a href="#">KT180333</a>
<i>Murina suilla</i>	<a href="#">GQ168905</a>	<i>Murina rongjiangensis</i>	<a href="#">MF359930</a>
<i>Murina puta</i>	<a href="#">GQ168901</a>	<i>Murina pluvialis</i>	<a href="#">JQ044689</a>
<i>Murina aenea</i>	<a href="#">GQ168906</a>	<i>Murina peninsularis</i>	<a href="#">GQ168911</a>
<i>Murina huttoni rubella</i>	<a href="#">KU521385</a>	<i>Myotis siligorensis</i>	<a href="#">FJ215679</a>
<i>Murina jaintiana</i>	<a href="#">JQ044690</a>	<i>Kerivoula furva</i>	<a href="#">MH208497</a>

Based on the reconstructed phylogenetic tree, the Chinese mainland *Harpiola* formed a monophyletic group with *H. isodon* (Fig. 3), which indicates a close relationship between them. However, the two taxa showed genetic distances of 4.8–5.1% between each other,

which exceeded the difference between *Murina recondita* and *Murina gracilis* (3.7%) and was not much lower than those between *Murina suilla* and *Murina florium* (7.7%), *Murina gracilis* and *Murina eleryi* (7.7%) and *Murina puta* and *Murina huttoni rubella* (7.2%) (Suppl. material 1). Thus, there are substantial sequence differences in the studied uniparentally inherited gene between Taiwanese and mainland *Harpiole* bats, but the taxonomic interpretation of these differences would be premature and additional material and nuclear genes should be included in subsequent analyses.




Figure 3. [doi](#)

Maximum Likelihood phylogenetic reconstruction of 25 Vespertilionidae samples using an 1140 bp alignment of the mitochondrial *cyt b* gene. Bootstrap (BS) values are indicated adjacent to nodes (nodes with BS < 0.50 are not labelled).

With a synthetic consideration of morphological and molecular evidence, we refer to the specimens from the Guanyin Mountains, Yuanyang, Yunnan as the first records of *H. isodon* on the Chinese mainland.

## Ecological notes

The specimens were captured at two mountain sites (23.03N, 102.95E and 22.99N, 102.99E) with mist nets. Their locations and the distribution map of *Harpiole isodon* are shown in Fig. 4. The habitat is a mid-mountain evergreen broad-leaved forest at elevations of 2,381 m and 2,463 m, respectively. The canopy is well closed and the forest is rich in shrubs and has small streams. There are no caves in this area, but there are many large trees with tree holes. We speculate that this insect-eating bat mainly roosts in tree cavities or under the dense canopy during the daytime.




Figure 4. [doi](#)

Distribution map of *Harpiola isodon*. Coordinate information is derived from Kuo et al. (2006), Kruskop et al. (2006) and the Global Biodiversity Information Facility (Anonymous 2024).

## Acknowledgements

We are very grateful to the Conservation Area Management Committee of Guanyin Mountains Provincial Nature Reserve, Prof. Rui-Chang Quan and Prof. Yun-Hong Tan (Xishuangbanna Tropical Botanical Garden, CAS) for their help in our fieldwork, also thanks to Prof. Zhen Liu (Kunming Institute of Zoology, CAS) for his help in our paper writing. The work was supported by the survey of Chiroptera species diversity and distribution in northwest and southwest of China (No: 2021FY100302) and the Project of Yuanyang Guanyin Mountains Provincial Nature Reserve Integrative Scientific Expedition (E2HX105B). The work of G. Csorba was supported by the National Research, Development and Innovation Fund of Hungary (NKFIH FK137778).

## References

- Anonymous (2024) GBIF Occurrence Download. <https://doi.org/10.15468/dl.pkm4hw>. Accessed on: 2024-3-01.
- Bates PJ, Harrison DL (1997) Bats of the Indian subcontinent. Harrison Zoological Museum Publication, Sevenoaks [ISBN 0951731319]
- Bhattacharyya TP (2002) Taxonomic status of the genus *Harpiola* Thomas, 1915 (Mammalia: Chiroptera: Vespertilionidae), with a report of the occurrence of *Harpiola grisea* (Peters, 1872) in Mizoram, India. Proceedings of the Zoological Society 55 (1): 73-76.

- Bickham JW, Wood CC, Patton JC (1995) Biogeographic implications of cytochrome b sequences and allozymes in sockeye (*Oncorhynchus nerka*). *Journal of Heredity* 86 (2): 140-144. <https://doi.org/10.1093/oxfordjournals.jhered.a111544>
- Corbet GB, Hill JE (1992) *The mammals of the Indomalayan region*. Natural History Museum and Oxford University Press, Oxford, 488 pp. [ISBN 0198546939]
- Dobson GE (1878) *Catalogue of the Chiroptera in the collection of the British Museum*. Cornell University Library, London, 567 pp. <https://doi.org/10.5962/bhl.title.8322>
- Ellerman JR, Morrison-Scott TC (1951) *Checklist of Palaearctic and Indian mammals 1758 to 1946*. Trustees of the British Museum (Natural History), London, 810 pp. <https://doi.org/10.1126/science.115.2990.431>
- Kalyanamoorthy S, Minh BQ, Wong TK, Haeseler A, Jermin LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* 14 (6): 587-589. <https://doi.org/10.1038/nmeth.4285>
- Koopman KF (1994) *Chiroptera: Systematics*. In: Niethammer J, Schliemann H, Starck D (Eds) *Handbook of Zoology*. Vol. 8. Walter de Gruyter, Berlin, 217 pp. [ISBN 3110140810].
- Kruskop SV, Kalyakin MV, Abramov AV (2006) First record of *Harpiola* (Chiroptera, Vespertilionidae) from Vietnam. *Russian Journal of Theriology* 5 (1): 13-16. <https://doi.org/10.15298/rusjtheriol.05.1.02>
- Kuo HC, Fang YP, Csorba G, Lee LL (2006) The definition of *Harpiola* (Vespertilionidae: Murininae) and the description of a new species from Taiwan. *Acta Chiropterologica* 8 (1): 11-19. [https://doi.org/10.3161/1733-5329\(2006\)8\[11:tдохvm\]2.0.co;2](https://doi.org/10.3161/1733-5329(2006)8[11:tдохvm]2.0.co;2)
- Maeda K (1980) Review on the classification of Little tube-nosed bats, *Murina aurata*, group. *Mammalia* 44 (4): 531-551. <https://doi.org/10.1515/mamm.1980.44.4.531>
- Ognev S (1928) *Mammals of Eastern Europe and Northern Asia*. Vol. 1. Insectivora and Chiroptera, Glavnauka, Moscow, 631 pp. <https://doi.org/10.5962/bhl.title.46316>
- Peters W (1880) Mittheilung über die von Hrn. In: Hilgendorf F (Ed.) *Japan gesammelten Chiropteren*. *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften*, 1880: 23-25 pp. URL: <https://biostor.org/reference/129543>
- Simmons N (2005) Order Chiroptera. 312–529pp. In: Wilson DE, Reeder DM (Eds) *Mammal species of the World: a taxonomic and geographic reference*. 3rd edition. The Johns Hopkins University Press, Baltimore, 2142 pp. [ISBN 0801882214].
- State Council Decree (1992) *Wildlife protective enforcement regulation of The People's Republic of China*. The Ministry of Forestry of PR China, s. n. [unknown pagination, in Chinese].
- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* 38 (7): 3022-3027. <https://doi.org/10.1093/molbev/msab120>
- Tate GHH (1941) Results of the Archbold expeditions no. 40. Notes on Vespertilionid bats of the subfamilies Miniopterinae, Murininae, Kerivoulinae and Nyctophilinae. *Bulletin of the American Museum of Natural History* 78: 567-597.
- Thomas O (1915) A special genus for the Himalayanbat known as *Murina grisea*. *Annals and Magazines of Natural History* 16 (8): 309-310. <https://doi.org/10.1080/00222931508693717>
- Wallin L (1969) The Japanese bat fauna. *Zoologiska Bidrag fran Uppsala* 37: 223-440. URL: <http://api.semanticscholar.org/CorpusID:87152921>

- Yoshiyuki M (1989) A systematic study of the Japanese Chiroptera. National Science Museum, Tokyo. URL: <https://api.semanticscholar.org/CorpusID:82665303>
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. Molecular Ecology Resources 20 (1): 348-355. <https://doi.org/10.1111/1755-0998.13096>.

## Supplementary material

### Suppl. material 1: Uncorrected pairwise genetic *P*-distance

**Authors:** Xin Mou

**Data type:** pairwise genetic *P*-distance

**Brief description:** Uncorrected pairwise genetic *P*-distance (%) amongst the species on 1140 bp of mitochondrial Cyt b.

[Download file](#) (22.96 kb)