



New records of two endemic troglobitic and threatened arachnids (Amblypygi and Opiliones) from limestone caves of Minas Gerais state, southeast Brazil

Bruno Gabriel O do Monte[‡], Jonas Eduardo Gallão^{‡,§}, Diego M von Schimonsky^{‡,§}, Maria Elina Bichuette[‡]

[‡] Laboratório de Estudos Subterrâneos, Universidade Federal de São Carlos, São Carlos, São Paulo, Brazil
[§] Programa de Pós-Graduação em Biologia Comparada-FFCLRP-USP, Ribeirão Preto, São Paulo, Brazil

Corresponding author: Bruno Gabriel O do Monte (bruno_gom@hotmail.com)

Academic editor: Adriano Kury

Received: 08 May 2015 | Accepted: 06 Nov 2015 | Published: 10 Nov 2015

Citation: do Monte B, Gallão J, von Schimonsky D, Bichuette M (2015) New records of two endemic troglobitic and threatened arachnids (Amblypygi and Opiliones) from limestone caves of Minas Gerais state, southeast Brazil. Biodiversity Data Journal 3: e5260. doi: [10.3897/BDJ.3.e5260](https://doi.org/10.3897/BDJ.3.e5260)

Abstract

Background

The endemic and threatened troglobites (organisms restricted to caves) *Charinus eleonora* (Amblypygi) and *landumoema uai* (Opiliones), both from Olhos d'Água cave, located at Peruaçu Caves National Park (southeast Brazil), have their distribution expanded for a new locality inside of the National Park (Lapa do Cipó cave), extending their distribution at least in 9.5km².

New information

This new data suggest that these arachnids can be in a differentiation process and/or there are several possibilities of dispersion in the karst of Peruaçu. Indeed, a revision of their

categorical status at IUCN Red List is necessary. We herein report a new distribution range (Lapa do Cipó cave) of the troglobitic species *I. uai* and *C. eleonora*, which are, to date, known to occur in the Olhos d'Água cave, located at the Peruaçu Caves National Park (PCNP).

Keywords

Charinus, *landumoema*, new distribution, Olhos d'Água cave, Lapa do Cipó cave

Introduction

In Brazil there are thirteen species of troglobitic (obligatory cave-dweller) opilionids, belonging to two families: Gonyleptidae Sundevall, 1833 (*Pachylospeleus strinatii* Šilhavý, 1974; *landumoema uai* Pinto-da-Rocha, 1996; *Giupponia chagasi* Pérez & Kury, 2002; *Discocyrtus pedrosoi* Kury, 2008; *Eusarcus elinae* Kury, 2008; *Spinopilar moria* Kury & Pérez-González, 2008; *landumoema setimapocu*, Hara & Pinto-da-Rocha, 2008; *landumoema* sp. n.; *Eusarcus* sp. n. 2; *Eusarcus* sp. n. 3 and two undescribed Pachylinae) and Escadabiidae Kury and Pérez, 2003 (*Spaeleolepts spaeleus* Soares, 1966) (Hara and Pinto-da-Rocha 2008). Out of these eight troglobitic opilionid species, seven Gonyleptidae are from Bahia, Minas Gerais, and São Paulo states and only one Escadabiidae species is from Minas Gerais State (Trajano and Bichuette 2010).

The troglobitic belonging to the genus *landumoema* Pinto-da-Rocha, 1996 consists of three species, *landumoema uai* Pinto-da-Rocha, 1996, *landumoema setimapocu* Hara and Pinto-da-Rocha, 2008 and *landumoema* sp. n. All of the species occur in the center-north of Minas Gerais State and each one of them is recorded only from their type-locality. For example, *I. uai* is found at the Olhos d'Água cave (Pinto-da-Rocha 1996).

The genus *Charinus* Simon, 1982 is the most diverse in Order Amblypygi and the Family Charinidae Quintero, 1986. This genus comprises 17 species in South America and 11 species in Brazil (Harvey 2003, Vasconcelos et al. 2014). Out of these, two are troglobitic species: *C. troglobius* Baptista and Giupponi, 2002 and *C. eleonora* Baptista and Giupponi, 2003. *C. eleonora* like *I. uai* is also known only from its type-locality, the Olhos d'Água cave at Itacarambi municipality in the north of Minas Gerais State.

Out of these, two are troglobitic species: *C. troglobius* Baptista and Giupponi, 2002 and *C. eleonora* Baptista and Giupponi, 2003.

However, endemism is not necessarily a characteristic of such obligatory cave-dweller arachnids in Brazil, since there are records of troglobitic species occurring in two or more caves in Brazil. For example, the opilionid *Discocyrtus pedrosoi* Kury, 2008 from Chapada Diamantina, central region of Bahia State, occurs in seven caves (J.E. Gallão, pers. obs.) and the opilionid *Giupponia chagasi* Pérez and Kury, 2002, as well as the amblypygid *C.*

troglobius, both from Serra do Ramalho karstic area, south of Bahia state, occur in two caves (Baptista and Giupponi 2002).

The ecological importance and fragility of the troglobitic species, *I. uai* and *C. eleonora*, are recognized by the International Union for Conservation of Nature (IUCN). Both these species are included in the IUCN Red List as Critically Endangered (CR), the higher risk category, highlighting their extremely vulnerability (IUCN 2014). However, neither a management plan nor an access control has been implemented for the caves inside a Peruaçu Caves National Park (PCNP), putting those species in risk.

We herein report a new distribution range (Lapa do Cipó cave) of the troglobitic species *I. uai* and *C. eleonora*, which are, to date, known to occur in the Olhos d'Água cave, located at the Peruaçu Caves National Park (PCNP).

Materials and methods

Study area

The Peruaçu Caves National Park (PCNP) located at the Peruaçu Basin of the São Francisco River Basin, is compound by large rocky outcrops with predominance of limestone rocks of Bambuí karst area (Auler et al. 2001). The region is located between the transition of Cerrado (savannah-like vegetation) and Caatinga (semiarid vegetation) phytophysionomies (Ab'Saber 1977); the dry season in this region occurs between April and November and the average temperature is 24°C (Nimer 1979).

The Olhos d'Água cave is the largest cave of Minas Gerais State, with approximately nine kilometers of horizontal projection (Auler et al. 2001), occurring at the PCNP. The cave is considered a spot of high biodiversity in Brazil, with more than seven troglobitic species (Deharveng and Bedos 2005, Trajano and Bichuette 2010).

We conducted fieldtrips to PCNP in June and August of 2014 for sampling caves (Fig. 1). The Lapa do Cipó cave (S 15.05611, W 44.18444) (Fig. 2) is located 6.5 km northwest of the Olhos d'Água cave entrance. Both caves are in different small drainages (part of Peruaçu basin – see Fig. 1) which can be an isolation factor for aquatic fauna, but not necessarily for terrestrial cave invertebrates, however hidrogeological studies are still needed.

To recognize the minimal occurrence area of the species, we did a triangulation with the three points of caves (see Fig. 1).




Figure 1.

Type-locality (Olhos d'Água cave) and new records (Lapa do Cipó cave) for the troglobitic *landumoema uai* genus *Charinus* is characterized by the following characteristics and *Charinus eleonora*. The soil level represents the relative altitudes in the area and the drainages are in the lowest level. Olhos d'Água cave resurgence is the main entrance for this cave. See the two separated drainages for both caves.



Figure 2.

Lapa do Cipó cave entrance; ca. 20 meters high. Both people in the bottom right are for scale.

Collection and identification

We employed the direct qualitative search and hand collecting sampling method targeting walls, under block rocks, organic matter, and unconsolidated substrate (wet and dry). The collected individuals were fixed in 70% ethanol.

Identification and diagnosis of species was conducted following the original description of taxa (Pinto-da-Rocha 1996, Baptista and Giupponi 2003). We compared individuals from Lapa do Cipó cave and those from the type-locality, Olhos d'Água cave. Additional reference material, deposited at Laboratório de Estudos Subterrâneos (LES) collection from Universidade Federal de São Carlos (UFSCar), São Carlos municipality from São Paulo state, were also used for the comparison.

In addition, for body measurements, we used the classical morphometric data specific for each group to confirm the identification. The measurements for opilionids and amblypygids were conducted as described by Acosta et al. (2007) and Quintero-Jr. (1981), respectively. We used a digital caliper with 0.01 mm accuracy for the measurements. In total, we measured 13 specimens of *I. uai* (seven from Olhos d'Água cave and six from Lapa do Cipó cave) and 10 specimens of *C. eleonora* (seven from Olhos d'Água cave and three from Lapa do Cipó cave).

Images were taken using a Leica DFC 295 camera attached to a Leica M205C stereomicroscope with a PlanApo (1.0) objective. The figures were produced using multiple frames of LAS software (Leica Application Suite v3.7).

Taxonomy

The genus *landumoema* is characterized by a single erect spine on eye mound, areas of dorsal scutum unarmed, and presence of mesal-subapical setae on the pedipalpal femur (Fig. 3a, b, c). The two species of this genus, *I. setimapocu* and *I. uai*, differ in terms of the curvature on femur IV as observed on *I. uai* and the direction of dorso-apical apophyses on male coxae IV; the latter is directed laterally backwards, away from the body in *I. uai*, while it is closer in *I. setimapocu*. In addition, in *I. uai* males, the tibial pedipalpal setae features lili conformation (Fig. 3a) (Pinto-da-Rocha 1996, Hara and Pinto-da-Rocha 2008).

The genus *Charinus* is characterized by the following characteristics: pedipalpal basitarsus with two long dorsal spines and one ventral; pedipalpal tibia expanded dorsally, with a spine and a setiferous tubercle distally in relation to its longest dorsal spine; trochanter with a well-developed ventral protuberance, with setiferous tubercles anteriorly projected (Armas and Pérez-González 2001). *Charinus troglobius* is characterized by an anterior depression on the carapace, in place of the absent median eye tubercle (Baptista and Giupponi 2002), whereas *C. eleonora* (Fig. 4a, b, c) is characterized by an indistinct median eye tubercle (Fig. 4a, b, c) usually with only two very small eye spots, or without eye spots in rare cases (Baptista and Giupponi 2003).



Figure 3.

landumoema uai, male from Lapa do Cipó cave.

a: Dorsal view of pedipalpal tibia-tarsus ectal and mesal setae lili

b: Dorsal view of habitus, showing coxa IV tuberculate laterally, with robust apical external apophysis and curved at 1/3 distal (on male) and femur IV curved laterally and dorsally at 1/3 from base (on male)

c: Lateral view of habitus, showing eye mound with erect high spine, with acuminate apex pointing slightly backwards.

Taxon treatments

Charinus eleonora Baptista & Giupponi 2003

Materials

- a. acceptedNameUsageID: *Charinus eleonora*; taxonID: *Charinus eleonora*; locationID: Laboratório de Estudos Subterrâneos / Universidade Federal de São Carlos; higherGeographyID: São Carlos, São Paulo State; higherGeography: Brazil; continent: South America; verbatimCoordinates: 15 06 49.0S 44 10 10.0W; geodeticDatum: WGS84; eventID: July 26, 2010; individualCount: 5; occurrenceDetails: Olhos d'Água cave; geologicalContextID: Bambuí geomorphological unit, limestone from Peruaçu karst area, Medium São Francisco basin, southeast Brazil; identificationID: LES 0400; LES03217;; identifiedBy: Bruno Gabriel O. do Monte; identificationReferences: Baptista & Giupponi 2003

- b. acceptedNameUsageID: *Charinus eleonorae*; taxonID: *Charinus eleonorae*; locationID: Laboratório de Estudos Subterrâneos / Universidade Federal de São Carlos; higherGeographyID: São Carlos, São Paulo State; higherGeography: Brazil; continent: South America; verbatimCoordinates: 15 06 49.0S 44 10 10.0W; verbatimSRS: WGS84; eventID: July 23-24, 2012; individualCount: 2; occurrenceDetails: Olhos d'Água cave; geologicalContextID: Bambuí geomorphological unit, limestone from Peruaçu karst area, Medium São Francisco basin, southeast Brazil; identificationID: LES03232; LES05873;; identifiedBy: Bruno Gabriel O. do Monte; identificationReferences: Baptista & Giupponi 2003
- c. acceptedNameUsageID: *Charinus eleonorae*; taxonID: *Charinus eleonorae*; locationID: Laboratório de Estudos Subterrâneos / Universidade Federal de São Carlos; higherGeographyID: São Carlos, São Paulo State; higherGeography: Brazil; continent: South America; verbatimCoordinates: 15 03 22.0S 44 11 04.0W; verbatimSRS: WGS84; eventID: June 05, 2014; individualCount: 3; occurrenceDetails: Lapa do Cipó cave; geologicalContextID: Bambuí geomorphological unit, limestone from Peruaçu karst area, Medium São Francisco basin, southeast Brazil; identificationID: LES05868; identifiedBy: Bruno Gabriel O. do Monte; identificationReferences: Baptista & Giupponi 2003




Figure 4.

Charinus eleonorae, female from Lapa do Cipó cave.

a: Carapace, dorsal view, showing mid-portion of carapace without median eyes

b: Habitus, dorsal view

c: Lateral view of right pedipalp showing spines.

Conservation

According to 2014 IUCN revision, this species is CR (Critically Endangered) category.

Taxon discussion

Expansion of occurrence of troglobitic species previously known for only a single cave.

landumoema uai Pinto-da-Rocha 1996

Materials

- a. acceptedNameUsageID: *landumoema uai*; taxonID: *landumoema uai*; locationID: Laboratório de Estudos Subterrâneos / Universidade Federal de São Carlos; higherGeographyID: São Carlos, São Paulo State; higherGeography: Brazil; continent: South America; verbatimCoordinates: 15 06 49.0S 44 10 10.0W; verbatimSRS: WGS84; eventID: July 26, 2010; individualCount: 7; occurrenceDetails: Olhos d'Água cave; geologicalContextID: Bambuí geomorphological unit, limestone from Peruaçu karst area, Medium São Francisco basin, southeast Brazil; identificationID: LES03214, LES03233; identifiedBy: Bruno Gabriel O. do Monte
- b. acceptedNameUsageID: *landumoema uai*; taxonID: *landumoema uai*; locationID: Laboratório de Estudos Subterrâneos / Universidade Federal de São Carlos; higherGeographyID: São Carlos, São Paulo State; higherGeography: Brazil; continent: South America; verbatimCoordinates: 15 03 22.0S 44 11 04.0W; verbatimSRS: WGS84; eventID: June 05, 2014; individualCount: 2; sex: MALE; occurrenceDetails: Lapa do Cipó cave; geologicalContextID: Bambuí geomorphological unit, limestone from Peruaçu karst area, Medium São Francisco basin, southeast Brazil; identificationID: LES05869, LES05870; identifiedBy: Bruno Gabriel O. do Monte
- c. acceptedNameUsageID: *landumoema uai*; taxonID: *landumoema uai*; locationID: Laboratório de Estudos Subterrâneos / Universidade Federal de São Carlos; higherGeographyID: São Carlos, São Paulo State; higherGeography: Brazil; continent: South America; verbatimCoordinates: 15 03 22.0S 44 11 04.0W; verbatimSRS: WGS84; eventID: August 26, 2014; individualCount: 4; sex: MALE; occurrenceDetails: Lapa do Cipó cave; geologicalContextID: Bambuí geomorphological unit, limestone from Peruaçu karst area, Medium São Francisco basin, southeast Brazil; identificationID: LES05871, LES05872; identifiedBy: Bruno Gabriel O. do Monte

Conservation

According to 2014 IUCN revision, this species is CR (Critically Endangered) category.

Taxon discussion

Expansion of occurrence of troglobitic species previously known for only a single cave.

Analysis

Individuals of *I. uai* and *C. eleonora*, recorded from Lapa do Cipó cave, are the first reported occurrence of these species from any other cave beyond their type-locality, Olhos d'Água cave. We observed that all individuals from both the species present the diagnostic characters for *I. uai* and *C. eleonora*, with no sharp differences in characters among the specimens from both localities. However, all the three *C. eleonora* specimens from Lapa do Cipó cave had only eyespots with the median eyes absent, which are reduced in the specimens from Olhos d'Água cave. Morphometric data of both the species from Lapa do Cipó cave are in agreement with the original description with subtle differences (Tables 1, 2).

Table 1.

Mean measurements values and standard error for the opilionid *landumoema uai* from Olhos d'Água and Lapa do Cipó caves. The fifth column brings the data of the original description – type-series (Pinto-da-Rocha 1996).

| | | Olhos d'Água n = 7 | Lapa do Cipó n = 6 | Type-series n = 9 |
|------------|------------|-----------------------|-----------------------|----------------------|
| Pedipalpus | Trochanter | 0.48(0.01) | 0.52(0.02) | 0.54 |
| | Femur | 2.03(0.05) | 1.92(0.08) | 1.96 |
| | Patella | 1.01(0.02) | 1(0.04) | 1.10 |
| | Tibia | 1.46(0.04) | 1.33(0.04) | 1.40 |
| | Tarsus | 1.01(0.03) | 1(0.03) | 1.10 |
| | Total | 6(0.11) | 5.78(0.18) | 6.10 |
| Leg I | Trochanter | 0.54(0.02) | 0.528(0.03) | 0.46 |
| | Femur | 4.57(0.06) | 4.09(0.18) | 4.40 |
| | Patella | 1.08(0.02) | 0.989(0.03) | 1.12 |
| | Tibia | 3.51(0.07) | 3.17(0.08) | 3.37 |
| | Metatarsus | 5.83(0.03) | 5.26(0.2) | 5.69 |
| | Tarsus | 2.48(0.24) | 3.72(1.1) | 2.75 |
| | Total | 18.03(0.26) | 17.76(1.1) | 17.79 |
| Leg II | Trochanter | 0.60(0.02) | 0.56(0.03) | 0.60 |
| | Femur | 9.24(0.07) | 8.12(0.3) | 8.64 |
| | Patella | 1.61(0.01) | 1.27(0.07) | 1.44 |
| | Tibia | 8.05(0.02) | 7.26(0.2) | 7.37 |
| | Metatarsus | 9.44(0.08) | 8.72(0.22) | 8.88 |
| | Tarsus | 10.16(0.18) | 9.75(0.14) | 9.84 |
| | Total | 37.31(1.3) | 32.63(3.56) | 36.77 |
| Leg III | Trochanter | 0.61(0.03) | 0.58(0.03) | 0.62 |

| | | | | |
|---------------------|------------|-------------|-------------|-------|
| | Femur | 6.21(0.06) | 5.6(0.18) | 5.70 |
| | Patella | 1.22(0.02) | 1.13(0.04) | 1.28 |
| | Tibia | 3.88(0.05) | 3.56(0.11) | 3.56 |
| | Metatarsus | 6.55(0.08) | 6.03(0.21) | 6.31 |
| | Tarsus | 3.10(0.05) | 2.98(0.61) | 2.87 |
| | Total | 21.60(0.17) | 18.24(1.57) | 20.34 |
| Leg IV | Trochanter | 1.05(0.03) | 0.99(0.16) | 1.15 |
| | Femur | 8.17(0.06) | 7.6(0.26) | 8.10 |
| | Patella | 1.42(0.07) | 1.42(0.09) | 1.69 |
| | Tibia | 5.94(0.07) | 5.37(0.21) | 5.56 |
| | Metatarsus | 8.85(0.19) | 8.25(0.28) | 8.56 |
| | Tarsus | 3.84(0.02) | 3.75(0.09) | 3.59 |
| | Total | 29.29(0.4) | 25.40(2.03) | 28.65 |
| Dorsal scute length | | 3.86(0.07) | 3.7(0.15) | 4.04 |
| Prosoma length | | 4.40(0.09) | 3.55(0.08) | 1.52 |
| Prosoma width | | 3.77(0.29) | 3.72(0.24) | 1.88 |
| Opisthosoma width | | 2.44(0.04) | 2.13(0.2) | 3.16 |

Table 2.

Mean measurements values and standard error for the amblypigid *Charinus eleonorae* from Olhos d'Água and Lapa do Cipó caves. The fourth column brings the data of the original description – type-series (Baptista and Giupponi 2003).

| | Olhos d' Água n = 7 | Lapa do Cipó n = 3 | Type-series n = 17 |
|---------------------|------------------------|-----------------------|-----------------------|
| Femur | 2.89 (0.36) | 3.09(0.01) | 4.2(3.4-5.6) |
| Tibia | 3.47(0.43) | 3.86(0.22) | 4.0(3.2-5.4) |
| Basitarsus | 1.47(0.17) | 1.76(0.11) | 1.8 (1.6-2.2) |
| Distitarsus | 1.15(0.07) | 1.35(0.06) | 1.2(1.1-1.4) |
| Tarsal claw | 0.73(0.03) | 0.85(0.01) | 0.9(0.8-1.1) |
| Total | 9.74(1.02) | 10.93(0.36) | |
| Cephalotorax length | 2.97(0.22) | 3.23(0.32) | 3.5(3.1-4.0) |
| Cephalotorax width | 3.15(0.25) | 3.52(0.19) | 4.3(3.9-5.1) |
| Abdomen length | 4.60(0.43) | 6.17(0.24) | 5.0(4.2-5.6) |
| Body length | 7.41(0.58) | 9.12(0.37) | 7.8(7.0-9.1) |

Discussion

The Olhos d'Água cave is the largest cave in the Minas Gerais State with a significant number of troglobitic species and restricted range distribution (Deharveng and Bedos 2005, Trajano and Bichuette 2010). Nevertheless, there is no information on other entrances or accesses except the known resurgence. Therefore, we considered that the hypogean habitats are isolated. However, the Olhos d'Água cave upstream entrance (sinkhole), known as Água d'Olhos (Piló 1989) may represent an alternative passageway to the cave fauna. To date, there are no geological studies that indicate a possible communication with the other caves in the region.

The occurrence of *I. uai* and *C. eleonora*e in Lapa do Cipó cave (distant 6.5 km northwest from Olhos d'Água cave) either indicates the existence of a complex system of subterranean microspaces (such as cracks and fissures), interconnecting both caves in the karst of Peruaçu; or the past existence of a whole unique system, reaching these two caves. In the first case, the dispersion can occur through the voids connected through cracks and fissures, typical of the MSS habitat (mesovoid shallow substratum, sensu Juberthie 2000). In the latter case, as defined by Ford and Williams 2007, which outlined that cave systems are integrated habitats linking output and input points of dissolution, which possibilities the flow of clastic sediments and, particularly, the fauna. These two hypothesis are not mutually exclusive.

In terms of median eyes, we observed that the *C. eleonora*e populations from Lapa do Cipó cave showed a higher frequency of individuals without them comparing to the observed for Olhos d'Água cave population. The absence of median eyes is considered a rare condition by (Baptista and Giupponi 2003). Our data (frequent absence of median eyes) strongly suggests that this variation must be more common than the authors considered and the description and proposed key differentiating *C. eleonora*e and *C. troglobius* must be reviewed. However, such variability has not been recorded for the highly troglomorphic species, *C. troglobius*, from Serra do Ramalho (Bahia), and therefore, in this case, the absence of median eyes is a robust diagnostic feature (Baptista and Giupponi 2003). Furthermore, the occurrence of such variability in *C. eleonora*e is a indicative of or relaxed pressure or even neutralism processes, testable hypothesis through population genetic studies.

For conservation purposes, the data presented herein, such as the occurrence area of 9.5km² made by triangulation of A, B and C points (Fig. 1), suggests a revision of the category status of these two species in the IUCN list (from critically endangered - CR to Endangered - EN). Finally, an implementation of the management plan for caves in the Peruaçu Caves National Park (PCNP) is urgent, as well as the regulation of cave tourism in the region.

Acknowledgements

We are grateful to Vandeir B. de Jesus (“Branco”) for help in the fieldtrips; to Peruaçu Caves National Park administrators for support. We also thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for financial support and grants to BGOM and DMS (doctoral scholarships); to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for grants to JEG (doctor scholarship - CNPq142276/2013-8) and MEB (fellowship); to Dr. Angélica M.P.M. Dias and Instituto Nacional de Ciência e Tecnologia de Hymenoptera Parasitóides da Região Sudeste Brasileira (INCT Hympar Sudeste) for providing us the use of the stereomicroscope Leica DFC 295; to L.B.R. Fernandes for image captures; to M.P. Bolfarini for photography of Lapa do Cipó cave; to the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) for collection permit (28992-4) and to the Programa de Pós-graduação em Ecologia e Recursos Naturais da Universidade Federal de São Carlos (PPGERN/ UFSCar) for the infrastructure to develop this work. A native speaker revised the work.

Author contributions

BGOM, JEG and DMvS collected, identified and contributed for writing.

MEB help in the identification, called attention to the new distribution and contributed for writing.

DMvS confeccioned the map and formatted the figures.

References

- Ab'Saber AN (1977) Os domínios morfoclimáticos na América do Sul. *Geomorfologia* 52: 1-21. [In Portuguese].
- Acosta LE, Pérez-Gonzalez A, Tourinho AL (2007) Methods and Techniques of Study: Methods for taxonomic study. In: Machado G, Pinto-da-Rocha R, Giribet G (Eds) *Harvestmen, the biology of Opiliones*. Harvard University Press, Cambridge, Massachusetts and London, 597 pp.
- Armas LF, Pérez-González A (2001) Los ambliopígididos de República Dominicana (Arachnida: Amblypygi). *Revista Ibérica de Aracnología* 3: 47-66. [In Spanish].
- Auler A, Rubbioli E, Brandi R (2001) As grandes cavernas do Brasil. 1, 1. Grupo Bambuí de Pesquisas Espeleológicas, Belo Horizonte, 228 pp. [In Portuguese].
- Baptista RLC, Giupponi APL (2002) A new troglomorphic *Charinus* from Brazil. *Revista Ibérica de Aracnología* 6: 105-110.
- Baptista RLC, Giupponi APL (2003) A new troglomorphic *Charinus* from Minas Gerais State, Brazil (Arachnida: Amblypygi: Charinidae). *Revista Ibérica de Aracnología* 7: 79-84.

- Deharveng L, Bedos A (2005) Diversity Patterns in the Tropics. Encyclopedia of Caves. URL: <http://dx.doi.org/10.1016/b978-0-12-383832-2.00032-3> DOI: [10.1016/b978-0-12-383832-2.00032-3](https://doi.org/10.1016/b978-0-12-383832-2.00032-3)
- Ford D, Williams P (2007) Karst hydrogeology and geomorphology. Wiley, Chichester, 562 pp. [ISBN 97804700849655] DOI: [10.1002/9781118684986](https://doi.org/10.1002/9781118684986)
- Hara MR, Pinto-da-Rocha R (2008) A new species of Brazilian troglobitic harvestman of the genus *landumoema* (Opiliones: Gonyleptidae). Zootaxa 1744: 50-58. DOI: [10.11646/zootaxa.1744.1.5](https://doi.org/10.11646/zootaxa.1744.1.5)
- Harvey MS (2003) Catalogue of the smaller arachnid orders of the world, Amblypygi, Uropygi, Schizomida, Palpigradi, Ricinulei and Solifugae . CSIRO, COLLINGWOOD, 385 pp.
- IUCN (2014) Guidelines for Using the IUCN Red List Categories and Criteria. <http://ir.iucnredlist.org/documents/RedListGuidelines.pdf>. Accession date: 2015 1 10.
- Juberthie C (2000) The diversity of the karstic and pseudo karstic hypogean habitats in the world. In: WILKENS H, CULVER DC, HUMPHREYS WF (Eds) Ecosystems of the World: Subterranean Ecosystems. Elsevier, Amsterdam, 791 pp. [ISBN 0444822992].
- Nimer E (1979) Climatologia do Brasil. SUPREN, Rio de Janeiro, 421 pp. [In Portuguese].
- Piló LB (1989) A morfologia cárstica do baixo curso do rio Peruaçu, Januária-Itacarambi, MG (Graduate Monography). UFMG, Belo Horizonte, 80 pp. [In Portuguese].
- Pinto-da-Rocha R (1996) *landumoema uai*, a new genus and species of troglobitic harvestman from Brazil (Arachnida, Opiliones, Gonyleptidae). Revista Brasileira de Zoologia 13 (4): 843-848. DOI: [10.1590/s0101-81751996000400005](https://doi.org/10.1590/s0101-81751996000400005)
- Quintero-Jr. D (1981) The amblypygid *Phrynus* in the Americas (Amblypygi, Phrynidae). Journal of Arachnology 9: 117-166.
- Trajano E, Bichuette ME (2010) Diversity of Brazilian subterranean invertebrates with a list of troglomorphic taxa. Subterranean Biology 7: 1-16.
- Vasconcelos AC, Giupponi APL, Ferreira RL (2014) A new species of *Charinus* from Minas Gerais State, Brazil, with comments on its sexual dimorphism (Arachnida: Amblypygi: Charinidae). Journal of Arachnology 42 (2): 155-162. DOI: [10.1636/h14-01.1](https://doi.org/10.1636/h14-01.1)