



Taxonomic Paper

Diversity of the longhorned beetles (Coleoptera: Cerambycidae) from Cuatro Ciénegas Basin, Coahuila, Mexico

Oscar Pérez-Flores[‡], Víctor H. Toledo-Hernández[§]

[‡] Colección Nacional de Insectos, Instituto de Biología, Universidad Nacional Autónoma de México, Delegación Coyoacán, Mexico City, Mexico

[§] Centro de Investigación en Biodiversidad y Conservación (CIByC), Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos, Mexico

Corresponding author: Oscar Pérez-Flores (oscar_skopt@ciencias.unam.mx)

Academic editor: Jennifer C. Girón

Received: 19 May 2020 | Accepted: 09 Jul 2020 | Published: 16 Jul 2020

Citation: Pérez-Flores O, Toledo-Hernández VH (2020) Diversity of the longhorned beetles (Coleoptera: Cerambycidae) from Cuatro Ciénegas Basin, Coahuila, Mexico. Biodiversity Data Journal 8: e54495.
<https://doi.org/10.3897/BDJ.8.e54495>

Abstract

Background

Cerambycidae is one the most diverse families in the order Coleoptera with more than 37,000 species described in all continents. Cerambicyd beetles have a worldwide distribution from sea level to montane sites. In Mexico, more than 1,600 species have been recorded. Nevertheless, the diversity and distribution of this family in the Mexican deserts is poorly known.

New information

A first checklist of Cerambycidae from seven localities of Cuatro Ciénegas Basin is presented. This study is the result of sampling carried out between 2009 to 2013. Some material from other collections is also included. The species list includes four subfamilies, 13 tribes, 32 genera and 37 species, from which 13 are new records for the state of

Coahuila and three species represent new records for Mexico. These results highlight the paucity of knowledge of insects in the deserts of Mexico.

Keywords

Cerambycid beetles, Chihuahuan desert, distribution, North America.

Introduction

Cerambycidae comprise one of the largest families of Coleoptera, represented by more than 37,000 species, described in more than 5,300 genera from eight subfamilies (Monné et al. 2017, Tavakilian and Chevillotte 2020). Cerambycid beetles are distributed worldwide wherever their hosts plants are found from sea level to montane sites as high as 4,200 m elevation (Bezark et al. 2013). Due to its ecological, biological and evolutionary importance, this family has been studied extensively. The diversity of the Cerambycidae in Mexico was summarised by Noguera (2014). However, knowledge of the diversity and distribution of the family Cerambycidae in the Mexican deserts is poor, with just some isolated records (Bezark 2020).


The Cuatro Ciénegas Basin (CCB) (Souza et al. 2012) is located at the south-eastern portion of the Chihuahuan Desert (Fig. 1). With more than 450,000 km², it is the largest desert in North America. This, along with its high biological diversity, makes it one of the most important deserts in the world (Morafka 1977). The CCB has a unique geological and biological history which dates from at least 40 mya in the Middle Eocene. At this point the region was isolated from the Atlantic Ocean by the rise of the Sierra Madre Oriental (Álvarez and Ojeda 2019). The subterranean water, which currently emerges, forms ponds and streams that distinguish this region from other surrounding desert areas. This, plus the isolation of the CCB, has resulted in a high degree of endemism in some taxonomic groups (Souza et al. 2012).

The fauna that occurs in the CCB is a mixture of species with Neotropical and Nearctic biogeographic origins or affinities. This important region of transition can represent the distributional limits of both groups (Souza et al. 2012, Álvarez and Ojeda 2019). This work is a result of a study of diverse taxonomic groups in the CCB (Álvarez and Ojeda 2019), undertaken for the purpose of providing a better understanding of its local diversity. There are no published records of cerambycids from this region. This study is the first approach that aims to contribute to the general knowledge of this group in the Mexican deserts.

Materials and methods

The dominant habitat in CCB is desertic shrub of the microphyte and rosetophyte variants, which include sotol, mesquite, secondary grassland and semi-aquatic vegetation associated with ponds. The climate is desertic dry (BW) (García 1964) with three well-defined seasons: rainy, cold dry and hot dry. All localities that were selected have

superficial water (except Río Cañón) and included all habitats present in CCB. Most habitats had some degree of perturbation due to different issues such as cattle grazing, agriculture and tourism. The localities were the following: (1) Churince ($26^{\circ}50'30.3''N$, $102^{\circ}08'9.9''W$), (2) Rancho PRONATURA ($26^{\circ}49'38.9''N$, $102^{\circ}01'24.4''W$), (3) Rancho Orozco ($26^{\circ}52'18.3''N$, $102^{\circ}05'16.8''W$), (4) Antiguos Mineros ($26^{\circ}46'57.7''N$, $102^{\circ}00'20.2''W$), (5) Poza Azul ($26^{\circ}55'21.2''N$, $102^{\circ}72'3.6''W$), (6) Río Cañón ($27^{\circ}00'33.7''N$, $102^{\circ}04'42.3''W$) and (7) San José del Anteojo ($26^{\circ}58'11.5''N$, $102^{\circ}07'26.4''W$) (Fig. 1). The distribution map was made using ArcGIS server (Bader 2005).



Fourteen field trips were performed during 2012 and 2013. Records from sporadic trips in 2009 and 2010 are also included. Sampling totalled 84 collecting days with at least three people collecting each day (09:00 to 16:00 h). Different sampling techniques (direct and indirect) were applied: by hand, sweep net and light traps. Cerambycid beetles were identified to species level with specialised literature (Chemsak 1991, Chemsak and Linsley

1965, Chemsak and Noguera 2005, Giesbert and Chemsak 1989, Giesbert and Chemsak 1993, Grzymala and Miller 2013, Linsley 1963, Linsley 1964, Linsley and Chemsak 1972, Noguera 2002, Santos-Silva 2007, Santos-Silva et al. 2016, Turnbow and Thomas 2002) and by comparison with identified specimens from the National Collection of Insects of the Biology Institute, UNAM. All specimens were deposited in the Insect Collection of the Zoological Museum “Alfonso L. Herrera” of the Faculty of Sciences, UNAM. The National Collection of Insects (CNIN) was revised and all the records belonging with CCB were included (Suppl. material 1).

List of Cerambycidae from Cuatro Ciénegas Basin. The taxonomy is based on Bezark (2020)

Subfamily Prioninae

Tribe Prionini

Derobrachus hovorei Santos-Silva, 2007

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénegas; locality: Rancho PRONATURA; verbatimCoordinates: 26°49'38.9"N, 102°01'24.4"W; samplingProtocol: Light trap; eventDate: 7-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB01; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénegas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: Light trap; eventDate: 20-Apr-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB02; basisOfRecord: Preserved Specimen

Host of: *Cercidium microphyllum* Rose and I. M. Johnston (Caesalpiniaceae), *Quercus* sp. (Fagaceae), *Prosopis juliflora* (Swartz) de Candolle (Mimosaceae), *Morus rubra* Linnaeus (Moraceae), *Citrus* sp. (Rutaceae), *Populus* sp. (Salicaceae), *Ulmus* sp. (Ulmaceae), *Vitis* sp. (Vitaceae). (Monné 2020)

Distribution: USA: Arizona, California, Nevada, New Mexico, Texas; MEXICO: Chihuahua, Coahuila, Durango, Nuevo León, Sonora, Tamaulipas.

Notes: MZ-FC

***Prionus (Prionus) poultoni* Lameere, 1912**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: La Becerra; samplingProtocol: By hand; eventDate: 15-Aug-14; habitat: Desertic shrub; recordedBy: Antonio Castro; collectionCode: CNIN_CCB01; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Chihuahua, Coahuila, Nuevo León, Oaxaca, Veracruz.

Notes: CNIN

Subfamily Lepturinae

Tribe Lepturini

***Meloemorpha aliena* (Bates, 1880)**

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'19.36", 102°09'04.38"W; samplingProtocol: By hand; eventDate: 21-Jul-15; habitat: Desertic shrub; recordedBy: Oscar Pérez; collectionCode: CNIN_CCB02; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'19.36"N, 102°09'04.38"W; samplingProtocol: By hand; eventDate: 21-Jul-15; habitat: Desertic shrub; recordedBy: Oscar Pérez; collectionCode: CNIN_CCB03; basisOfRecord: Preserved Specimen

Host of: *Pinus pseudostrobus* Lindley, *P. rufa* Endlicher (Pinaceae). (Monné 2020)

Distribution: MEXICO: Ciudad de México, Coahuila (new state record), Guerrero, Hidalgo, México, Michoacán, Morelos, Nuevo León, Veracruz; GUATEMALA.

Notes: CNIN

***Typocerus sinuatus* (Newman, 1841)**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: Sweeping; eventDate: 12-Apr-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB03; basisOfRecord: Preserved Specimen

Distribution: USA: Colorado, New Mexico; MEXICO: Coahuila (new country record).

Notes: MZ-FC

Subfamily Cerambycinae

Tribe Bothriospilini

Knulliana cincta subsp. *cincta* (Drury, 1773)

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: San José del Antejo; verbatimCoordinates: 26°58'11.5"N, 102°07'26.4"W; samplingProtocol: By hand; eventDate: 23-May-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB04; basisOfRecord: Preserved Specimen

Distribution: USA: Texas; MEXICO: Coahuila (new state record), Nuevo León, Sonora.

Notes: MZ-FC

Tribe Clytini

Megacyllene antennata (White, 1855)

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: By hand; eventDate: 24-Sep-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB05; basisOfRecord: Preserved Specimen

Host of: *Celtis reticulata* Torr. (Cannabaceae), *Prosopis juliflora* (Swartz) de Candolle, *Senegalia greggii* A. Gray (Mimosaceae). (Monné 2020)

Distribution: USA: California, Texas; MEXICO: Baja California Norte, Coahuila (new state record), Sinaloa, Sonora.

Notes: MZ-FC

Neoclytus mucronatus subsp. *mucronatus* (Fabricius, 1775)

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Poza Azul; samplingProtocol: By hand; eventDate: 17-Oct-16; habitat: Desertic shrub; recordedBy: Rodolfo Trejo; collectionCode: CNIN_CCB04; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Poza Azul; samplingProtocol: By hand; eventDate: 17-Oct-16; habitat: Desertic shrub; recordedBy: Rodolfo Trejo; collectionCode: CNIN_CCB05; basisOfRecord: Preserved Specimen

- c. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Poza Azul; samplingProtocol: By hand; eventDate: 17-Oct-16; habitat: Desertic shrub; recordedBy: Rodolfo Trejo; collectionCode: CNIN_CCB06; basisOfRecord: Preserved Specimen

Host of: *Acer rubrum* Linné (Aceraceae), *Celtis laevigata* Willdnow (Cannabaceae), *Juniperus virginiana* Linné (Cupressaceae), *Diospyros virginiana* Linné (Ebenaceae), *Quercus* sp. (Fagaceae), *Carya glabra* (Miller) Sweet, *C. illinoiensis* (Wangenheim) K. Koch, *C. ovata* (Miller) K. Koch, *C. tomentosa* Nuttall (Juglandaceae), *Pinus contorta* Douglas ex Loudon, *P. resinosa* Aiton (Pinaceae), *Ulmus* sp. (Ulmaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Coahuila (new country record).

Notes: CNIN

Placosternus erythropus (Chevrolat, 1835)

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: Sweeping; eventDate: 23-Oct-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB06; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: Sweeping; eventDate: 23-Oct-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB07; basisOfRecord: Preserved Specimen
- c. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Poza Azul; verbatimCoordinates: 26°55'21.2"N, 102°72'3.6"W; samplingProtocol: Sweeping; eventDate: 24-Aug-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB08; basisOfRecord: Preserved Specimen

Host of: *Baccharis neglecta* Nuttall (Asteraceae), *Acacia hindsii* Bentham, *Prosopis dulcis* Kunth, *P. juliflora* (Swartz) de Candolle (Mimosaceae), *Condalia globosa* I. M. Johnston (Rhamnaceae), *Crataegus pubescens* Steudel, *Malus sylvestris* Miller, *Prunus capulin* Zuccarini, *P. domestica* Linné, *P. persica* (Linné) Batsch, *Pyrus communis* Willdenow (Rosaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Ciudad de México, Coahuila, Durango, Guanajuato, Hidalgo, Jalisco, México, Morelos, Puebla, Sonora, Tabasco, Tamaulipas, Veracruz; GUATEMALA; COSTA RICA

Notes: MZ-FC

***Tanyochraethes hololeucus* (Bates, 1892)**

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Antiguos Mineros; verbatimCoordinates: 26°46'57.7"N, 102°00'20.2"W; samplingProtocol: By hand; eventDate: 22-Nov-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB09; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Antiguos Mineros; verbatimCoordinates: 26°46'57.7"N, 102°00'20.2"W; samplingProtocol: By hand; eventDate: 22-Nov-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB10; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Coahuila, Durango, Tamaulipas.

Notes: MZ-FC

Tribe Eburiini

***Eburia juanitae* Chemsak and Linsley, 1970**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Río Cañón; verbatimCoordinates: 27°00'33.7"N, 102°04'42.3"W; samplingProtocol: Sweeping; eventDate: 7-Jul-10; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB11; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Coahuila, Jalisco, Morelos, Oaxaca, Sinaloa.

Notes: MZ-FC

***Eburia maccartyi* Noguera, 2002**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Poza Azul; verbatimCoordinates: 26°55'28.03"N, 102°06'56.46"W; samplingProtocol: Sweeping; eventDate: 29-Jul-15; habitat: Desertic shrub; recordedBy: Oscar Pérez; collectionCode: CNIN_CCB07; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Coahuila, Colima, Guanajuato, Guerrero, Jalisco, Nayarit, Oaxaca.

Notes: CNIN

Eburia paraegrota Chemsak and Linsley, 1973

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: San José del Antejo; verbatimCoordinates: 26°58'11.5"N, 102°07'26.4"W; samplingProtocol: By hand; eventDate: 23-Aug-09; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB12; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: San José del Antejo; verbatimCoordinates: 26°58'11.5"N, 102°07'26.4"W; samplingProtocol: By hand; eventDate: 23-Aug-09; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB13; basisOfRecord: Preserved Specimen
- c. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: San José del Antejo; verbatimCoordinates: 26°58'11.5"N, 102°07'26.4"W; samplingProtocol: By hand; eventDate: 23-Aug-09; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB14; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Coahuila, Colima, Jalisco, Sinaloa.

Notes: MZ-FC

Tribe Elaphidiini

Aneflus (Aneflus) obscurus (LeConte, 1873)

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Río Cañón; verbatimCoordinates: 27°00'33.7"N, 102°04'42.3"W; samplingProtocol: Light trap; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB15; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Río Cañón; verbatimCoordinates: 27°00'33.7"N, 102°04'42.3"W; samplingProtocol: Light trap; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB16; basisOfRecord: Preserved Specimen

Host of: *Acacia* sp. (Mimosaceae). (Monné 2020)

Distribution: USA: Arizona, Texas; MEXICO: Chihuahua, Coahuila.

Notes: MZ-FC

Aneflus (Aneflus) prolixus subsp. *insoletus* Chemsak and Linsley, 1963

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: Light trap; eventDate: 20-Aug-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB17; basisOfRecord: Preserved Specimen

Host of: *Acacia berlandieri* Bentham (Mimosaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Chihuahua, Coahuila, Durango, Hidalgo, San Luis Potosí, Tamaulipas.

Notes: MZ-FC

Aneflomorpha rectilinea subsp. *rectilinea* Casey, 1924

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelgas; locality: Rancho PRONATURA; verbatimCoordinates: 26°49'38.9"N, 102°01'24.4"W; samplingProtocol: Light trap; eventDate: 6-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB18; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelgas; locality: Rancho PRONATURA; verbatimCoordinates: 26°49'38.9"N, 102°01'24.4"W; samplingProtocol: Light trap; eventDate: 7-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB19; basisOfRecord: Preserved Specimen

Host of: *Rhus aromatica* Aitchison (Anacardiaceae), *Baccharis sarothroides* A. Gray (Asteraceae), *Quercus* sp. (Fagaceae). (Monné 2020)

Distribution: USA: Arizona, New Mexico, Texas; MEXICO: Baja California Norte, Coahuila (new state record), Jalisco, Sinaloa.

Notes: MZ-FC

Aneflomorpha wernerii Chemsak, 1962

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelgas; locality: Poza Azul; verbatimCoordinates: 26°55'21.2"N, 102°72'3.6"W; samplingProtocol: Light trap; eventDate: 12-Apr-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB20; basisOfRecord: Preserved Specimen

Distribution: USA: Texas; MEXICO: Coahuila (new country record).

Notes: MZ-FC

Anelaphus moestus subsp. *moestus* (LeConte, 1854)

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelgas; locality: Antiguos Mineros; verbatimCoordinates: 26°46'57.7"N, 102°00'20.2"W; samplingProtocol: Light trap; eventDate: 21-Jun-09; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB21; basisOfRecord: Preserved Specimen

Host of: *Rhus aromatica* Aitchison (Anacardiaceae), *Grindelia* sp. (Asteraceae), *Celtis laevigata* Willdenow, *C. lindheimeri* Engelmann, *C. reticulata* Torrey (Cannabaceae), *Sophora secundiflora* (Ortega) de Candolle (Fabaceae), *Quercus* sp. (Fagaceae), *Juglans nigra* Linnaeus (Juglandaceae), *Pithecellobium flexicaule* (Bentham) Coulter, *Prosopis juliflora* (Swartz) de Candolle, *Vachellia rigidula* (Bentham) Seigler and Ebiner, (Mimosaceae), *Morus rubra* Linnaeus (Moraceae), *Pinus cembroides* Zuccarini (Pinaceae), *Zanthoxylum clava-herculis* Linnaeus (Rutaceae). (Monné 2020)

Distribution: USA: Arizona, Texas; MEXICO: Coahuila.

Notes: MZ-FC

***Anoplomorpha rinconia* (Casey, 1924)**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Río Cañón; verbatimCoordinates: 27°00'33.7"N, 102°04'42.3"W; samplingProtocol: Light trap; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB22; basisOfRecord: Preserved Specimen

Host of: *Quercus grisea* Liebmamn (Fagaceae). (Monné 2020)

Distribution: USA: Arizona; MEXICO: Baja California Norte, Coahuila (new state record), Oaxaca, Sinaloa, Sonora.

Notes: MZ-FC

***Elaphidionopsis fasciatipennis* Linsley, 1936**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho PRONATURA; verbatimCoordinates: 26°49'38.9"N, 102°01'24.4"W; samplingProtocol: By hand; eventDate: 8-Jul-10; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB23; basisOfRecord: Preserved Specimen

Host of: *Celtis reticulata* Torrey (Cannabaceae), *Dermatophyllum secundiflorum* (Ortega) Gandhi and Reveal (Fabaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Coahuila.

Notes: MZ-FC

***Neaneflus brevispinus* Chemsak, 1962**

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: Light

- trap; eventDate: 20-Aug-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB24; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénegas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: Light trap; eventDate: 20-Aug-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB25; basisOfRecord: Preserved Specimen

Host of: *Dalea formosa* Torrey (Fabaceae). (Monné 2020)

Distribution: USA: California, Texas; MEXICO: Coahuila (new state record), Zacatecas.

Notes: MZ-FC

***Stenosphenus dolosus* Horn, 1885**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénegas; locality: Poza Azul; verbatimCoordinates: 26°55'28.03"N, 102°06'56.46"W; samplingProtocol: Sweeping; eventDate: 29-Jul-15; habitat: Desertic shrub; recordedBy: Oscar Pérez; collectionCode: CNIN_CCB08; basisOfRecord: Preserved Specimen

Host of: *Baccharis halimifolia* Linnaeus (Asteraceae), *Acacia rigidula* Bentham, *Leucaena* sp., *Prosopis juliflora* (Swartz) de Candolle, *Vachellia farnesiana* (Linnaeus) Wight and Arn., *V. rigidula* (Benth.) Seigler and Ebinger (Mimosaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Coahuila, Nuevo León, San Luis Potosí, Tamaulipas, Veracruz.

Notes: CNIN

Tribe Hesperophanini

***Haplidus laticeps* Knull, 1941**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénegas; locality: Río Cañón; verbatimCoordinates: 27°00'33.7"N, 102°04'42.3"W; samplingProtocol: Light trap; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB26; basisOfRecord: Preserved Specimen

Host of: *Acacia constricta* Bentham ex Gray, *Prosopis juliflora* (Swartz) de Candolle (Mimosaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Chihuahua, Coahuila (new state record).

Notes: MZ-FC

Tribe Methiini

Methia mormona Linell, 1897

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Río Cañón; verbatimCoordinates: 27°00'33.7"N, 102°04'42.3"W; samplingProtocol: Sweeping; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB27; basisOfRecord: Preserved Specimen

Host of: *Berberis harrisoniana* Kearney and Peebles (Berberidaceae), *Juniperus deppeana* Steudel (Cupressaceae), *Juglans major* A. A. Heller (Juglandaceae), *Salix* sp. (Salicaceae). (Monné 2020)

Distribution: USA: Arizona, Colorado, New Mexico, Utah; MEXICO: Chihuahua, Coahuila (new state record).

Notes: MZ-FC

Tribe Rhopalophorini

Rhopalophora angustata Schaeffer, 1905

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Emilio Bichara; samplingProtocol: By hand; eventDate: 25-Sep-16; habitat: Desertic shrub; recordedBy: Cecilia Fernández; collectionCode: CNIN_CCB09; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Emilio Bichara; samplingProtocol: By hand; eventDate: 25-Sep-16; habitat: Desertic shrub; recordedBy: Cecilia Fernández; collectionCode: CNIN_CCB10; basisOfRecord: Preserved Specimen

Host of: *Baccharis neglecta* Nuttall (Asteraceae), *Celtis berlandieri* Klotzsch (Cannabaceae), *Diospyros texana* Scheele (Ebenaceae), *Eysenhardtia polystachya* (Ortega) Sargent, *E. texana* Scheele (Fabaceae), *Pithecellobium flexicaule* (Bentham) Coulter, *P. pallens* (Bentham) Standley (Mimosaceae), *Citrus sinensis* (Linnaeus) Osbeck, *Zanthoxylum fagara* (Linnaeus) Sargent (Rutaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Coahuila (new state record), Nuevo León, Tamaulipas.

Notes: CNIN

Tribe Tillomorphini

Euderces basimaculatus Giesbert and Chemsak, 1997

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: PRONATURA; samplingProtocol: By hand; eventDate: 21-Aug-15; habitat: Desertic shrub; recordedBy: Mayra Cervantes; collectionCode: CNIN_CCB11; basisOfRecord: Preserved Specimen

Host of: *Lysiloma acapulcense* (Kunth.) Benth., *Haematoxylum brasiletto* H. Karst., *Vachellia pennatula* (Schltdl. and Cham.) Seigler and Ebinger (Fabaceae). (Monné 2020)

Distribution: MEXICO: Coahuila (new state record), Colima, Durango, Guerrero, Hidalgo, Jalisco, Michoacán, Nayarit, Oaxaca, Puebla.

Notes: CNIN

Tribe Trachyderini

Chemsakiella michelbacheri (Chemsak, 1984)

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: PRONATURA; samplingProtocol: By hand; eventDate: 25-Aug-15; habitat: Desertic shrub; recordedBy: Mayra Cervantes; collectionCode: CNIN_CCB12; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Coahuila, Tamaulipas, Zacatecas.

Notes: CNIN

Crossidius mexicanus Chemsak and Noguera, 1997

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; samplingProtocol: By hand; eventDate: 18-Aug-14; habitat: Desertic shrub; recordedBy: Ángel Neria; collectionCode: CNIN_CCB13; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'19.36"N, 102°09'04.38"W; samplingProtocol: Sweeping; eventDate: 21-Jul-15; habitat: Desertic shrub; recordedBy: Oscar Pérez; collectionCode: CNIN_CCB14; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Coahuila, Nuevo León, Zacatecas.

Notes: CNIN

***Crossidius suturalis* subsp. *suturalis* LeConte, 1858**

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: By hand; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB28; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: By hand; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB29; basisOfRecord: Preserved Specimen
- c. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: By hand; eventDate: 4-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB30; basisOfRecord: Preserved Specimen
- d. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: By hand; eventDate: 26-Apr-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB31; basisOfRecord: Preserved Specimen
- e. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: By hand; eventDate: 29-Oct-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB32; basisOfRecord: Preserved Specimen
- f. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: By hand; eventDate: 29-Oct-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB33; basisOfRecord: Preserved Specimen

Host of: *Chrysothamnus viscidiflorus* (Hooker) Nuttall, *Gutierrezia longifolia* Greene, *G. lucida* Greene, *G. wrightii* A. Gray, *Haplopappus hartwegi* (Gray) Blake, *H. pluriflorus* (Gray) Hall, *Xanthocephalum glutinosum* (Sprengel) Shinners, *X. sarothrae* (Pursh) Shinners (Asteraceae). (Monné 2020)

Distribution: USA: Arizona, New Mexico, Texas; MEXICO: Chihuahua, Coahuila (new state record), Sonora.

Notes: MZ-FC

***Elytroleptus divisus* (LeConte, 1884)**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'19.36"N, 102°09'04.38"W; samplingProtocol: Sweeping; eventDate: 21-Jul-15; habitat: Desertic shrub; recordedBy: Oscar Pérez; collectionCode: CNIN_CCB15; basisOfRecord: Preserved Specimen

Distribution: USA: Texas; MEXICO: Coahuila, Nuevo León, Querétaro, San Luis Potosí, Tamaulipas.

Notes: CNIN

Neocrossidius trivittatus (Bates, 1880)

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: San José del Antejo; verbatimCoordinates: 26°58'11.5"N, 102°07'26.4"W; samplingProtocol: Sweeping; eventDate: 23-May-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB34; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: San José del Antejo; verbatimCoordinates: 26°58'11.5"N, 102°07'26.4"W; samplingProtocol: Sweeping; eventDate: 23-May-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB35; basisOfRecord: Preserved Specimen

Distribution: MEXICO: Chihuahua, Ciudad de México, Coahuila, Durango, Michoacán, Morelos.

Notes: MZ-FC

Parevander hovorei Giesbert, 1984

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Antiguos Mineros; verbatimCoordinates: 26°46'57.7"N, 102°00'20.2"W; samplingProtocol: By hand; eventDate: 8-Jul-10; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB36; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Antiguos Mineros; verbatimCoordinates: 26°46'57.7"N, 102°00'20.2"W; samplingProtocol: By hand; eventDate: 8-Jul-10; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB37; basisOfRecord: Preserved Specimen
- c. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Antiguos Mineros; verbatimCoordinates: 26°46'57.7"N, 102°00'20.2"W; samplingProtocol: By hand; eventDate: 8-Jul-10; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB38; basisOfRecord: Preserved Specimen

Host of: *Helianthus* sp., *Verbesina* sp., *Viguiera* sp. (Asteraceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Coahuila, Hidalgo, Nuevo León, San Luis Potosí, Tamaulipas.

Notes: MZ-FC

***Plionoma suturalis* (LeConte, 1858)**

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Río Cañón; verbatimCoordinates: 27°00'33.7"N, 102°04'42.3"W; samplingProtocol: Sweeping; eventDate: 31-May-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB39; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: Sweeping; eventDate: 30-Apr-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB40; basisOfRecord: Preserved Specimen

Host of: *Prosopis juliflora* (Swartz) de Candolle, *Vachellia farnesiana* (Linnaeus) Wight and Arn. (Mimosaceae). (Monné 2020)

Distribution: USA: California, Texas; MEXICO: Baja California Norte, Coahuila (new state record), Sonora.

Notes: MZ-FC

***Sphaenothecus bilineatus* (Gory in Guérin-Méneville, 1831)**

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: San José del Antejo; verbatimCoordinates: 26°58'11.5"N, 102°07'26.4"W; samplingProtocol: Sweeping; eventDate: 23-May-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB41; basisOfRecord: Preserved Specimen

Host of: *Baccharis* sp. (Asteraceae), *Cassia* sp. (Caesalpiniaceae), *Leucaena pulverulenta* (Schlechtendal) Benth, *Prosopis juliflora* (Swartz) de Candolle, *Vachellia farnesiana* (Linnaeus) Wight and Arn. (Mimosaceae), *Ficus* sp. (Moraceae), *Coffea arabica* Linnaeus (Rubiaceae), *Ulmus crassifolia* Nuttall (Ulmaceae). (Monné 2020)

Distribution: USA: California, Texas; MEXICO: Baja California, Chiapas, Coahuila, Hidalgo, Jalisco, Morelos, Oaxaca, Sonora, Tamaulipas; HONDURAS; NICARAGUA.

Notes: MZ-FC

***Stenaspis solitaria* (Say, 1824)**

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: By hand; eventDate: 3-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB42; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: By hand;

eventDate: 3-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB43; basisOfRecord: Preserved Specimen

Host of: *Acacia berlandieri* Bentham, *Prosopis juliflora* (Swartz) de Candolle (Mimosaceae). (Monné 2020)

Distribution: USA: Arizona, California, Texas; MEXICO: Baja California Norte, Coahuila (new state record), Durango, Sonora.

Notes: MZ-FC

Tylosis jimenezi Dugés, 1879

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: Sweeping; eventDate: 3-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB44; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: Sweeping; eventDate: 3-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB45; basisOfRecord: Preserved Specimen
- c. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Rancho Orozco; verbatimCoordinates: 26°52'18.3"N, 102°05'16.8"W; samplingProtocol: Sweeping; eventDate: 3-Jul-13; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB46; basisOfRecord: Preserved Specimen

Host of: *Sphaeralcea* sp. (Malvaceae). (Monné 2020)

Distribution: USA: Texas; MEXICO: Ciudad de México, Coahuila, Durango, Guanajuato.

Notes: MZ-FC

Subfamily Lamiinae

Tribe Acanthocinini

Moneilema armatum LeConte, 1853

Material

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Churince; verbatimCoordinates: 26°50'30.3"N, 102°08'9.9"W; samplingProtocol: By hand; eventDate: 6-Jun-10; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB47; basisOfRecord: Preserved Specimen

Host of: *Astrophytum asterias* (Zuccarini) Lemaire, *Homalocephala texensis* Britton and Rose, *Marginatocereus marginatus* (de Candolle) Beckeberg, *Opuntia*

arborescens Engelmann, *O. arbuscula* Engelmann, *O. engelmanni* Salm-Dyck, *O. imbricata* de Candolle, *O. leptocaulis* de Candolle, *O. lindheimeri* Engelmann, *O. macrocentra* Engelmann, *O. megacantha* Salm-Dyck, *O. robusta* Wendland, *O. spinosior* Toumey, *O. violacea* Engelmann, *Platyopuntia* sp. (Cactaceae). (Monné 2020)

Distribution: USA: Colorado, Kansas; MEXICO: Coahuila, Tamaulipas.

Notes: MZ-FC

Tribe Onciderini

Oncideres rhodosticta Bates, 1885

Materials

- a. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Poza Azul; verbatimCoordinates: 26°55'21.2"N, 102°72'3.6"W; samplingProtocol: Light trap; eventDate: 24-Aug-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB48; basisOfRecord: Preserved Specimen
- b. country: Mexico; stateProvince: Coahuila; municipality: Cuatro Ciénelas; locality: Poza Azul; verbatimCoordinates: 26°55'21.2"N, 102°72'3.6"W; samplingProtocol: Light trap; eventDate: 24-Aug-12; habitat: Desertic shrub; recordedBy: Marysol Trujano and Uri García; collectionCode: MZ-FC_CCB49; basisOfRecord: Preserved Specimen

Host of: *Parkinsonia aculeata* Linnaeus (Caesalpiniaceae), *Casuarina cunninghamiana* Miquel (Casuarinaceae), *Sarcobatus vermiculatus* (Hooker) Torrey (Chenopodiaceae), *Acacia greggii* A.Gray, *Mimosa* sp., *Pithecellobium flexicaule* (Bentham) Coulter, *Prosopis juliflora* (Swartz) de Candolle, *P. glandulosa* Torrey, *Vachellia farnesiana* (Linnaeus) Wight. and Arn. (Mimosaceae). (Monné 2020)

Distribution: USA: Arizona, New Mexico, Texas; MEXICO: Baja California Norte, Chihuahua, Coahuila, Durango, Sonora, Tamaulipas.

Notes: MZ-FC

Analysis

As this study is the first species list of Cerambycidae fauna in a Mexican desert environment, it greatly increases the knowledge of this group in Mexico. We recorded 37 species from 32 genera and identified 13 tribes from four subfamilies (Prioninae, Lepturinae, Cerambycinae and Lamiinae) Table 1. The subfamily Cerambycinae had the greatest number of species at 31 (84% of the total). The other subfamilies only registered two species each. This revealed a pattern similar to that of other habitats such as tropical dry forest (e.g. Toledo et al. 2002, Noguera et al. 2012), but in this case, the results were even more pronounced.

Table 1.

Number of tribes, genera and species of the cerambycids from Cuatro Ciénegas Basin

Subfamily	No. of tribes	No. of genera	No. of species
Prioninae	1	2	2
Lepturinae	1	2	2
Cerambycinae	9	26	31
Lamiinae	2	2	2
Total	13	32	37

Discussion

Despite the fact that Mexico has extensive area of deserts, faunistic studies relating to Cerambycidae have previously not been conducted in this or similar xerophilous habitats. In Mexico, the family Cerambycidae has been studied mainly in tropical dry forest (e.g. Chemsak and Noguera 1993, Noguera et al. 2012). Only sporadic records from desertic areas have been reported (Bezark 2020).

MacKay et al. (1987) recorded 44 species from the northern region of the Chihuahuan desert near Las Cruces, New Mexico, USA. Compared to our study, they obtained the same species distribution in subfamilies. Cerambycinae were extremely abundant, followed by Prioninae and Lamiinae, both being well represented. Only one species of Lepturinae was recorded. MacKay et al. (1987) argue that Lepturinae species depend on moisture for development, hence their species richness in arid environments is poor. On the other hand, Cerambycinae presents the greatest number of genera with diurnal habits (Linsley 1959). In the case of CCB, the genera of Cerambycinae were more common in daytime activity. This is presumably due to different types of adult feeding which is anthophilous in several genera of Cerambycinae (Švácha and Lawrence 2014).

Of the species recorded in this study, *Meloemorpha aliena* (Bates), *Knulliana cincta cincta* (Drury), *Megacyllene antennata* (White), *Aneflomorpha rectilinea rectilinea* Casey, *Anopliomorpha rinconia* (Casey), *Neaneplus brevispinus* Chemsak, *Haplidus laticeps* Knull, *Methia mormona* Linell, *Rhopalophora angustata* Schaeffer, *Euderces basimaculatus* Giesbert & Chemsak, *Crossidius suturalis suturalis* LeConte, *Plionoma suturalis* (LeConte) and *Stenaspis solitaria* (Say) are new records for the state of Coahuila. *Typocerus sinuatus* (Newman), *Neoclytus mucronatus mucronatus* (Fabricius) and *Aneflomorpha wernerii* Chemsak are reported for the first time in Mexico. These three species have been previously recorded in the southern United States (Texas).

Possibly due in part to extensive collecting efforts, the states of Veracruz (406), Oaxaca (343), Jalisco (327) and Chiapas (315) (Noguera 2014) have the highest species records of Cerambycidae (numbers in parenthesis indicate the numbers of species recorded from

the corresponding state). Noguera (2014) indicates the species richness from Coahuila at 24 species. Nevertheless, Bezark (2020) in the electronic catalogue of Cerambycidae recorded 43 species from Coahuila. Our work represents a considerable contribution and updated inventory of Mexican Cerambycidae in the Coahuila region. It presents new distributional records that represent a 41% increase in species richness. It is clear that it is necessary to continue exploring more regions of Mexico and publishing updated Cerambycidae distribution data to complete our knowledge, not only of desert environments, but also of other Mexican ecosystems present in the country.

The distribution of species that were recorded in CCB shows a mixture between cerambycid species with distributional records in United States and Mexico (Bezark 2020). Many of these species have diurnal habits and an affinity for desert environments. This is useful for understanding the distributional patterns and evolutionary history of cerambycid beetles. The ecological factors and historical events from the *Altiplano Mexicano* are relevant for determining the geographic distribution of insect species from CCB (Álvarez and Ojeda 2019). Additionally, the presence of host plants in these desert ecosystems is fundamental to their existence in these environments (Pinkava 2016).

The CCB is a unique region with different components from Neotropical and Nearctic biogeographic provinces. This makes it an essential crossroads to understand the evolution of several faunal groups (Álvarez and Ojeda 2019). This work represents the first species list of Cerambycidae in a Mexican desert (CCB, Chihuahuan desert) and Coahuila. Cerambycids provide important ecological services for desert ecosystems. They act as pollinators (MacKay et al. 1987), active agents of decomposition and recyclers of organic matter. As ecosystem agents, these beetles are indispensable for the health and permanency of CCB environments.

Acknowledgements

The authors thank M. Trujano-Ortega and U. García-Vázquez for providing the material to realise this study; CONABIO (JF065) to A. Nieto, for fieldwork; WWF-Alianza Carlos Slim (L039) and the Theodore Roosevelt Memorial Fund (American Museum of Natural History) for resources provided to M. Trujano-Ortega and U. García-Vázquez. We also thank PRONATURA A. C., DESUVALLE A.C. and CONANP for their assistance with fieldwork and C. Mayorga (CNIN-IBUNAM) for allowing access to the collections under their care. We would also like to thank Hans Clebsch for comments on the manuscript and helping to improve its English content.

References

- Álvarez F, Ojeda M (2019) Animal Diversity and Biogeography of the Cuatro Ciénebas Basin. Springer, Cham, Switzerland, 231 pp. <https://doi.org/10.1007/978-3-030-11262-2>
- Bader E (2005) ArcGIS Server Administrator and Developer Guide. Redlands, California.

- Bezark LG, Tyson WH, Schiff NM (2013) New species of Cerambycidae from Panama, with new distribution records (Coleoptera, Cerambycidae). Zootaxa 3608 (4): 273-277. <https://doi.org/10.11646/zootaxa.3608.4.5>
- Bezark LG (2020) Checklist of the Oxypeltidae, Vesperidae, Disteniidae and Cerambycidae, (Coleoptera) of the Western Hemisphere. <http://bezbycids.com/byciddb/wdefault.asp?w=n>. Accessed on: 2020-5-20.
- Chemsak JA, Linsley EG (1965) New genera and species of North American Cerambycidae (Coleoptera). The Pan-Pacific Entomologist 41 (3): 141-153. URL: <https://www.biodiversitylibrary.org/page/53605987>
- Chemsak JA (1991) New Mexican and Central American Elaphidiini (Coleoptera: Cerambycidae). Anales del Instituto de Biología 62 (3): 469-480.
- Chemsak JA, Noguera FA (1993) Annotated checklist of the Cerambycidae of the Estacion de Biología Chamela, Jalisco, Mexico (Coleoptera) with descriptions of new genera and species. Folia Entomológica Mexicana 89: 55-102.
- Chemsak JA, Noguera FA (2005) New Mexican species of the Elaphidionine genus *Aneflomorpha* Casey, 1912 (Coleoptera: Cerambycidae), with a key to the Mexican species. The Pan-Pacific Entomologist 81: 20-33.
- García E (1964) Modificaciones al sistema de clasificación climática de Köppen (para adaptarlo a las condiciones de la República Mexicana). Offset Larios, D. F., Mexico, 246 pp.
- Giesbert EF, Chemsak JA (1989) The genus *Stenosphenus* Haldeman (Coleoptera: Cerambycidae). The Pan-Pacific Entomologist 65 (3): 269-301. URL: <https://www.biodiversitylibrary.org/page/56180393>
- Giesbert EF, Chemsak JA (1993) A review of the Rhopalophorini (Coleoptera: Cerambycidae) of North and Central America. Insecta Mundi 7 (1-2): 27-64. URL: <https://journals.fvc.org/mundi/article/view/24723>
- Grzymala TL, Miller KB (2013) Taxonomic revision and phylogenetic analysis of the genus *Elytroleptus* Dugés (Coleoptera: Cerambycidae: Cerambycinae: Trachyderini). Zootaxa 3659 (1): 1-62. <https://doi.org/10.11646/zootaxa.3659.1.1>
- Linsley EG (1959) Ecology of Cerambycidae. Annual Review of Entomology 4 (1): 99-138. <https://doi.org/10.1146/annurev.en.04.010159.000531>
- Linsley EG (1963) The Cerambycidae of North America. Part IV. Taxonomy and classification of the subfamily Cerambycinae, tribes Elaphidionini through Rhinotragini. 21. University of California Publications in Entomology, California, Los Angeles, 165 pp.
- Linsley EG (1964) The Cerambycidae of North America. Part V. Taxonomy and classification of the subfamily Cerambycinae, tribes Callichromatini through Ancylocerini. 22. University of California Publications in Entomology, California, Los Angeles, 197 pp.
- Linsley EG, Chemsak JA (1972) Cerambycidae of North America. Part VI. No. 1. Taxonomy and classification of the subfamily Lepturinae. 69. University of California Publications in Entomology, California, Los Angeles, 138 pp.
- MacKay WP, Zak JC, Hovore FT (1987) Cerambycid beetles (Coleoptera: Cerambycidae) of the northern Chihuahuan Desert (South Central New Mexico). The Coleopterists Bulletin 41 (4): 361-369. URL: <https://www.jstor.org/stable/4008520>
- Monné ML, Monné MA, Wang Q (2017) General morphology, classification and biology of Cerambycidae. In: Wang Q (Ed.) Cerambycidae of the world. Biology and Pest Management. CRC Press, Florida, U.S.A, 642 pp.

- Morafka DJ (1977) A biogeographical analysis of the Chihuahuan Desert through its herpetofauna. *Biogeographica* 9: 1-313.
- Noguera FA (2002) Revisión taxonómica de las especies del género *Eburia* Lepeletier y A.-Serville in Lacordaire de Norte y Centroamérica (Coleoptera: Cerambycidae). *Folia Entomológica Mexicana* 41 (1): 1-167.
- Noguera FA, Zaragoza-Caballero S, Rodríguez-Palafox A, González-Soriano E, Ramírez-García E, Ayala R, Ortega-Huerta MA (2012) Cerambícidos (Coleoptera: Cerambycidae) del bosque tropical caducifolio en Santiago Dominguillo, Oaxaca, México. *Revista Mexicana de Biodiversidad* 83 (3): 611-622.
- Noguera FA (2014) Biodiversidad de Cerambycidae (Coleoptera) en México. *Revista Mexicana de Biodiversidad* 85: 290-297. <https://doi.org/10.7550/rmb.32966>
- Pinkava DJ (2016) Flora de la región del bolson de Cuatro Ciénegas, Coahuila, México, I. *Botanical Sciences* 38: 35-73. <https://doi.org/10.17129/botsci.1168>
- Santos-Silva A (2007) Revisão do gênero *Derobrachus* Audinet-Serville, 1832 (Coleoptera, Cerambycidae, Prioninae). *Arquivos de Zoologia* 38 (1): 1-94. <https://doi.org/10.11606/issn.2176-7793.v38i1p1-94>
- Santos-Silva A, Nearns EH, Swift IP (2016) Revision of the American species of the genus *Prionus* Geoffroy, 1762 (Coleoptera, Cerambycidae, Prioninae, Prionini). *Zootaxa* 4134 (1): 1-103. <https://doi.org/10.11646/zootaxa.4134.1.1>
- Souza V, Siebert JL, Escalante AE, Elser JJ, Eguiarte LE (2012) The cuatro ciéregos basin in Coahuila, Mexico: an astrobiological precambrian park. *Astrobiology* 12 (7): 641-647. <https://doi.org/10.1089/ast.2011.0675>
- Švácha PO, Lawrence JF (2014) Cerambycidae Latreille, 1802. In: Leschen RAB, Beutel RG (Eds) *Arthropoda: Insecta: Coleoptera. Morphology and Systematics Phytophaga*. Walter de Gruyter, Handbook of Zoology, Berlin, Germany, 675 pp.
- Tavakilian G, Chevillotte H (2020) TITAN: Cerambycidae database. <http://titan.gbif.fr/>. Accessed on: 2020-5-06.
- Toledo VH, Noguera FA, Chemsak JA, Hovore FT, Giesbert EF (2002) The cerambycid fauna of the tropical dry forest of "El Aguacero," Chiapas, México (Coleoptera: Cerambycidae). *The Coleopterists Bulletin* 56 (4): 515-532. [https://doi.org/10.1649/0010-065X\(2002\)056\[0515:TCFOTT\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2002)056[0515:TCFOTT]2.0.CO;2)
- Turnbow RH, Thomas MC (2002) Cerambycidae Leach 1815. In: Arnett RH, Thomas MC, Skelley PE, Frank JH (Eds) *American Beetles, Volume II: Polyphaga: Scarabaeoidea through Curculionoidea*. 2. CRC press, Florida, Boca Raton, 861 pp. <https://doi.org/10.1201/9781420041231.ch8>

Supplementary material

Suppl. material 1: Checklist_Taxon_CCB [doi](#)

Authors: Pérez-Flores, O.

Data type: Taxon information and occurrence information

[Download file](#) (52.51 kb)