



Research Article

First systematic inventory of the jumping plant lice of Luxembourg (Hemiptera, Sternorrhyncha, Psylloidea)

Carmelo Rapisarda[†], Alexander M. Weigand[§], Paul Braun[§], Michael Eickermann[¶]

[†] Dipartimento di Agricoltura, Alimentazione e Ambiente (Di3A), Università degli Studi di Catania, Catania, Italy

[§] Musée National d'Histoire Naturelle (MNHN), Luxembourg, Luxembourg

[¶] Luxembourg Institute of Science and Technology, Belvaux, Luxembourg

Corresponding author: Michael Eickermann (michael.eickermann@list.lu)

Academic editor: Yasen Mutafchiev

Received: 03 Nov 2021 | Accepted: 01 Jul 2022 | Published: 04 Aug 2022

Citation: Rapisarda C, Weigand AM, Braun P, Eickermann M (2022) First systematic inventory of the jumping plant lice of Luxembourg (Hemiptera, Sternorrhyncha, Psylloidea). Biodiversity Data Journal 10: e77571.

<https://doi.org/10.3897/BDJ.10.e77571>

Abstract

Psyllids (superfamily Psylloidea), also known as jumping plant lice, are a group of plant-sap sucking Hemiptera having significant pest status for crops, forest trees and ornamental plants. Only seven species of psyllids have been recorded in Luxembourg so far. An additional group of seven species has been recorded exclusively, based on the findings of their galls or specific plant deformations; but no mention exists in literature on the actual collection of the inducing insect in Luxembourg. To fill this knowledge gap, field collections were carried out during the years 2019–2020. In addition, samples from 1999–2000 stored in the wet collection of the Musée National d'Histoire Naturelle de Luxembourg were studied. This research, in combination with information coming from literature, allowed us to list 48 species of the families Aphalaridae (5 species), Liviidae (5), Psyllidae (24) and Triozidae (14), though the presence of one species within the last family (*Trioza rhamni*) needs to be confirmed. Brief information on geographical distribution, biology and (if available) illustrations of diagnostic characters are provided on the psyllid species detected in Luxembourg so far.

Keywords

Psyllids, Luxembourg, species diversity, biology, ecology

Introduction

Psyllids (superfamily Psylloidea), also known as jumping plant lice, are a group of plant-sap sucking Hemiptera which may have significant pest status for crops, forest trees and ornamental plants due to their copious production of honeydew, their frequent coating with waxy secretions (smearing the canopy of infested plants), the injection of toxic saliva (causing necrosis, deformations or galls) and, last but not least, their responsibility in transmitting many pathogens to plants, mainly bacteria and especially phytoplasmas (Hodkinson 1974, Hodkinson 1984, Burckhardt 1994a, Burckhardt 2005, Munyaneza 2010, Haapalainen 2014, Ben Othmen et al. 2019). In spite of their importance, psyllids are still poorly known in Luxembourg; this is why their species composition, diversity and distribution need to be properly assessed.

According to literature, *Spanioneura buxi* (Linnaeus), *S. fonscolombii* Foerster and *Trichochermes walkeri* (Foerster) are the only species of Psylloidea whose presence in Luxembourg has been reported till now based on adult specimens (Baugnée 2001, O'Connor and Malumphy 2011, Eickermann et al. 2020). However, additional records of psyllids can be also considered, deriving from rich cecidological literature dealing with the presence of plants with galls or deformations probably induced by Psylloidea. Yet, these records need to be confirmed by finding the insects to which the effects on plants are attributed. In particular, Lambinon and Schneider (2004) investigated deformations caused to plants by *Cacopsylla mali* (Schmidberger), *C. melanoneura* (Foerster), *Livia junci* (Schrink), *Spanioneura buxi* (Linnaeus), *Psyllopsis fraxini* (Linnaeus), *Lauritrioza alacris* (Flor), *Trioza flavipennis* Foerster, *T. remota* Foerster and *T. rhamni* (Schrink), in addition to the ones caused by *T. walkeri* (already reported for Luxembourg). For *P. fraxini*, the authors reported the rear of the psyllid from leaf galls collected on *Fraxinus excelsior* L. from many localities in Luxembourg (Bettembourg, Bonnevoie, Bridel and Kleinbettingen). Similarly, they reported findings of eggs of *T. flavipennis* in pit galls found on leaves of *Aegopodium podagraria* L. For all remaining cecidia described, however, no indication was given on the finding of the causing insect. Deformations caused by three additional psyllid species [*Camarotoscena speciosa* (Flor), *Trioza centranthi* (Vallot) and *T. urticae* (Linnaeus)] were reported by Lambinon et al. (2012), who did not mention any finding of their causing insects. Deformations by *L. alacris* and *T. centranthi* and galls by *T. remota* were reported again by Schneider (2016) from new localities in Luxembourg. Only for *T. remota*, the author referred to the presence of an insect immature within each pit gall found on leaves of *Quercus petraea* (Matt.) Liebl. and *Q. robur* L. Cecidia caused by *C. speciosa*, *L. junci* and *T. flavipennis* were reported again by Burton et al. (2019), who explicitly mentioned for the first time the finding of nymphs of *C. speciosa* and their abundant wax secretion on the leaf deformations of *Populus nigra* L. and confirmed the occurrence of a *T. flavipennis* nymph in the concave leaf gall produced on *A. podagraria*.

Considering the available literature data, seven psyllid species were known for Luxembourg, based on observations of specimens (*Camarotoscena speciosa*, *Psyllopsis fraxini*, *Spanioneura buxi*, *S. fonscolombii*, *Trichochermes walkeri*, *Trioza flavigaster* and *T. remota*) and seven further species were only recorded, based on the findings of galls or deformations they cause to plants (*Cacopsylla mali*, *C. melanoneura*, *Livia junci*, *Lauritrioza alacris*, *Trioza centranthi*, *T. rhamni* and *T. urticae*).

Even when hypothesising that all psyllid records, based on findings of their galls, will be confirmed by collection of insect specimens, still the number of psyllid species presently known in Luxembourg is remarkably low, for example, when compared with faunas of neighbouring countries in the “Benelux” region”: 69 species known in The Netherlands (den Bieman et al. 2019); at least 56 species in Belgium (Ouvrard 2021). Present knowledge on the psyllid fauna of Luxembourg appears weak also when considering the richness and composition of the regional flora (especially for plant groups that may host psyllid species), as assessed by Lambinon and Verloove (2012). The lack of a systematic inventory of the psyllids of Luxembourg, in combination with their ecological and partly agricultural significance, gave rise to the current manuscript.

Data resources

Study area

The Grand Duchy of Luxembourg is characterised by a temperate, semi-oceanic climate. Even though the area of the country is small (2,586 km²) with a maximal Euclidean distance of 82 km from north to south and 57 km from east to west, Luxembourg offers quite diverse physiogeography with different climatological characteristics, associated vegetation and anthropogenic land use. Traditionally, Luxembourg is divided into two main ecoregions, the Oesling (32% of total area) and the Gutland (68%) (Dohet et al. 2008). The Oesling is located in the north of the country, in the border region between Belgium and Luxembourg – and is the eastern part of woody mountains called the Ardennes with highest altitude of 450 m a.s.l. Annual precipitation ranges from 800–1000 mm and annual mean air temperature is 7.5°C (Goergen et al. 2013). The region is characterised by meadows, pastures and forests of coniferous/deciduous trees. The Gutland shows a higher level of anthropogenic disturbance (including larger cities and industrial areas) and a longer vegetational period in comparison to the Oesling (Goergen et al. 2013). Annual precipitation ranges from 700–800 mm and annual mean air temperature is 9°C (Goergen et al. 2013). It can be divided into four sub-ecoregions: the Western and the Eastern Gutland, the Moselle Valley and the Minette Basin. Land use is quite diverse in the Gutland. Pastures are still common, while the acreage of arable land is increasing. Extended deciduous forests are typical for the Gutland (Dietz and Pir 2009). For the Western Gutland, secondary sandstones and sandy soils are common (Stevens et al. 2010) and the amount of precipitation is higher compared to the eastern part (Goergen et al. 2013). The Eastern Gutland is characterised by loam-loess-based soils and a more variable topography (Stevens et al. 2010). The Minette Basin is located in the southwestern part of Luxembourg, next to the French border. It represents the former mining

district, due to the rich resources of iron ore. The Moselle Valley – along the border to Germany – represents the smallest ecoregion (1% of total area of Luxembourg). It is sunnier and also drier (less than 700 mm) in comparison to all other areas of the country. Due to these very specific climatic conditions, vineyards are prevalent and agriculture shows a higher level of intensification (Dohet et al. 2008). For further information about climatological characteristics of the different regions in Luxembourg, see Eickermann et al. (2014).

Field collection of psyllids, their preparation and identification (2019/2020)

Adult psyllids were collected during one year (July 2019 to June 2020) by beating host or shelter plants over a sweeping-net, from which the specimens were captured into plastic tubes containing 70% ethanol, thereby exploiting their natural aptitude to jump. When easily visible on the plants, nymphs were also collected by picking them directly and storing them in plastic tubes as described for the adults. If necessary, plants on which samples were collected (or parts of them) were stored in plastic bags and taken to the laboratory for their specific identification, according to Lambinon and Verloove (2012).

In the lab, all specimens collected in each tube were observed using a dissecting microscope (LEICA MZ7.5); in this phase, adults were separated by sex, counted and (where possible) identified. For samples whose observation under a dissecting microscope allowed the species identification, all specimens (adults and nymphs, if present) were stored in glass tubes containing 70% ethanol, each marked with a progressive collection number, corresponding to those of the general collection register containing complete data on the date and location of collection (including the geographical coordinates) and on the plant(s) on which the sample was recorded. In case a closer examination was needed to allow species identification, a maximum of six specimens (if available, three of each sex) were mounted on permanent microscope slides. For slide preparation, selected adults were firstly cleaned in 70% ethanol under gentle heating, then left to clear overnight in 10% potassium hydroxide (KOH), rinsed in a solution of 20% glacial acetic acid, dehydrated in 95% ethanol (for 10 minutes), placed in xylene for ≥ 10 minutes, mounted in Canada Balsam and allowed to dry in an incubator for 15 days at 35°C.

Each specimen was dissected in a drop of mounting medium before being mounted on the slide, in order to mount all different parts of the body separately: head (mounted dorsal uppermost), pronotum and forelegs, mouth parts, mesoscutum and forewings (dorsally), ventral mesothorax and mid-legs, metathorax, hind legs and abdomen (laterally, with well-exposed genitalia).

Photographs of morphological details (head, forewing, male and female terminalia) were made at the Musée National d'Histoire Naturelle de Luxembourg (MNHN), from ethanol-preserved specimens of most of the species collected, using a Keyence VHX-6000 digital microscope.

With the exception of some special cases, for which we used literature specifically concerning the taxonomic group of the species to be identified, the material was identified

by using the taxonomic keys of Hodkinson and White (1979) and Ossiannilsson (1992), following the classification and nomenclature proposed by Burckhardt et al. (2021).

All the material studied, i.e. wet and slide mounted material, is stored at the MNHN, apart from several specimens being part of abundant field collected samples, which are stored in the collections of the museum of the University of Catania (Italy).

Study of psyllid material from Malaise and Moericke traps (1999/2000)

During the years 1999-2000, an entomological collection campaign was carried out in Luxembourg by Evelyne Carrières, focussing on the national inventory of hoverflies (Syrphidae) (Carrières 2003). The project was financed by the MNHN. Several Malaise and yellow Moericke traps were deployed in various areas of the country and regularly controlled every 2-4 weeks after initial installation. Many of the bulk samples collected during this campaign (and stored in the MNHN wet collection) contained psyllid specimens, which, as part of the research aimed at compiling this manuscript, were sorted out, observed under a LEICA dissecting microscope, separated by sex, counted and (where possible) identified as already reported in the previous section. Additionally, in this case, all material identified by the simple use of a dissecting microscope has been stored in glass tubes containing 70% ethanol and marked with progressive collection numbers (different from those used for the field collected material). A general register has been prepared, containing complete data on collection localities (including the geographical coordinates) and the dates of exposure and removal of the traps. In case a more sophisticated approach was necessary to identify some material, permanent slides were prepared and identifications performed as described in the previous section. All material studied, both ethanol preserved or mounted on permanent slides, is stored at the MNHN.

Results

Overall, the reliable bibliographic references (e.g. those reporting the actual collection of Psylloidea specimens), the field collections we made during 2019/2020 and the investigated Malaise trap material from 1999/2000 stored at the MNHN allowed us to record a total of 47 species, belonging to the families Aphalaridae (5 species), Liviidae (5), Psyllidae (24) and Trioziidae (13). For one additional species [*T. rhamni* (Schrank)], belonging to the family Trioziidae, only records in literature are available relating to the presence of the galls it causes on the host plant; therefore, its presence in Luxembourg is highly probable, but needs to be confirmed by observations of psyllid specimens. The total dataset originating from the analysed 1999/2000 and 2019/2020 material can be retrieved from <https://doi.org/10.15468/svhf53> (Rapisarda et al. 2021).

In the following, brief information is provided for all 48 species whose presence has been ascertained or, as reported above for *T. rhamni*, are very likely in Luxembourg. Depicted are head structures (Figs 1, 2, 3, 4), forewings (Figs 5, 6, 7, 8), female and male terminalia (Figs 9, 10, 11, 12, 13, 14, 15, 16).




Figure 1. [doi](#)

Overview of head structures of psyllid species in Luxembourg (Aphalaridae, Liviidae).

A *Aphalara* sp. gr. *polygoni* Foerster, 1848 **B** *Craspedolepta nebulosa* (Zetterstedt, 1828) **C** *Craspedolepta nervosa* (Foerster, 1848) **D** *Craspedolepta subpunctata* (Foerster, 1848) **E** *Rhinocola aceris* (Linnaeus, 1758) **F** *Livia junci* (Schrank, 1789) **G** *Psyllopsis fraxini* (Linnaeus, 1758) **H** *Psyllopsis fraxinicola* (Foerster, 1848) **I** *Strophingia ericae* (Curtis, 1835).

Aphalaridae

Aphalara sp. gr. *polygoni* Foerster, 1848

(Figs 1, 5, 9)

Findings in Luxembourg. Personal field collection by the authors: Oesling: Fléiber (1 ♀, 20.VIII.2019, by general sweeping with net, including clover, *Phacelia* sp. and *Atriplex* sp.), Fussekaul (1 ♀, 22.VIII.2019, on *Rumex* sp.), Goebelsmuehle (2 ♀♀, 20.VIII.2019, on *Rumex* sp.); West Gutland: Eschdorf (2 ♀♀, 14.VIII.2019, on *Rumex* sp.).

Material studied in the MNHN collection: Minette: Kockelscheuer, Conter Jans Boesch (1 ♀, 2.IX - 27.IX.1999, Malaise).

New record for Luxembourg: no member of the genus *Aphalara* Foerster has been recorded in Luxembourg so far; the identification of this taxon needs to be validated by the investigation of male specimens.

According to Burckhardt and Lauterer (1997b) and Cho et al. (2017b), species of this group have a Palaearctic distribution and live on plants of the family Polygonaceae, overwintering on shelter plants. Due to a high similarity of species within the genus *Aphalara* Foerster and the recent clarification of their taxonomic status (Burckhardt and Lauterer 1997b), many records of *A. polygoni* in various countries are doubtful. Hence, we decided to be more conservative by referring to the species group *A. polygoni* instead.




Figure 2. doi

Overview of head structures of psyllid species in Luxembourg (Psyllidae I).

A *Arytaina genistae* (Latreille, 1804) **B** *Arytainilla spartiophila* (Foerster, 1848) **C** *Cacopsylla affinis* (Löw, 1880) **D** *Cacopsylla ambigua* (Foerster, 1848) **E** *Cacopsylla crataegi* (Schrank, 1801) **F** *Cacopsylla mali* (Schmidberger, 1836) **G** *Cacopsylla melanoneura* (Foerster, 1848) **H** *Cacopsylla nigrita* (Zetterstedt, 1828) **I** *Cacopsylla peregrina* (Foerster, 1848) **J** *Cacopsylla* sp. gr. *pruni* (Scopoli, 1763) **K** *Cacopsylla pulchra* (Zetterstedt, 1838) **L** *Cacopsylla pyri* (Linnaeus, 1758) **M** *Cacopsylla pyricola* (Foerster, 1848) **N** *Cacopsylla pyrisuga* (Foerster, 1848) **O** *Cacopsylla rhamnicola* (Scott, 1876) **P** *Cacopsylla visci* (Curtis, 1835).




Figure 3. [doi](#)

Overview of head structures of psyllid species in Luxembourg (Psyllidae II).

A *Livilla ulicis* Curtis, 1836 **B** *Psylla alni* (Linnaeus, 1758) **C** *Psylla betulae* (Linnaeus, 1758) **D** *Psylla foersteri* Flor, 1861 **E** *Psylla hartigii* Flor, 1861 **F** *Spanioneura buxi* (Linnaeus, 1758) **G** *Spanioneura fonscolombii* Foerster, 1848.

***Craspedolepta nebulosa* (Zetterstedt, 1828)**

(Figs 1, 5, 9, 13)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Arsdorf (7 ♂♂, 5 ♀♀, 8.VI.2020, on *Epilobium angustifolium*).

New record for Luxembourg.

Geographical distribution. Apart from North Africa, *C. nebulosa* is widely distributed in the Holarctic Region, from Far East Russia to nearly all north and central Europe, through Central Asia and was recorded also in North America (Canada, USA) (Hodkinson and White 1979, Burckhardt 1983, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Tishetshkin 2007, O'Connor and Malumphy 2011, Percy et al. 2012, Drohojowska and Burckhardt 2014, Ouvrard et al. 2015, den Bieman et al. 2019, Drohojowska and Klasa 2019).

Biology. Monophagous on *Epilobium angustifolium* L. (Onagraceae) (Hodkinson and White 1979, Burckhardt 1983, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Tishetshkin 2007, Percy et al. 2012), on which it performs one generation per

year and overwinters as a nymph (usually 4th instar) on the roots. This psyllid causes both foliar (optional) and root galls. The former consist in downward foldings of the leaf margin, which in June-July, protect the eggs (arranged in a long line along the lower margin) and subsequently the young nymphs; these very soon (already at the 1st or 2nd instar) migrate to the roots, where they overwinter (Sampò 1975). Root galls are large, thread-like and tangled (Lauterer 1976).




Figure 4. [doi](#)

Overview of head structures of psyllid species in Luxembourg (Trioziidae).

- A** *Bactericera albiventris* (Foerster, 1848)
- B** *Bactericera curvatineris* (Foerster, 1848)
- C** *Bactericera substriola* Ossiannilsson, 1992
- D** *Eryngiofaga lautereri* Loginova, 1977
- E** *Lauritrioza alacris* (Flor, 1861)
- F** *Trioza abdominalis* Flor, 1861
- G** *Trioza cirsii* Löw, 1881
- H** *Trioza galii* Foerster, 1848
- I** *Trioza remota* Foerster, 1848
- J** *Trioza urticae* (Linnaeus, 1758).

***Craspedolepta nervosa* (Foerster, 1848)**

(Figs 1, 5, 9)

Findings in Luxembourg. Material studied in the MNHN collection: East Gutland: Godbrange, Schléidelbierg (1 ♀, 27.V - 9.VI.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Eurosiberian distribution, from Yakutskiya and Siberia, through Mongolia, Central European Russia and Iraq, to nearly all Europe (Wagner and Franz 1961, Loginova 1968, Loginova 1974, Hodkinson and White 1979, Burckhardt 1983, Rapisarda 1985, Burckhardt and Kofler 1991, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Lauterer 1994, Lauterer 2001, Seljak 2006, Ripka 2009, Malenovský et al. 2011, O'Connor and Malumphy 2011, Malenovský and Lauterer 2012, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, den Bieman et al. 2019, Drohojowska and Klasa 2019).

Biology. Oligophagous on plants of the genera *Achillea* L. (especially *A. millefolium* L., but also *A. gerberi* Willd., *A. micrantha* Willd., *A. nobilis* L., *A. ptarmica* L.) and *Cirsium* Miller [especially *C. arvense* (L.) Scop.] (Asteraceae) (Wagner and Franz 1961, Loginova 1968, Loginova 1974, Hodkinson and White 1979, Burckhardt 1983, Burckhardt and Kofler 1991, Ossiannilsson 1992, Conci et al. 1993, Lauterer 1994, Conci et al. 1996, Lauterer 2001, Seljak 2006, Ripka 2008, Hodkinson 2009, Malenovský et al. 2011, Malenovský and Lauterer 2012). It develops one generation a year and overwinters as a nymph on the roots of its host plants (Lauterer 1991).




Figure 5. [doi](#)

Overview of forewings of psyllid species in Luxembourg (Aphalaridae, Liviidae).

A *Aphalara* sp. gr. *polygoni* Foerster, 1848 **B** *Craspedolepta nebulosa* (Zetterstedt, 1828) **C** *Craspedolepta nervosa* (Foerster, 1848) **D** *Craspedolepta subpunctata* (Foerster, 1848) **E** *Rhinocola aceris* (Linnaeus, 1758) **F** *Livia junci* (Schrank, 1789) **G** *Psylloopsis fraxini* (Linnaeus, 1758) **H** *Psylloopsis fraxinicola* (Foerster, 1848) **I** *Strophingia ericae* (Curtis, 1835).

***Craspedolepta subpunctata* (Foerster, 1848)**

(Figs 1, 5, 9, 13)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Arsdorf (4 ♂♂, 3 ♀♀, 3 nymphs, 8.VI.2020, on *Epilobium angustifolium*).




Figure 6. [doi](#)

Overview of forewings of psyllid species in Luxembourg (Psyllidae I).

A *Arytaina genistae* (Latreille, 1804) **B** *Arytainilla spartiophila* (Foerster, 1848) **C** *Cacopsylla affinis* (Löw, 1880) **D** *Cacopsylla ambigua* (Foerster, 1848) **E** *Cacopsylla crataegi* (Schrank, 1801) **F** *Cacopsylla mali* (Schmidberger, 1836) **G** *Cacopsylla melanoneura* (Foerster, 1848) **H** *Cacopsylla nigrita* (Zetterstedt, 1828) **I** *Cacopsylla peregrina* (Foerster, 1848) **J** *Cacopsylla* sp. gr. *pruni* (Scopoli, 1763) **K** *Cacopsylla pulchra* (Zetterstedt, 1838) **L** *Cacopsylla pyri* (Linnaeus, 1758) **M** *Cacopsylla pyricola* (Foerster, 1848) **N** *Cacopsylla pyrisuga* (Foerster, 1848) **O** *Cacopsylla rhamnicola* (Scott, 1876) **P** *Cacopsylla visci* (Curtis, 1835).

New record for Luxembourg.

Geographical distribution. Similar to *Craspedolepta nebulosa*, it is widely distributed in the north and central part of the Holarctic Region: from Far East Russia, through Central Asia, to nearly all north and central Europe and North America (Canada, USA) (Hodkinson and White 1979, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Burckhardt 1994b, Conci et al. 1996, Tishetshkin 2007, Gertsson 2010, O'Connor and Malumphy 2011, Serbina et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Monophagous on *Epilobium angustifolium* L. (Onagraceae). According to Lauterer and Baudys (1968), adults fly in late May/early June and lay eggs on leaves and stems in June; the 1st instar nymphs migrate to the roots, where they cause small galls, which grow with the development of the subsequent nymphal stages (2nd-4th instars); the 4th instar nymphs appear in late August – early September and spend the winter in the root galls; during the following May, these nymphs abandon the galls, climb up to the aerial part of the plant and produce the 5th instar nymphs, which in turn give rise to the new adults.

The gall caused by *C. subpunctata* is a conglomerate of tangled, enlarged and deformed rootlets, which reaches its maximum size (up to about one centimetre) in mid-September, when it contains one to three 4th instar nymphs.




Figure 7. [doi](#)

Overview of forewings of psyllid species in Luxembourg (Psyllidae II).

A *Livilla ulicis* Curtis, 1836 **B** *Psylla alni* (Linnaeus, 1758) **C** *Psylla betulae* (Linnaeus, 1758) **D** *Psylla foersteri* Flor, 1861 **E** *Psylla hartigii* Flor, 1861 **F** *Spanioneura buxi* (Linnaeus, 1758) **G** *Spanioneura fonscolombii* Foerster, 1848.

Rhinocola aceris (Linnaeus, 1758)

(Figs 1, 5, 9, 13)

Findings in Luxembourg. Personal field collection by the authors: Oesling: Marnach (1 ♀, 2.VI.2020, on *Pyrus communis*, occasional plant); West Gutland: Dondelange (1 ♂, 3 ♀♀, 28.V.2020, on *Acer campestre*), Esch-sur-Sûre (13 ♂♂, 24 ♀♀, 15.V.2020, on *A. campestre*; 3 ♂♂, 4 ♀♀, 15.V.2020, on *Acer* sp.), Préitzerdaul (1 ♂, 1 ♀, 20.V.2020, on *Populus tremula*, occasional plant); Moselle: Erpeldange (1 ♀, 7.V.2020, on *Crataegus monogyna*, occasional plant), Mondorf-les-Bains (8 ♂♂, 12 ♀♀, 18.V.2020, on *A. campestre*; 1 ♀, 18.V.2020, on *Salix* sp., occasional plant), Moutfort (3 ♂♂, 3 ♀♀, 1.VI.2020, on *Acer platanoides*), Remich (9 ♂♂, 18 ♀♀, 8.V.2020, on *Acer* sp.).

Material studied in the MNHN collection: East Gutland: Niederanven, Aarnesch (1 ♀, 9.VI - 25.VI.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Widely distributed in the central-western part of the Palaearctic Region, with records from middle Asia, Caucasian Region (Armenia, Georgia), Turkey, nearly all Europe (except Ireland and Iberian Peninsula) and North Africa (Tunisia)

(Hodkinson and White 1979, Burckhardt 1988b, Burckhardt and Lauterer 1989, Ossiannilsson 1992, Burckhardt and Önuçar 1993, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, Gertsson 2010, Lauterer 2011, Malenovský et al. 2011, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, Drohojowska and Kłasa 2019).

Biology. Strictly oligophagous on *Acer* spp. (Sapindaceae). Host plants: *Acer campestre* L., *A. platanoides* L., *A. pseudoplatanus* L., *A. tataricus* L. (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1984, Burckhardt 1988a, Burckhardt and Lauterer 1989, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Lauterer 2001, Burckhardt 2002, Seljak 2006, Ripka 2008, Hodkinson 2009, Malenovský et al. 2011). Adults were also collected on *Betula pendula* Roth (Betulaceae), *Buxus sempervirens* L. (Buxaceae) and *Ulmus* spp. (Ulmaceae) (Burckhardt 1988a, Ripka 2008, Kanturski and Drohojowska 2013) which are “occasional plants”. *Rhinocola aceris* performs one generation per year and, according to Lauterer (1991), overwinters in the egg stage and spends the summer in parapausa on its host plants.




Figure 8. doi

Overview of forewings of psyllid species in Luxembourg (Trioziidae).

A *Bactericera albiventris* (Foerster, 1848) **B** *Bactericera curvattnervis* (Foerster, 1848) **C** *Bactericera substriola* Ossiannilsson, 1992 **D** *Eryngiofaga lautereri* Loginova, 1977 **E** *Lauritrioza alacris* (Flor, 1861) **F** *Trioza abdominalis* Flor, 1861 **G** *Trioza cirsii* Löw, 1881 **H** *Trioza galii* Foerster, 1848 **I** *Trioza remota* Foerster, 1848 **J** *Trioza urticae* (Linnaeus, 1758).

Liviidae

Camarotoscena speciosa (Flor, 1861)

Findings in Luxembourg. This species has not been found by the authors in Luxembourg so far. Nevertheless, findings of its leaf deformations produced on *Populus* sp. and *P. nigra* were reported by Lambinon et al. (2012) (Eastern Gutland: Ettelbruck) and the occurrence of its nymphs in the above galls was described by Burton et al. (2019) (Eastern Gutland: Mersch).

Geographical distribution. Present in almost all of Europe, *C. speciosa* extends its distribution also to the Middle East (Iraq, Turkey) and Central Asia (Caucasian Region, Mongolia, former south European Russia, Tadzhikistan, Turkmenistan, Xinjiang Chinese autonomous region) (Baeva and Kankina 1971, Hodkinson and White 1979, Ossiannilsson 1992, Burckhardt and Önuçar 1993, Conci et al. 1993, Rapisarda 1994, Nokkala 1995, Conci et al. 1996, Burckhardt and Mifsud 2003, Drees 2005, Seljak 2006, Ripka 2008, Mustafa et al. 2014, Serbina et al. 2015, den Bieman et al. 2019).




Figure 9. doi

Overview of female terminalia of psyllid species in Luxembourg (Aphalaridae, Liviidae).

A *Aphałara* sp. gr. *polygoni* Foerster, 1848 **B** *Craspedolepta nebulosa* (Zetterstedt, 1828) **C** *Craspedolepta nervosa* (Foerster, 1848) **D** *Craspedolepta subpunctata* (Foerster, 1848) **E** *Rhinocola aceris* (Linné, 1758) **F** *Livia junci* (Schrank, 1789) **G** *Psyllopsis fraxini* (Linnaeus, 1758) **H** *Psyllopsis fraxinicola* (Foerster, 1848) **I** *Strophingia ericae* (Curtis, 1835).

Biology. Oligophagous on plants of the genus *Populus* L. (amongst which, *P. alba* L., *P. nigra* L. and *P. tremula* L. are important hosts in Europe) (Salicaceae) (Hodkinson and White 1979, Ossiannilsson 1992, Burckhardt and Mifsud 2003, Tomasi 2003, Seljak 2006, Ripka and Kiss 2008, Mustafa et al. 2014), *C. speciosa* performs only one annual generation and overwinters as adults on shelter plants (conifers). Leaf galls produced by this psyllid, consisting in a winding up of the leaf margin (sometimes turning to reddish), are reported in older literature (Houard 1908, Houard 1909, Houard 1913, Ross and Hedicke 1927, Buhr 1964, Buhr 1965); however, *C. speciosa* does not seem to produce significant damage to infested poplar plants.




Figure 10. [doi](#)

Overview of female terminalia of psyllid species in Luxembourg (Psyllidae I).

A *Arytaina genistae* (Latreille, 1804) **B** *Arytainilla spartiophila* (Foerster, 1848) **C** *Cacopsylla ambigua* (Foerster, 1848) **D** *Cacopsylla crataegi* (Schrank, 1801) **E** *Cacopsylla mali* (Schmidberger, 1836) **F** *Cacopsylla melanoneura* (Foerster, 1848) **G** *Cacopsylla nigrita* (Zetterstedt, 1828) **H** *Cacopsylla peregrina* (Foerster, 1848) **I** *Cacopsylla* sp. gr. *pruni* (Scopoli, 1763) **J** *Cacopsylla pulchra* (Zetterstedt, 1838) **K** *Cacopsylla pyri* (Linnaeus, 1758) **L** *Cacopsylla pyricola* (Foerster, 1848) **M** *Cacopsylla pyrisuga* (Foerster, 1848) **N** *Cacopsylla rhamnicola* (Scott, 1876) **O** *Cacopsylla visci* (Curtis, 1835).

***Livia junci* (Schrank, 1789)**

(Figs 1, 5, 9, 13)

Findings in Luxembourg. Material studied in the MNHNL collection: Oesling: Eselborn, Bréichen (2 ♂♂, 2 ♀♀, 24.VIII - 22.IX.2000; 1 ♂, 19.X - 7.XI.2000, Malaise), Hoffelt, Sporbech (4 ♂♂, 7 ♀♀, 22.IX - 19.X.2000, Malaise), Sonlez, Pamer (10 ♂♂, 6 ♀♀, 3.VIII - 24.VIII.2000; 10 ♂♂, 3 ♀♀, 24.VIII - 22.IX.2000; 10 ♂♂, 6 ♀♀, 22.IX - 19.X.2000, Malaise); East Gutland: Wilferdange, Conzefenn (1 ♂, 3 ♀♀, 3.VIII - 24.VIII.2000; 1 ♂, 3 ♀♀, 22.IX - 19.X.2000, Malaise).

No explicit mention exists in literature on previous findings of this species in Luxembourg, though deformations it causes to inflorescences of *Juncus articulatus* L. are reported by Lambinon and Schneider (2004) and by Schneider (2016) (West Gutland: Hollenfels; Moselle: Remerschen). These authors, however, do not report any findings of this psyllid in or nearby the galls. Therefore, the present records represent the first direct observations of *L. junci* in Luxembourg.

Geographical distribution. *Livia junci* is spread all over the Palaearctic Region and has been found in nearly all of Europe, north Africa (Algeria, Morocco), the Middle East (Cyprus, Iran, Lebanon, Turkey) and central, south and eastern Asia (Caucasian Region, India, Kazakhstan, Kyrgyzstan, Primorsky Krai, Siberia, Tadzhikistan, Turkmenistan) (Hodkinson and White 1979, Ossiannilsson 1992, Burckhardt and Lauterer 1993, Burckhardt and Önuçar 1993, Conci et al. 1993, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Hodkinson and Bird 2000, Seljak 2006, Ripka 2008, Malenovský et al. 2011, O'Connor and Malumphy 2011, Burckhardt et al. 2018, den Bieman et al. 2019).

Biology. Oligophagous on many species of the genus *Juncus* L. (Juncaceae) (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988a, Hodkinson 1986, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt 2005, Zeidan-Gèze and Burckhardt 1998, Hodkinson and Bird 2000, Seljak 2006, Ripka 2008, Drohojowska 2009a, Hodkinson 2009, Drohojowska 2009b, Malenovský et al. 2011, Burckhardt et al. 2018). It performs one generation per year and overwinters as adults on shelter plants (conifers). During spring, adults fly back to their host plants, on which they lay eggs. After hatching, nymphs move to the young shoots and start producing characteristic galls by transforming the inflorescences into masses of reddish small leaflets, very close to each other due to the sharp shortening of internodes (Darboux and Houard 1901, Kieffer 1901, Houard 1908, Houard 1909, Houard 1913, Ross and Hedicke 1927, Buhr 1964, Buhr 1965).

***Psyllopsis fraxini* (Linnaeus, 1758)**

(Figs 1, 5, 9, 13)

Findings in Luxembourg. Galls produced by this psyllid on *Fraxinus excelsior* L. are reported by Lambinon and Schneider (2004) (Oesling: Allerborn, Beiler, Biwisch, Clervaux, Hautbellain, Hosingen, Kirchermillen, between Moersdorf and Langsur [sic!], Vianden; Moselle: Ahn, Bous, Deisermillen, Fausermillen, Grevenmacher, Mertert, Mondorf-les-

Bains, Stadbredimus, Wasserbillig; Minette: Bettembourg, Belvaux, Dudelange, Esch-sur-Alzette, Gantenbeinsmühle, Hesperange, Rumelange, Schiff lange, Noertzange, Oberkorn, Pétange, Prënzbierg; West Gutland: Beckerich, Bereldange, Bonnevoie, Bridel, Capellen, Cents, Grass, Haardt, Hamm, Hollerich, Howald, Kleinbettingen, Luxembourg Ville, Pulvermühle, Schleifmühle, Stadtgrund, Steinfort, Steinsel, Walferdange, Windhof; East Gutland: Beringen, Bettendorf, Betzdorf, Bürgerkräiz, Colmar-Berg, Diekirch, Ettelbruck, Gilsdorf, Hagelsdorf, Lintgen, Lorentzweiler, Marxmillen, Manternach, Moesdorf, Neudorf, Niederanven, Oetrange, Roodt-sur-Syre, Rosport, Uebersyren, Wecker, Warken). According to the same authors, the lab rearing from galls collected in some of the above locations always produced adults of *P. fraxini*.




Figure 11. [doi](#)

Overview of female terminalia of psyllid species in Luxembourg (Psyllidae II).

A *Livilla ulicis* Curtis, 1836 **B** *Psylla alni* (Linnaeus, 1758) **C** *Psylla betulae* (Linnaeus, 1758) **D** *Psylla foersteri* Flor, 1861 **E** *Psylla hartigii* Flor, 1861 **F** *Spanioneura buxi* (Linnaeus, 1758) **G** *Spanioneura fonscolombii* Foerster, 1848.

Personal field collection by the authors: Minette: Kayl/Tetange (1 ♂, 2 ♀♀, 23.VII.2019, on *F. excelsior*).

Geographical distribution. Central Asian - European chorotype, diffused from the Himalayan Region (Uttaranchal) westwards to Iran and great parts of Europe (Hodkinson and White 1979, Ossiannilsson 1992, Burckhardt and Lauterer 1993, Conci et al. 1993, Seljak 2006, Ripka 2008, Gertsson 2010, Malenovský et al. 2011, O'Connor and

Malumphy 2011, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, Burckhardt et al. 2018, Drohojowska and Klasa 2019, den Bieman et al. 2019). Introduced to North America (USA; Hodkinson 1988) and the Australian Region (New Zealand and Tasmania; Hollis 2004, Martoni et al. 2018).

Biology. Strictly oligophagous on ashes (*Fraxinus* L., Oleaceae). According to the literature, this psyllid is common especially on *Fraxinus excelsior* L., but can be collected also on *F. americana* L., *F. angustifolia* Vahl, *F. aurea* Willd., *F. mandschurica* Rupr., *F. ornus* L., *F. oxycarpa* Willd., *F. pendula* (Aiton) Hoffmanns, *F. pennsylvanica* Marshall (Hodkinson and White 1979, Burckhardt 1983, Hodkinson 1988, Seljak 2006, Ripka 2008, Burrows 2012). On its host plants, *P. fraxini* performs 1-2 generations per year and overwinters in the egg stage.

This psyllid produces showy galls, widely described in literature (e.g. Darboux and Houard 1901, Houard 1908, Houard 1909, Houard 1913, Ross and Hedicke 1927, Buhr 1964, Buhr 1965, Sampò 1975), consisting in the downrolling of the leaf margin; the rolled up part becomes dilated, thickened, turgescent, conspicuously cross-linked in red and violet.




Figure 12. [doi](#)

Overview of female terminalia of psyllid species in Luxembourg (Triozidae).

A *Bactericera albiventris* (Foerster, 1848) **B** *Bactericera curvattnervis* (Foerster, 1848) **C** *Lauritrioza alacris* (Flor, 1861) **D** *Trioza abdominalalis* Flor, 1861 **E** *Trioza cirsii* Löw, 1881 **F** *Trioza galii* Foerster, 1848 **G** *Trioza remota* Foerster, 1848 **H** *Trioza urticae* (Linnaeus, 1758).

***Psyllopsis fraxinicola* (Foerster, 1848)**

(Figs 1, 5, 9, 13)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Eschdorf (1 ♀♀, 14.VIII.2019, on *Fraxinus excelsior*; 1 ♀♀, 14.VIII.2019, on *Prunus* sp., occasional plant); Minette: Kayl/Tetange (3 ♂♂, 3 ♀♀, 23.VII.2019, on *F. excelsior*); Mosel: Bech/Kleinmacher (1 ♀, 30.VII.2019, on *F. excelsior*), Mondorf-les-Bains (1 ♂, 1 ♀, 1.VIII.2019, on *Populus cinerea*, occasional plant), Remich (1 ♂, 1.VIII.2019, on *F. excelsior*).

New record for Luxembourg.

Geographical distribution. Turanic-European-Mediterranean chorotype, diffused from Central Asia (Kazakhstan) westwards to the Caucasian Region (Armenia, Georgia), Turkey, nearly all of Europe and North Africa (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988a, Burckhardt 1989, Burckhardt and Halperin 1992, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Önuçar 1993, Burckhardt 2005, Seljak 2006, Ripka 2008, Malumphy et al. 2009, Gertsson 2010, Malenovský et al. 2011, O'Connor and Malumphy 2011, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, Holzinger et al. 2017, Spodek et al. 2017, den Bieman et al. 2019). Introduced to the Americas (Canada, USA, South America; Hodkinson 1988, Percy et al. 2012, Castillo Carrillo et al. 2016) and the Australian Region (Australia, New Zealand; Hollis 2004, Percy et al. 2012, Martoni et al. 2018).

Biology. Strictly oligophagous on ashes (*Fraxinus* L., Oleaceae). Especially common on *Fraxinus excelsior* L. According to literature, this psyllid has been collected also on *F. angustifolia* Vahl, *F. dipetala* Hook. & Am., *F. ornus* L., *F. syriaca* Boiss. (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988a, Burckhardt 1989, Hodkinson 1988, Burckhardt 2005, Seljak 2006, Ripka 2008, Drohojowska 2009a, Percy et al. 2012, Spodek et al. 2017), but it is not sure if they are host or casual plants. On its host plants, *P. fraxinicola* performs 1-2 generations per year and overwinters in the egg stage.

***Strophingia ericae* (Curtis, 1835)**

(Figs 1, 5, 9, 13)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Brouch (4 ♀♀, 15.V.2020, on *Calluna vulgaris*), Esch-sur-Sûre (1 ♂, 15.V.2020, on *C. vulgaris*). Material studied in the MNHN collection: Oesling: Lellingen, Op Baerel (1 ♀, 8.VI - 20.VI. 2000, Moericke).

New record for Luxembourg.

Geographical distribution. *Strophingia ericae* is a typical European chorotype and is widespread in almost all Europe, though less common in the Mediterranean part of the continent (Iberian Peninsula, central and south Italy, Balkan Peninsula) (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996,

Burckhardt 2005, Seljak 2006, Gertsson 2010, O'Connor and Malumphy 2011, Serbina et al. 2015, den Bieman et al. 2019).

Biology. Oligophagous on Ericaceae, with *Calluna vulgaris* (L.) Hull as main host plant (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Seljak 2006), but also recorded on *Erica cinerea* L. and *Vaccinium uliginosum* L. (Hodkinson and White 1979, Hodkinson 1981, Burckhardt 2005). Eventual other plants should be considered as occasional hosts. Winter is spent by this psyllid as a nymph on the host plants. In north England, where the biology of *S. ericae* has been thoroughly investigated (Hodkinson 1973a, Hodkinson 1973b, Parkinson and Whittaker 1975), this psyllid develops its cycle in one year at low altitude and in two years at higher altitudes, thus presenting two physiological "races", characterised also by very small morphological differences. Studies on population dynamics of *S. ericae* are reported by Hodkinson (1973b).




Figure 13. doi

Overview of male terminalia of psyllid species in Luxembourg (Aphalaridae, Liviidae).

A *Craspedolepta nebula* (Zetterstedt, 1828) **B** *Craspedolepta subpunctata* (Foerster, 1848)
C *Rhinocola aceris* (Linnaeus, 1758) **D** *Livia junci* (Schrank, 1789) **E** *Psyllopsis fraxini* (Linnaeus, 1758) **F** *Psyllopsis fraxinicola* (Foerster, 1848) **G** *Strophingia ericae* (Curtis, 1835).

Psyllidae

Figs 2, 6, 10

***Arytaina genistae* (Latreille, 1804)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: Oesling: Arsdorf (22 ♂♂, 39 ♀♀, 22.VIII.2019, on *Cytisus scoparius*), Berlé (1 ♂, 2 ♀♀, 20.VIII.2019, on *C. scoparius*); West Gutland: Bertrange (1 ♀, 7.VI.2020, on *C. scoparius*), Bridel (7 ♂♂, 11 ♀♀, 6.VIII.2019, on *C. scoparius*), Brouch (1 ♂, 1 ♀, 6.IX.2019, by general sweeping with net); Minette: Kayl/Tetange (1 ♀, 23.VII.2019, on *C. scoparius*).

Material studied in the MNHN collection: Oesling: Basbellain, Klengelbaach (1 ♀, 24.VIII - 22.IX.2000; 1 ♀, 22.IX - 19.X.2000, Malaise), Eselborn, Bréichen (1 ♀, 3.VIII - 24.VIII.2000, Malaise), Goebelsmühle (3 ♂♂, 2 ♀♀, 13.IV - 27.IV.2000; 2 ♂♂, 5 ♀♀, 25.V - 8.VI.2000; 8 ♂♂, 4 ♀♀, 20.VI - 6.VII.2000; 13 ♂♂, 13 ♀♀, 6.VII - 3.VIII.2000; 1 ♂, 2 ♀♀, 3.VIII - 24.VIII.2000; 4 ♂♂, 1 ♀, 24.VIII - 22.IX.2000; 1 ♂, 2 ♀♀, 22.IX - 19.X.2000, Malaise), Hoscheid, Molberlay (2 ♂♂, 1 ♀, 13.IV - 27.IV.2000; 4 ♂♂, 11 ♀♀, 20.VI - 6.VII.2000; 11 ♂♂, 14 ♀♀, 6.VII - 3.VIII.2000; 4 ♂♂, 8 ♀♀, 3.VIII - 24.VIII.2000; 2 ♂♂, 5 ♀♀, 24.VIII - 22.IX.2000; 1 ♀, 22.IX - 19.X.2000, Malaise), Lellingen, Op Baerel (2 ♂♂, 8.VI - 20.VI.2000; 3 ♂♂, 20.VI - 6.VII.2000, Malaise).

New record for Luxembourg.

Geographical distribution. Widespread in Europe and adventive in the USA, Canada (Nova Scotia) and New Zealand (Hodkinson and White 1979, Ossiannilsson 1992, Hodkinson and Hollis 1987, Hodkinson 1988, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Percy et al. 2012, Serbina et al. 2015, Bleach 2019, den Bieman et al. 2019). Some of the geographical reports need confirmation, due to likely confusion with related species.

Biology. Probably oligophagous on plants of the genus *Cytisus* L. [especially *C. scoparius* (L.) Link] (Fabaceae) (Hodkinson and White 1979, Burckhardt 1983, Hodkinson and Hollis 1987, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt 2005, Seljak 2006, Syrett et al. 2007, Percy et al. 2012); its reports on other Genistinae (e.g. *Chamaecytisus* spp., *Genista tinctoria* L. and *Ulex europaeus* L.) are doubtful or need to be confirmed. This psyllid performs 2-3 (in southern Europe maybe more) generations per year, with an almost continuous development; it spends the winter in all developmental stages (especially as an adult) on its host plants.

***Arytainilla spartiophila* (Foerster, 1848)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: Oesling: Urspeilt (3 ♂♂, 16 ♀♀, 2.VI.2020, on *Cytisus scoparius*); West Gutland: Arsdorf (30 ♂♂, 20 ♀♀, 20.V.2020, on *C. scoparius*; 4 ♀♀, 20.V.2020, by general sweeping), Bertrange (1 ♂, 24 ♀♀,

7.VI.2020, on *C. scoparius*), Brouch (40 ♂♂, 46 ♀♀, 15.V.2020, on *C. scoparius*), Dondelange (1 ♂, 1 ♀, 28.V.2020, by general sweeping), Elvange/Schweich (1 ♀, 28.V.2020, on *Prunus spinosa*, occasional plant; 1 ♂, 28.V.2020, on *Salix viminalis*, occasional plant), Esch-sur-Sûre (11 ♂♂, 12 ♀♀, 15.V.2020, on *C. scoparius*; 4 ♂♂, 7 ♀♀, 15.V.2020, on *P. spinosa*, occasional plant), Goesdorf/Bockholtz (1 ♂, 1 ♀, 20.V.2020, on *Acer pseudoplatanus*, occasional plant; 2 ♂♂, 6 ♀♀, 20.V.2020, on *Sonchus* sp., occasional plant), Hobscheid (4 ♂♂, 6 ♀♀, 28.V.2020, on *C. scoparius*; 1 ♂, 1 ♀, 28.V.2020, on *Malus* sp., occasional plant), Noerdange (1 ♀, 28.V.2020, on *Pyrus communis*, occasional plant), Strassen (19 ♂♂, 4 ♀♀, 28.V.2020, on *C. scoparius*; 3 ♂♂, 8.VI.2020, on *C. scoparius*); Minette: Belvaux (8 ♂♂, 3 ♀♀, 7.V.2020, on *C. scoparius*), Kayl (1 ♀, 20.V.2020, on *Malus* sp., occasional plant); Mosel: Mondorf (1 ♂, 18.V.2020, on *Acer campestre*, occasional plant; 1 ♂, 18.V.2020, on *Crataegus monogyna*, occasional plant).




Figure 14. [doi](#)

Overview of male terminalia of psyllid species in Luxembourg (Psyllidae I).

A *Arytaina genistae* (Latreille, 1804) **B** *Arytainilla spartiophila* (Foerster, 1848) **C** *Cacopsylla affinis* (Löw, 1880) **D** *Cacopsylla ambigua* (Foerster, 1848) **E** *Cacopsylla crataegi* (Schrank, 1801) **F** *Cacopsylla mali* (Schmidberger, 1836) **G** *Cacopsylla melanoneura* (Foerster, 1848) **H** *Cacopsylla peregrina* (Foerster, 1848) **I** *Cacopsylla* sp. gr. *pruni* (Scopoli, 1763) **J** *Cacopsylla pulchra* (Zetterstedt, 1838) **K** *Cacopsylla pyri* (Linnaeus, 1758) **L** *Cacopsylla pyricola* (Foerster, 1848) **M** *Cacopsylla pyrisuga* (Foerster, 1848).

Material studied in the MNHN collection: Oesling: Goebelsmühle (313 ♂♂, 594 ♀♀, 11.V - 25.V.2000; 384 ♂♂, 611 ♀♀, 1 nymph, 25.V - 8.VI.2000; 5 ♂♂, 6 ♀♀, 20.VI - 6.VII.2000; 4 ♂♂, 5 ♀♀, 6.VII - 3.VIII.2000; 1 ♂, 1 ♀, 24.VIII - 22.IX.2000; 1 ♂, 22.IX - 19.X.2000, Malaise), Hoscheid, Molberlay (1 ♀, 20.VI - 6.VII.2000, Malaise), Lellingen, Op Baerel (88 ♂♂, 87 ♀♀, 25.V - 8.VI.2000; 25 ♂♂, 33 ♀♀, 8.VI - 20.VI.2000; 2 ♂♂, 20.VI - 6.VII.2000, Malaise); West Gutland: Capellen, Werwelslach (1 ♂, 2 ♀♀, 27.V - 9.VI.1999, Malaise); East Gutland: Godbrange, Schléidelbierg (3 ♂♂, 7 ♀♀, 27.V - 9.VI.1999, Moericke), Niederanven, Aarnesch (3 ♂♂, 2 ♀♀, 27.V - 9.VI.1999, Malaise); Minette: Schiffange, Kayl, Brucherbierg (3 ♂♂, 5 ♀♀, 27.V - 9.VI.2000, Malaise); Moselle: Canach, Wéngertsbierg (6 ♀♀, 27.V - 9.VI.1999; 1 ♀, 9.VI - 25.VI.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Widespread in central-western and southern Europe (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Malumphy et al. 2009, O'Connor and Malumphy 2011, den Bieman et al. 2019). Introduced to North America (Canada, USA), Australia and New Zealand (Hodkinson 1988, Percy et al. 2012).

Biology. *Arytainilla spartiophila* is monophagous on *Cytisus scoparius* (L.) Link (Fabaceae), on which it spends its entire life cycle, performing a single generation per year and overwintering as egg. In the Northern Hemisphere, adults start to fly in mid-April and occur on the plants till the first half of June; from the second half of May, males start to decline in number and populations of this psyllid become female-biased (Wheeler 2017). For its exclusive monophagy on *C. scoparius* and the large populations, this psyllid can build up on Scotch broom in many European countries (especially in Great Britain) causing substantial damage; it has been artificially introduced and released for biological control in exotic habitats (such as California and New Zealand) where its host plant became invasive (Syrett et al. 2007, Hogg et al. 2015).

***Cacopsylla affinis* (Löw, 1880)**

(Figs 2, 6, 14)

Findings in Luxembourg. Material studied in the MNHN collection: Oesling: Lellingen, Op Baerel (1 ♂, 25.V - 8.VI.2000, Malaise).

In the absence of male specimens, it is impossible to morphologically distinguish this species from *Cacopsylla melanoneura* (Foerster); for this reason, collections in Luxembourg of female specimens, here attributed to the more common species *C. melanoneura*, could also refer to *C. affinis*.

New record for Luxembourg.

Geographical distribution. Distributed in most of Europe, eastwards to Turkey and the Caucasian Region (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Önuçar 1993, Conci et al. 1996,

Seljak 2006, Tedeschi et al. 2008, Ripka 2008, Ripka 2009, Malenovský et al. 2011, den Bieman et al. 2019).

Biology. Oligophagous on hawthorns (*Crataegus* spp.) (Rosaceae) (Burckhardt 1983, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Hodkinson 2009, Malenovský et al. 2011), *C. affinis* performs one generation per year and overwinters as adult on shelter plants, especially conifers and Fagaceae.

Economic significance. A controversial phytosanitary importance is attributed to this species as potential vector of phytopathogenic microorganisms, especially '*Candidatus Liberibacter europaeus*' (Tedeschi et al. 2009, Camerota et al. 2012) which should be considered, however, as an endophyte rather than a real pathogen (Raddadi et al. 2011).




Figure 15. doi

Overview of male terminalia of psyllid species in Luxembourg (Psyllidae II).

A *Livilla ulicis* Curtis, 1836 **B** *Psylla alni* (Linnaeus, 1758) **C** *Psylla betulae* (Linnaeus, 1758) **D** *Psylla foersteri* Flor, 1861 **E** *Psylla hartigii* Flor, 1861 **F** *Spanioneura buxi* (Linnaeus, 1758) **G** *Spanioneura fonscolombii* Foerster, 1848.

Cacopsylla ambigua (Foerster, 1848)

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Brouch (3 ♂♂, 8 ♀♀, 15.V.2020, on *Salix caprea*; 1 ♀, 15.V.2020, on *Salix viminalis*), Elvange/Schweich (2 ♂♂, 28.V.2020, on *S. viminalis*), Strassen (3 ♂♂, 5 ♀♀, 27.IV.2020, on *S. caprea*); East Gutland: Rodenburg (1 ♀, 27.V.2020, on *S. caprea*).

Material studied in the MNHN collection: East Gutland: Wilferdange, Conzefenn (1 ♀, 6.VII - 3.VIII.2000, Malaise).

New record for Luxembourg.

Geographical distribution. *Cacopsylla ambigua* is an Eurasian chorotype, especially having a wide distribution in Europe (Hodkinson and White 1979, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Lauterer and Burckhardt 1997, Labina 2006, Seljak 2006, Labina 2008, Ripka 2008, Inoue 2010, O'Connor and Malumphy 2011, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, Zendedel et al. 2016, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Strictly oligophagous on *Salix* spp. (Salicaceae), being reported in literature from *S. alba* L., *S. atrocinerea* Brot., *S. aurita* L., *S. caprea* L., *S. cinerea* L., *S. elaeagnos* Scop., *S. incana* Schrank, *S. lapponum* L., *S. purpurea* L. and *S. viminalis* L. (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Lauterer and Burckhardt 1997). On its host plants, it performs the entire developmental cycle, showing one or two generations per year and overwintering as egg or, according to Lauterer (1976), as 1st or 2nd instar nymph.

Cacopsylla crataegi (Schrank, 1801)

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Strassen (1 ♂, 2 ♀♀, 27.IV.2020, on *Crataegus monogyna*).

Material studied in the MNHN collection: West Gutland: Capellen, Werwelslach (1 ♀, 18.V - 27.V.1999, Moericke); East Gutland: Godbrange, Schléidelbierg (4 ♂♂, 1 ♀, 6.IV - 22.IV.1999, Malaise); Minette: Niedercorn, Giele Botter (2 ♂♂, 25.VI - 12.VII.1999, Malaise); Moselle: Canach, Wéngertsbierg (1 ♂, 1 ♀, 22.IV - 11.V.1999; 1 ♂, 11.V - 27.V.1999; 1 ♂, 27.V - 9.VI.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Widely distributed in the Palaearctic Region, from central-south Asia (west Himalayan Region, India, Iran, Caucasian Region) to nearly all parts of Europe and North Africa (Algeria, Morocco) (Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Lauterer 1993,

Conci et al. 1996, Seljak 2006, Ripka 2008, Ripka and Kiss 2008, Gertsson 2010, Malenovský et al. 2011, O'Connor and Malumphy 2011, Kanturski and Drohojowska 2013, Oettl and Schlink 2015, Ouvrard et al. 2015, Serbina et al. 2015, Burckhardt et al. 2018, Drohojowska and Klasa 2019, den Bieman et al. 2019).




Figure 16. doi

Overview of male terminalia of psyllid species in Luxembourg (Trioziidae).

A *Bactericera substriola* Ossiannilsson, 1992 **B** *Eryngiofaga lautereri* Loginova, 1977 **C** *Trioza abdominalis* Flor, 1861 **D** *Trioza cirsii* Löw, 1881 **E** *Trioza galii* Foerster, 1848 **F** *Trioza remota* Foerster, 1848 **G** *Trioza urticae* (Linnaeus, 1758).

Biology. Strictly oligophagous on hawthorns (*Crataegus* spp.) (Rosaceae), *C. crataegi* is mainly reported from *Crataegus monogyna* Jacq. and *C. oxyacantha* L. (Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996), being found also on other species, especially in Asia, such as *C. coccinea* L. or *C. pentagyna* Waldst. & Kit. ex Willd. (Ossiannilsson 1992). Besides its host plants and various occasional species, this psyllid can be found frequently on other Rosaceae, such as *Malus* spp., *Mespilus* spp. or *Sorbus* spp. (Burckhardt 1989,

Ripka 2008, Oettl and Schlink 2015). *Cacopsylla crataegi* performs only one generation per year and overwinters as an adult on shelter plants (conifers).

***Cacopsylla mali* (Schmidberger, 1836)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Dondelange (4 ♂♂, 2 ♀♀, 28.V.2020, on *Malus* sp.); Minette: Kayl (3 ♂♂, 6 ♀♀, 20.V.2020, on *Malus* sp.).

No explicit mention exists in literature on findings of this insect in Luxembourg, though the leaf deformations it produces on *Malus* sp. are reported by Lambinon and Schneider (2004) (West Gutland: Kirchberg). The authors, however, do not report any finding of this psyllid in or nearby the galls. Therefore, the present record represents the first direct observation of *C. mali* in Luxembourg.

Geographical distribution. *Cacopsylla mali* is widely distributed in the Eurasian Region: though not properly recorded in central Asia, it seems to occur from the Russian Far East, Japan and the Korean Peninsula to most of Europe (Hodkinson and White 1979, Burckhardt 1983, Kwon 1983, Burckhardt 1988b, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Önuçar 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, Malumphy et al. 2009, Inoue 2010, Lauterer 2011, Malenovský et al. 2011, O'Connor and Malumphy 2011, Kanturski and Drohojowska 2013, Serbina et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019). It is known also from the Nearctic (USA, Canada) and Afrotropical (South Africa) Regions, as well as from Australia (Burckhardt 1988b, Hodkinson 1988, Burckhardt 1994a, Wheeler and Hoebeke 2005, Inoue 2010), where it likely has been introduced.

Biology. Strictly oligophagous on various species of the genus *Malus* Mill. (Rosaceae). In Europe, it can be found mainly on *M. domestica* Borkh. and *M. sylvestris* Mill. (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Burckhardt 1994a, Conci et al. 1996, Ripka 2008, Inoue 2010). *Malus asiatica* Nakai, *M. baccata* (L.) Borkh. and *M. transitoria* (Batalin) C.K. Schneid. are reported as host plants of this psyllid in Asia (Kwon 1983, Li 2011). *Cacopsylla mali* performs one generation per year and overwinters as egg on its host plants. It often causes leaf alterations, corrugations and distortions (Buhr 1964, Buhr 1965). Similar to other species of the genus *Cacopsylla*, it may show a typical summer migration: part of the adults, which fed on the host plant for 2-3 weeks after emergence, migrate to other species of trees or shrubs, to re-immigrate to the host plants in September. Adults show a summer reproductive "parapause", which ends in autumn with a reactivation with their oogenesis (Lauterer 1991).

Economic significance. *Cacopsylla mali* is a secondary pest of apple trees in central Europe. In spite of old reports as a very harmful species, the damage it causes to crops is usually negligible. In recent studies, 'Candidatus Phytoplasma mali', the etiological agent of the Apple Proliferation (AP) disease, has been detected also in various psyllid species, including *C. mali*, different from the two known vectors of this pathogen [*Cacopsylla*

melanoneura (Foerster) and *C. picta* (Foerster)] (Miñarro et al. 2016); yet the real potential of *C. mali* to transmit the disease is still unclear and warrants further investigation.

***Cacopsylla melanoneura* (Foerster, 1848)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: Oesling: Marnach (9 ♂♂, 6 ♀♀, 2.VI.2020, on *Crataegus monogyna*); West Gutland: Arsdorf (1 ♂, 7 ♀♀, 1 nymph, 20.V.2020, on *C. monogyna*), Dondelange (4 ♂♂, 2 ♀♀, 28.V.2020, on *C. monogyna*), Elvange/Schweich (2 ♂♂, 2 ♀♀, 28.V.2020, on *C. monogyna*; 4 ♀♀, 28.V.2020, on *Prunus spinosa*, occasional plant), Esch-sur-Sûre (1 ♂, 1 ♀, 15.V.2020, on *Acer* sp., occasional plant), Préitzerdaul (1 ♂, 20.V.2020, on *C. monogyna*), Strassen (1 ♂, 3 ♀♀, 27.IV.2020, on *C. monogyna*), Useldange (1 ♂, 1 ♀, 28.V.2020, on *C. monogyna*); East Gutland: Bettendorf (1 ♀, 27.V.2020, by general sweeping with net, from *Salix* sp. and *Populus* sp., occasional plants); Moselle: Elvange/Burmerange (3 ♂♂, 5 ♀♀, 19.V.2020, on *C. monogyna*; 1 ♀, 19.V.2020, by general sweeping with net), Erpeldange (6 ♂♂, 3 ♀♀, 7.V.2020, on *C. monogyna*), Mondorf (4 ♂♂, 5 ♀♀, 18.V.2020, on *C. monogyna*; 1 ♀, 28.V.2020, on *Acer campestre*, occasional plant; 1 ♀, 28.V.2020, by general sweeping with net).

Material studied in the MNHN collection: Oesling: Basbellain, Klengelbaach (1 ♀, 22.IX - 19.X.2000, Malaise), Lellingen, Op Baerel (3 ♂♂, 8 ♀♀, 25.V - 8.VI.2000, Malaise); West Gutland: Bertrange, Brill (8 ♂♂, 14 ♀♀, 15.III - 19.III.1999, Moericke), Capellen, Werwelslach (2 ♂♂, 1 ♀, 13.IV - 22.IV.1999; 1 ♀, 22.IV - 28.IV.1999; 2 ♀♀, 18.V - 27.V.1999; 1 ♂, 2 ♀♀, 27.V - 9.VI.1999, Malaise); East Gutland: Godbrange, Schléidelbierg (44 ♂♂, 37 ♀♀, 6.IV - 22.IV.1999; 10 ♂♂, 2 ♀♀, 22.IV - 11.V.1999; 8 ♂♂, 11 ♀♀, 18.V - 27.V.1999; 23 ♂♂, 23 ♀♀, 27.V - 9.VI.1999; 2 ♂♂, 5 ♀♀, 9.VI - 25.VI.1999; 1 ♂, 25.VI - 8.VII.1999, Malaise), Koedange, Poenn (1 ♀, 11.III - 15.III.1999, Moericke), Niederanven, Aarnesch (5 ♂♂, 4 ♀♀, 22.IV - 11.V.1999; 2 ♀♀, 11.V - 27.V.1999; 1 ♀, 9.VI - 25.VI.1999, Malaise); Minette: Kockelscheuer, Conter Jans Boesch (1 ♀, 22.IV - 28.IV.1999, Malaise), Niedercorn, Giele Botter (2 ♂♂, 25.VI - 12.VII.1999, Malaise), Schiffange, Kayl, Brucherbierg (2 ♂♂, 3 ♀♀, 22.IV - 11.V.1999; 1 ♂, 1 ♀, 11.V - 27.V.1999; 2 ♂♂, 2 ♀♀, 27.V - 9.VI.1999, Malaise); Moselle: Canach, Wéngertsbierg (6 ♂♂, 5 ♀♀, 22.IV - 11.V.1999; 3 ♀♀, 11.V - 27.V.1999, Malaise).

In the absence of male specimens, it is impossible to morphologically distinguish this species from *Cacopsylla affinis* (Löw); for this reason, collections in Luxembourg of female specimens here attributed to *C. melanoneura* could refer to *C. affinis*. Identification tools of the two species by molecular methods have been studied by Tedeschi and Nardi (2010).

No explicit mention exists in literature on findings of *C. melanoneura* in Luxembourg, though its leaf deformations produced on *Crataegus monogyna* Jacq. are reported by Lambinon and Schneider (2004) (Minette: Bettembourg; Moselle: Canach; West Gutland: Bonnevoie, Kleinbettingen, Steinfort; East Gutland: Oberanven), who do not report, however, any finding of this psyllid in or nearby the galls. Therefore, the present records are the first direct findings of *C. melanoneura* in Luxembourg.

Geographical distribution. Palearctic chorotype, widespread and common from the Far East Asia (Japan) to almost all Europe and North Africa (Algeria), through various confirmed reports from Central Asia [Mongolia, Russia (Irkutsk and Siberia), Tadzhikistan, Uzbekistan] (Loginova 1968, Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Lauterer 1999, Baugnée et al. 2002, Seljak 2006, Ripka 2008, Malumphy et al. 2009, Gertsson 2010, Malenovský et al. 2011, O'Connor and Malumphy 2011, Chireceanu and Fătu 2012, Ouvrard et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Oligophagous on many Rosaceae, *C. melanoneura* primarily lives on plant species of the genus *Crataegus* L. (especially frequent on *C. monogyna* Jacq. and *C. oxyacantha* L., but also reported on *C. laevigata* (Poir.) DC. and *C. maximowiczii* C.K. Schneid. (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988a, Burckhardt 1989, Ossiannilsson 1992, Seljak 2006, Ripka 2008, Hodkinson 2009, Malenovský et al. 2011). It can live even on *Malus* spp. (*M. communis* Desf., *M. domestica* Borck.), *Mespilus* spp. (*M. germanica* L.), *Prunus* spp. (*P. armeniaca* L.) and *Pyrus* spp. (*Pyrus communis* L.) (Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, Hodkinson 2009, Malumphy et al. 2009). Furthermore, it can frequently be found on other occasional plants. It shows only one generation per year and overwinters in the adult stage on shelter plants (conifers); aestivation and overwintering habits of this species have been studied in detail by Pizzinat et al. (2011).

Economic significance. On hawthorn *C. melanoneura* is not a harmful species, if leaf deformations often caused by this insect are excluded, i.e. yellow to blood-red folds of the leaf margins. Nevertheless, phytosanitary importance of this species has recently been highlighted, for its ability to host and transmit phytopathogenic microorganisms of various cultivated Rosaceae (especially apple trees), such as '*Candidatus Liberibacter europaeus*' and '*Candidatus Phytoplasma mali*', the latter being the causative agent of the Apple Proliferation (AP) disease (Mayer et al. 2009, Tedeschi et al. 2009, Tedeschi et al. 2012, Camerota et al. 2012, Jarausch et al. 2012, Oettl and Schlink 2015, Kaya et al. 2016, Miñarro et al. 2016, Oppedisano et al. 2020)

Given the phytosanitary importance of this species as a potential vector of phytopathogenic microorganisms, its actual distribution and pathogenicity in Luxembourg should be further investigated.

***Cacopsylla nigrita* (Zetterstedt, 1828)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Material studied in the MNHN collection: East Gutland: Wilferdange, Conzefenn (1 ♀, 22.IX - 19.X.2000, Malaise).

New record for Luxembourg.

Geographical distribution. Widely distributed in Europe, except its most western (e.g. Great Britain, France and Iberian Peninsula) and Mediterranean parts and reported also from the

Caucasian Region and eastwards to Japan (Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Lauterer and Burckhardt 1997, Seljak et al. 2008).

Biology. Strictly oligophagous on *Salix* spp. (Salicaceae), *C. nigrita* is reported especially on *S. caprea* L., *S. elaeagnos* Scop., *S. foetida* Schleicher, *S. helvetica* Vill., *S. lapponum* L., *S. myrsinifolia* Salisb., *S. phylicifolia* L., *S. purpurea* L. and *S. waldsteniana* Willd. (Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Ripka 2008, Seljak et al. 2008). It performs only one generation per year and overwinters as an adult on shelter plants (conifers).

***Cacopsylla peregrina* (Foerster, 1848)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: Oesling: Marnach (11 ♂♂, 17 ♀♀, 2.VI.2020, on *Crataegus monogyna*), Wahl (2 ♂♂, 2 ♀♀, 14.VIII.2019, on *C. monogyna*); West Gutland: Arsdorf (2 ♀♀, 20.V.2020, on *C. monogyna*), Dondelange (3 ♀♀, 28.V.2020, on *C. monogyna*; 1 ♂, 1 ♀, 28.V.2020, on *Alnus glutinosa*, occasional plant), Elvange/Schweich (7 ♂♂, 7 ♀♀, 28.V.2020, on *C. monogyna*), Esch-sur-Sûre (1 ♀, 15.V.2020, on *C. monogyna*; 1 ♀, 15.V.2020, on *Acer* sp., occasional plant), Strassen (4 ♀♀, 27.IV.2020, on *C. monogyna*), Useldange (2 ♂♂, 2 ♀♀, 28.V.2020, on *C. monogyna*); Minette: Belvaux (9 ♂♂, 2 ♀♀, 8.VIII.2019, on *C. monogyna*; 1 ♂, 8.VIII.2019, on *Euonymus* sp., occasional plant), Dudelange (5 ♂♂, 3 ♀♀, 23.VII.2019, on *C. monogyna*), Kayl (4 ♂♂, 7 ♀♀, 11.VI.2020, on *Malus* sp., occasional plant), Kayl/Tetange (14 ♂♂, 10 ♀♀, 23.VII.2019, on *C. monogyna*), Vesquenhaff (7 ♂♂, 17 ♀♀, 30.VII.2019, on *C. monogyna*); Moselle: Elvange/Burmerange (8 ♂♂, 2 ♀♀, 19.V.2020, on *C. monogyna*), Erpeldange (15 ♂♂, 13 ♀♀, 3 nymphs, 7.V.2020, on *C. monogyna*), Mondorf (4 ♂♂, 3 ♀♀, 18.V.2020, on *C. monogyna*; 2 ♀♀, 18.V.2020, on *Acer campestre*, occasional plant), Moutfort (1 ♀, 1.VI.2020, on *Acer platanoides*, occasional plant), Remerschen (1 ♀, 28.V.2020, on *Alnus glutinosa*, occasional plant).

Material studied in the MNHN collection: Oesling: Lellingen, Op Baerel (2 ♂♂, 8.VI - 20.VI.2000, Malaise); East Gutland: Godbrange, Schléidelbierg (7 ♂♂, 4 ♀♀, 18.V - 27.V.1999; 9 ♂♂, 4 ♀♀, 27.V - 9.VI.1999; 14 ♂♂, 10 ♀♀, 9.VI - 25.VI.1999; 1 ♂, 2 ♀♀, 25.VI - 8.VII.1999; 4 ♂♂, 4 ♀♀, 8.VII - 22.VII.1999; 3 ♂♂, 1 ♀, 22.VII - 5.VIII.1999; 2 ♂♂, 5.VIII - 19.VIII.1999; 1 ♀, 19.VIII - 2.IX.1999; 1 ♂, 2 ♀♀, 2.IX - 27.IX.1999, Malaise), Niederanven, Aarnesch (1 ♀, 11.V - 27.V.1999; 2 ♀♀, 27.V - 9.VI.1999; 3 ♂♂, 1 ♀, 9.VI - 25.VI.1999; 1 ♀, 5.VIII - 19.VIII.1999; 2 ♀♀, 2.IX - 27.IX.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Widely distributed in the Palaearctic Region, where it is recorded from Japan and Far East Asia (Inoue 2010) and westwards through the temperate areas of Asia (Burckhardt 2005) and Turkey (Burckhardt and Önuçar 1993), to nearly all Europe and North Africa (Hodkinson and White 1979, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt 2005, Seljak 2006, Ripka 2008, Malumphy et al. 2009, Gertsson 2010, Inoue 2010, Lauterer 2011, Malenovský et al. 2011, O'Connor

and Malumphy 2011, Kanturski and Drohojowska 2013, Serbina et al. 2015, Drohojowska and Kłasa 2019, den Bieman et al. 2019). Recently recorded also from North America (Canada and USA; Wheeler and Stoops 2001, Wheeler and Hoebeke 2005), where it has been introduced.

Biology. Strictly oligophagous on hawthorns (*Crataegus* L., Rosaceae) (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt 2005, Wheeler and Hoebeke 2005, Seljak 2006, Ripka 2008, Hodkinson 2009, Malumphy et al. 2009, Inoue 2010, Lauterer 2011, Malenovský et al. 2011). In Europe, preferably found on *Crataegus monogyna* Jacq. and *C. oxyacantha* L.; also frequent on other hawthorn species, such as on *C. arnoldiana* Sarg., *C. laevigata* (Poir.) DC or *C. maximowiczii* C.K. Schneid., especially in other geographical areas. On its host plants, this insect performs a single generation per year, with long-living adults during summer, when they frequently spread around also on occasional plants. Eggs are laid in late summer on the host plants (hawthorns) and overwinter. In recent studies, *C. peregrina* showed to be moderately associated with '*Candidatus Liberibacter europaeus*' (Tedeschi et al. 2009, Camerota et al. 2012), a phloem-limited Gram-negative bacterium infecting pear plants though producing no specific symptoms; therefore, thought to be an endophyte rather than a pathogen (Raddadi et al. 2011).

***Cacopsylla* sp. gr. *pruni* (Scopoli, 1763)**

(Figs 2, 6, 10, 14)

Two cryptic species, formerly considered as biotypes A and B, have been evidenced within *C. pruni*. They are ecologically and morphologically indistinguishable and partly overlap in their distributions (Sauvion et al. 2021), but show clear genetic differentiation with no hybrids detected (Sauvion et al. 2007, Peccoud et al. 2013, Peccoud et al. 2018). The identity of the material collected in Luxembourg still needs to be checked by molecular methods.

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Arsdorf (1 ♂, 20.V.2020, on *Prunus spinosa*), Strassen (1 ♀, 27.IV.2020, on *P. spinosa*); Moselle: Elvange/Burmerange (1 ♀, 19.V.2020, on *P. spinosa*).

Material studied in the MNHN collection: Oesling: Goebelsmühle (1 ♀, 25.V - 8.VI.2000, Malaise), Lellingen, Op Baerel (1 ♀, 25.V - 8.VI.2000; 1 ♀, 8.VI - 20.VI.2000; 2 ♂♂, 1 ♀, 20.VI - 6.VII.2000, Malaise); West Gutland: Capellen, Werwelslach (2 ♂♂, 7 ♀♀, 13.IV - 22.IV.1999; 3 ♀♀, 22.IV - 28.IV.1999; 6 ♂♂, 12 ♀♀, 11.V - 18.V.1999; 1 ♀, 18.V - 27.V.1999; 1 ♂, 1 ♀, 27.V - 9.VI.1999, Malaise); East Gutland: Godbrange, Schléidelbierg (1 ♂, 1 ♀, 6.IV - 22.IV.1999; 2 ♀♀, 11.V - 18.V.1999; 2 ♀♀, 18.V - 27.V.1999, Malaise), Koedange, Poenn (1 ♀, 22.VII - 5.VIII.1999, Malaise), Niederanven, Aarnesch (1 ♂, 1 ♀, 22.IV - 11.V.1999, Malaise); Minette: Schifflange, Kayl, Brucherbierg (1 ♀, 22.IV - 11.V.1999, Malaise); Moselle: Canach, Wéngertsbierg (2 ♂♂, 1 ♀, 22.IV - 11.V.1999; 2 ♂♂, 1 ♀♀, 11.V - 27.V.1999; 1 ♀, 25.VI - 8.VII.1999, 1 ♂, 5.VIII - 19.VIII.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Central Asian-European chorotype; *C. pruni* has been recorded from the Mongolian Region (Irkutsk) westwards to the Iran, Caucasian Region (Azerbaijan, Georgia), Turkey and to nearly all of Europe (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Burckhardt and Lauterer 1993, Conci et al. 1993, Conci et al. 1996, Burckhardt 2005, Seljak 2006, Sauvion et al. 2007, Ripka 2008, Gertsson 2010, Lauterer 2011, Malenovský et al. 2011, O'Connor and Malumphy 2011, Steffek et al. 2012, Kanturski and Drohojowska 2013, Drohojowska and Burckhardt 2014, Ouvrard et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Though its adults have been found by chance also on occasional plants, such as *Crataegus* spp. or *Malus domestica* Borckh. (Oettl and Schlink 2015), *C. pruni* is strictly oligophagous on plants of the genus *Prunus* L. (Rosaceae), with *Prunus spinosa* L. being the most frequent host plant, but it can also be observed on *P. armeniaca* L., *P. avium* (L.) L., *P. cerasifera* Ehrh., *P. domestica* L., *P. insititia* L., *P. padus* L., *P. persica* (L.) Stokes and *P. salicina* Lindl. (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt 2005, Seljak 2006, Sauvion et al. 2007, Ripka 2008, Hodkinson 2009, Lauterer 2011, Malenovský et al. 2011, Steffek et al. 2012, Jarausch and Jarausch 2016). This psyllid species performs one generation per year and overwinters as an adult on shelter plants (conifers). Recent studies allowed us to understand the feeding behaviour on its winter shelter plants, through the application of electrical penetration graph (EPG) recordings and survival bioassays on different conifer species, as well as the analysis of chemical composition of their plant sap (Gallinger and Gross 2018).

Economic significance. Over the past few years, *C. pruni* has taken on a remarkable phytosanitary significance, for its ability to transmit '*Candidatus Phytoplasma prunorum*' (Carraro et al. 1998, Jarausch et al. 2001, Carraro et al. 2004), the causing agent of a complex of economically important disorders on *Prunus* plants (including cultivated *P. armeniaca*, *P. domestica* and *P. persica*), which are collectively referred to as European Stone Fruit Yellows (ESFY). Recently, it has been demonstrated how both putative species of the *C. pruni* complex can transmit and spread the pathogen (Marie-Jeanne et al. 2020). Over the past two decades, ESFY has been detected in most southern and central European countries, Middle East and North Africa (Sertkaya et al. 2005, Marcone et al. 2010, Ben Khalifa et al. 2011, Cieślińska 2011, Tedeschi et al. 2013, Allahverdi et al. 2014, Valasevich and Schneider 2016, Warabieda et al. 2017, Andrianjaka-Camps et al. 2018, Jarausch et al. 2019, Riedle-Bauer et al. 2019b), thus being one of the most serious pests in European stone fruit production and a potential threat for fruit crops worldwide, causing important economic damage.

Given its phytosanitary importance as a potential vector of serious phytopathogenic microorganisms, it would be worth monitoring the distribution and pathogenicity of the separate taxonomic entities within the *Cacopsylla pruni* complex in Luxembourg.

***Cacopsylla pulchra* (Zetterstedt, 1838)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: East Gutland: Rodenburg (1 ♂, 2 ♀♀, 27.V.2020, on *Salix caprea*); Moselle: Mondorf (1 ♂, 2 ♀♀, 18.V.2020, on *Salix* sp.).

Material studied in the MNHN collection: Minette: Kockelscheuer, Conter Jans Boesch (1 ♀, 13.IV - 22.IV.1999, Malaise), Schifflange, Kayl, Brucherberg (1 ♂, 11.V - 27.V.1999, Malaise).

New record for Luxembourg.

Geographical distribution. *Cacopsylla pulchra* is a Eurasian chorotype, widely distributed in Europe (except the southern Balkan Peninsula) and recorded also from the Caucasian Region and eastwards to Central (Mongolia) and East Asia (Far East Russia, Japan, Korean Peninsula) (Hodkinson and White 1979, Kwon 1983, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Lauterer and Burckhardt 1997, Seljak 2006, Ripka 2008, Gertsson 2010, Inoue 2010, Malenovský et al. 2011, O'Connor and Malumphy 2011, Ouvrard et al. 2015, Serbina et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Strictly oligophagous on *Salix* spp. (Salicaceae), *C. pulchra* has been recorded on many species of willows, such as *S. apennina* Skvortsov, *S. atrocinerea* Brot., *S. caprea* L., *S. cinerea* L., *S. elaeagnos* Scop., *S. foetida* Schleicher, *S. gracilistyla* Miq., *S. integra* Thunb., *S. myrsinifolia* Salisb., *S. pentandra* L., *S. purpurea* L., *S. repens* L. and *S. viminalis* L. (Hodkinson and White 1979, Burckhardt 1983, Kwon 1983, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Park 1996, Seljak 2006, Malenovský et al. 2011, Cho et al. 2017a). It performs one generation per year and overwinters as an adult on shelter plants (conifers).

***Cacopsylla pyri* (Linnaeus, 1758)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: Moselle: Burmerange (6 ♂♂, 8 ♀♀, 8.V.2020, on *Pyrus communis*), Elvange/Burmerange (2 ♂♂, 3 ♀♀, 19.V.2020, on *P. communis*).

New record for Luxembourg.

Geographical distribution. *Cacopsylla pyri* is common throughout Europe, especially in the central-southern fruit growing areas of the continent and is recorded also from the Caucasian Region (Armenia, Georgia), Middle East (Iran), central Asia (Kazakhstan) eastwards up to China (Xinjiang) (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Lauterer 1993, Burckhardt and Önuçar 1993, Conci et al. 1996, Aguiar and Martin 2001, Agusti et al. 2003, Seljak 2006, Ripka 2008, O'Connor and Malumphy 2011, Sanchez and Ortín-Angulo 2011, Sanchez and Ortín-Angulo 2012, Drohojowska and Burckhardt 2014, Bufaur and

Harizanova 2015, Oettl and Schlink 2015, Pekár et al. 2015, Serbina et al. 2015, Cho et al. 2017b, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Oligophagous on plants of the genus *Pyrus* L., mostly *Pyrus communis* L. (Rosaceae) (Hodkinson and White 1979, Burckhardt 1983, Burckhardt and Hodkinson 1986, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Burckhardt 1994a, Conci et al. 1996, Aguiar and Martin 2001, Seljak 2006, Ripka 2008, Hodkinson 2009, Li 2011, Sanchez and Ortín-Angulo 2011, Civolani et al. 2013). According to Cravedi et al. (2008), it performs up to 5-7 generations per year, overwintering as adults (especially females in ovarian diapause) in bark crevices of the same pear trees, but also on other occasional plants. Egg deployment starts in early spring, with the first sunny days (2-3 consecutive days with maximum temperatures over 10°C), initially within the cracks of the twigs, then in those of smaller branches and afterwards, as soon as trees start to vegetate, on perules, petioles and leaflets. Newly-emerged nymphs tend to settle during this spring period on the newly-spread leaflets and around the calycin zone of the newly-formed small fruits. Later in the season, adults of the summer generations lay eggs on tender shoots, along the main vein of the underside of the leaves, where nymphs tend to remain and produce a huge amount of honeydew. Late summer and autumn generations have normally a reduced fertility; and population levels are usually reduced with consequent limited harm, although late summer pullulations may also occur (especially in warmer climates).

Economic significance. *Cacopsylla pyri* is the most common psyllid species on all cultivated pear varieties in western Europe and regionally often considered a key pest. Nymphs of the summer generation produce a large amount of honeydew, damaging both plants and fruits. In the case of heavy infestations, the honeydew can smear all green parts of the plant, reducing photosynthesis and respiration, as well as making fruits suffer a significant commercial depreciation. *Cacopsylla pyri* can also be dangerous, especially to young plants, as a vector of '*Candidatus Phytoplasma pyri*', the agent of Pear Decline disease, which is transmitted in Europe by both nymphs and adults of this psyllid (Facundo et al. 2017, Riedle-Bauer et al. 2019a). Transmission from one year to the next can be sustained by infected overwintering adult specimens.

Numerous species of entomophagous insects are known as natural antagonists of *C. pyri*, both parasitoids, such as *Prionomitus mitratus* (Dalman) or *Trechnites psyllae* (Ruschka) and predators, such as species of the genera *Anthocoris* Fallen and *Orius* Wolff (Hemiptera, Anthocoridae). In particular, the predation by *Anthocoris nemoralis* (Fabr.) is highly significant; and the integrated control of this pear psyllid must be primarily based on techniques aimed at protecting and favouring (through augmentative releases) the action of this natural antagonist. This is why, in case of strong infestation requiring chemical control, low impact applications with selective insecticides having deterrent effects on oviposition (such as plant oils) (Erler and Tosun 2017) or with products having washing properties on the honeydew (such as salts of fatty acids) can be applied in mid-spring against the second generation nymphs, if intervention thresholds are exceeded.

In order to reduce the incidence of the Pear Decline phytoplasma, especially on young plants, control should be directed against overwintering adults of *C. pyri*, with the main

objective of reducing the vector population before the spring vegetation of plants, thus preventing the transmission on time.

***Cacopsylla pyricola* (Foerster, 1848)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Esch-sur-Sûre (1 ♂, 3 ♀♀, 15.V.2020, on *Pyrus communis*), Noerdange (1 ♂, 1 ♀, 28.V.2020, on *P. communis*); East Gutland: Bettendorf (3 ♂♂, 6 ♀♀, 27.V.2020, on *P. communis*), Olingen (6 ♂♂, 6 ♀♀, 27.V.2020, on *P. communis*); Moselle: Burmerange (1 ♀, 8.V.2020, on *P. communis*), Elvange (1 ♀, 1.VIII.2019, on *P. communis*), Elvange/Burmerange (2 ♀♀, 19.V. 2020, on *P. communis*).

Material studied in the MNHN collection: Minette: Schiffange, Brucherberg (2 ♂♂, 22.IV - 11.V.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Before the revisions of Burckhardt and Hodkinson (1986) and Cho et al. (2017b), respectively, of the west and east Palaearctic pear psyllids, *C. pyricola* was reported from the Asiatic Far East, Caucasus and all of Europe; and was also indicated as introduced in North and South America. At present, its occurrence is confirmed only in Europe (including the Caucasian Region) and North America (Burckhardt 1988a, Hodkinson 1988, Ossiannilsson 1992, Burckhardt and Lauterer 1993, Conci et al. 1993, Burckhardt 1994a, Conci et al. 1996, Agusti et al. 2003, Seljak 2006, Ripka 2008, Gertsson 2010, Inoue 2010, O'Connor and Malumphy 2011, Percy et al. 2012, Kanturski and Drohojowska 2013, Cooper and Horton 2015, Castillo Carrillo et al. 2016, Zendedel et al. 2016, Cho et al. 2017b, Holzinger et al. 2017, Drohojowska and Klasa 2019, den Bieman et al. 2019). As for all other European pear psyllids, the presence of *C. pyri* in East Asia was also questioned, based on recent DNA barcoding studies (Cho et al. 2020).

Biology. Oligophagous on plants of the genus *Pyrus* L. (Rosaceae), especially on *P. communis* L., but also frequently recorded on *P. calleryana* Decne., *P. pyrasater* (L.) and *P. ussuriensis* Maxim. (Burckhardt 1983, Burckhardt 1988a, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Burckhardt 1994a, Conci et al. 1996, Seljak 2006, Ripka 2008, Hodkinson 2009, Guédot et al. 2009, Inoue 2010, Percy et al. 2012, Cooper and Horton 2015, Spodek et al. 2017). It performs 3-5 generations per year, overwintering as adults on the host plants (but also on other occasional fruit or forestry plants in the surrounding area). In early spring, eggs are laid in groups on the buds; later in the season and throughout all summer, they are mostly laid on the underside of the leaves, along the middle vein or in the grooves of the leaf stalks.

Economic significance. In Europe, *C. pyricola* is the second most important and widespread species of pear psyllids, after *C. pyri*, to which we refer for information on harmfulness and control methods. Apart from the negative effects on infested plants consisting in the suction of sap and production of abundant honeydew, *C. pyricola* is also a

vector of the pathogenic microorganisms causing Pear Decline disease ('*Candidatus Phytoplasma pyri*') (Facundo et al. 2017, Cruz et al. 2018, Riedle-Bauer et al. 2019a); for such damage, it is particularly feared in North America.

***Cacopsylla pyrisuga* (Foerster, 1848)**

(Figs 2, 6, 10, 14)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Esch-sur-Sûre (1 ♂, 15.V.2020, on *Pyrus communis*), Noerdange (1 ♂, 28.V.2020, on *P. communis*); Moselle: Burmerange (1 ♀, 8.V.2020, on *P. communis*).

New record for Luxembourg.

Geographical distribution. Eurasian chorotype, widely distributed in Europe and diffused eastwards, through Turkey, the Middle East, the Caucasian Region and Iran (Ramírez Gómez 1956, Hodkinson and White 1979, Burckhardt 1983, Kwon 1983, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Lauterer 1993, Burckhardt and Önuçar 1993, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Lauterer 1999, Seljak 2006, Ripka 2008, Inoue 2010, Malenovský et al. 2011, Jerinić-Prodanović and Protić 2013, Kanturski and Drohojowska 2013, Serbina et al. 2015, Spodek et al. 2017, Drohojowska and Klasa 2019, den Bieman et al. 2019), though findings in the extreme eastern part of Asia have been recently referred to other taxa (Cho et al. 2017b, Cho et al. 2020).

Biology. Strictly oligophagous on *Pyrus* spp. (Rosaceae), with records in Europe especially on *P. amygdaliformis* Vill., *P. communis* L. and *P. elaeagnifolia* Pall.; in Asia, also reported from many other *Pyrus* species, amongst which *P. pashia* Buch.-Ham. ex D. Don, *Pyrus pyrifolia* (Burm. f.) Nakai, *P. salicifolia* Pall. and *P. ussurensis* Maxim. ex Rupr. (Hodkinson and White 1979, Burckhardt 1983, Kwon 1983, Burckhardt 1988a, Ossiannilsson 1992, Conci et al. 1993, Burckhardt 1994a, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Seljak 2006, Ripka 2008, Hodkinson 2009, Inoue 2010, Malenovský et al. 2011). It performs one generation per year and overwinters as an adult on shelter plants (conifers). Sometimes, it causes deformations to leaves, which appear wrinkled and folded (Houard 1908, Houard 1909, Houard 1913).

Economic significance. In the past, *C. pyrisuga* has been reported to be harmful to pears. Current knowledge suggests that it does not cause direct damage, being apparently better adapted to wild pears or old varieties of cultivated pear trees, especially in hilly areas and in non-intensive orchards. However, it has been recently confirmed as a vector of "*Ca. Phytoplasma pyri*", the causing agent of Pear Decline (Riedle-Bauer et al. 2022).

***Cacopsylla rhamnicola* (Scott, 1876)**

(Figs 2, 6, 10)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Brouch (3 ♀♀, 15.V.2020, on *Salix caprea*, occasional plant), Strassen (2 ♀♀, 27.IV.2020; 1 ♀, 8.VI. 2020, by general sweeping with net).

New record for Luxembourg.

Geographical distribution. Widely distributed from central Asia to nearly all Europe (Hodkinson and White 1979, Burckhardt 1988b, Ossiannilsson 1992, Burckhardt and Önuçar 1993, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, Ripka 2009, Gertsson 2010, Malenovský et al. 2011, O'Connor and Malumphy 2011, Ouvrard et al. 2015, den Bieman et al. 2019).

Biology. Strictly oligophagous on *Rhamnus* spp. (Rhamnaceae), especially *R. cathartica* L., but also reported on *R. alpinus* L., *R. fallax* Boiss., *R. imeretina* Booth et al. and *R. saxatilis* Jacq. (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988b, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, Ripka 2009, Hodkinson 2009, Malenovský et al. 2011); records in Luxembourg refer only to occasional plants, so far. *Cacopsylla rhamnicola* performs only one generation per year and overwinters as an adult on shelter plants (conifers).

***Cacopsylla ulmi* (Foerster, 1848)**

Findings in Luxembourg. Personal field collection by the authors: Moselle: Elvange/Burmerange, (2 ♂♂, 1 ♀, 1.VI.2020, by sweeping with net on a mix with *Acer platanoides* + *Populus cinerea*, occasional plants).

New record for Luxembourg.

Geographical distribution. *Cacopsylla ulmi* is a Eurasian chorotype, occurring in most parts of central and northern Europe, the Caucasian Region and central Asia (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Hansen 1996, Seljak 2006, Ripka 2008, Lauterer 2011, Malenovský et al. 2011, Kanturski and Drohojowska 2013, Serbina et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Strictly oligophagous on elm trees (*Ulmus* spp., Ulmaceae), *C. ulmi* has been especially found on *Ulmus effusa* Willd., *U. glabra* Huds., *U. laevis* Pall., *U. minor* Mill. and *U. pedunculata* Foug. (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Hansen 1996, Seljak 2006, Ripka 2008, Hodkinson 2009, Lauterer 2011, Malenovský et al. 2011). It performs one generation per year and spends the winter as an egg on the host plants. According to Lauterer (1991), part of its population migrates during the summer months to other plant species (mainly trees), which are used for shelter.

***Cacopsylla visci* (Curtis, 1835)**

(Figs 2, 6, 10)

Findings in Luxembourg. Personal field collection by the authors: Moselle: Elvange/Burmerange (5 ♀♀, 19.V.2020, on *Viscum album*).

New record for Luxembourg.

Geographical distribution. *Cacopsylla visci* is likely a Palaearctic chorotype. It is reported from nearly all of Europe (Hodkinson and White 1979, Lauterer 1979, Conci et al. 1993, Conci et al. 1996, Hansen and Hodkinson 2006, Seljak 2006, Ripka 2008, Varga et al. 2012, Ouvrard et al. 2015, Burckhardt et al. 2017, Struwe et al. 2009, Drohojowska and Klasa 2019, den Bieman et al. 2019), from North Africa (Morocco) (Inoue 2010), the Middle East and the Caucasian Region (Inoue 2010, Burckhardt et al. 2017), to Far East Asia (Japan and South Korea) (Inoue 2010, Cho et al. 2017a).

Biology. *Cacopsylla visci* lives only on Santalales of the genera *Loranthus* Jacq. (*L. europaeus* Jacq.) (Loranthaceae) and *Viscum* L. (*V. album* L., *V. laxum* Boiss. & Reut.) (Santalaceae) (Hodkinson and White 1979, Burckhardt 1983, Hansen and Hodkinson 2006, Seljak 2006, Hodkinson 2009, Inoue 2010). According to Bin (1970), it performs 2-3 generations per year, overwintering in the nymphal stage on its host plants, to which it causes showy deformations to the leaves, which are folded in a C-shape.

***Livilla ulicis* Curtis, 1836**

(Figs 3, 7, 11, 15)

Findings in Luxembourg. Material studied in the MNHN collection: East Gutland: Niederanven, Aarnesch (4 ♂♂, 3 ♀♀, 27.V - 9.VI.1999; 1 ♀, 9.VI - 25.VI.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Central European chorotype, never recorded till now in the northernmost part of the continent (Scandinavia), as well as in the “Benelux” Region and the Iberian Peninsula (Hodkinson and White 1979, Burckhardt 1983, Hodkinson and Hollis 1987, Conci et al. 1993, Conci et al. 1996, Seljak 2006, O’Connor and Malumphy 2011).

Biology. Oligophagous on plants of the genus *Genista* L. (Fabaceae); especially recorded from *G. tinctoria* L. and *G. germanica* L. (Hodkinson and White 1979, Burckhardt 1983, Hodkinson and Hollis 1987, Conci et al. 1993, Conci et al. 1996, Seljak 2006); records on other Fabaceae, such as *Cytisus scoparius* (L.) Link or *Ulex europaeus* L., are not confirmed. *Livilla ulicis* performs one generation per year and probably spends the winter in the egg stage on its host plants.

***Psylla alni* (Linnaeus, 1758)**

(Figs 3, 7, 11, 15)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Dondelange (5 ♂♂, 2 ♀♀, 28.V.2020, on *Alnus glutinosa*; 1 ♂, 28.V.2020, on *Crataegus monogyna*, occasional plant), Elvange/Schweich (1 ♂, 28.V.2020, on *Prunus spinosa*, occasional plant), Goesdorf/Bockholtz (7 ♂♂, 4 ♀♀, 20.V.2020, on *A. glutinosa*), Hobscheid (1 ♂, 28.V.2020, on *A. glutinosa*), Préitzerdaul (7 ♂♂, 5 ♀♀, 20.V.2020, on *A. glutinosa*); East Gutland: Bettendorf (2 ♂♂, 2 ♀♀, 27.V.2020, on *A. glutinosa*), Rodenburg (3 ♂♂, 4 ♀♀, 27.V.2020, on *A. glutinosa*); Minette: Belvaux (1 ♂, 1 ♀, 19.VII.2019, on *A. glutinosa*), Kayl (1 ♂, 20.V.2020, on *Malus* sp., occasional plant).

New record for Luxembourg.

Geographical distribution. Distributed in the Holarctic Region, with a wide presence in the Palaearctic (though not reported from North Africa) and in North America (Canada, Greenland, USA) (Hodkinson and White 1979, Kwon 1983, Yang 1984, Hodkinson 1986, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Önuçar 1993, Conci et al. 1996, Labina 2006, Seljak 2006, Ripka 2008, Gertsson 2010, Inoue 2010, Malenovský et al. 2011, O'Connor and Malumphy 2011, Percy et al. 2012, Kanturski and Drohojowska 2013, Drohojowska and Burckhardt 2014, Ouvrard et al. 2015, Serbina et al. 2015, Holzinger et al. 2017, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Strictly oligophagous on various species of alders (*Alnus* Mill., Betulaceae). In Europe, on *Alnus glutinosa* (L.) Gaertn. and *A. incana* (L.) Moench and sometimes on *A. viridis* (Chaix) DC.; in Asia, also on *A. hirsuta* Turcz. ex Rupr. and *A. japonica* (Thunb.) Steud.; in North America, on *A. rhombifolia* Nutt. (Hodkinson and White 1979, Yang 1984, Hodkinson 1986, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, Inoue 2010, Percy et al. 2012). On its host plants, this psyllid species performs only one generation per year, overwintering as an egg which is laid inside the shoots.

***Psylla betulae* (Linnaeus, 1758)**

(Figs 3, 7, 11, 15)

Findings in Luxembourg. Personal field collection by the authors: Moselle: Remerschen (3 ♂♂, 4 ♀♀, 28.V.2020, on *Betula pendula*).

New record for Luxembourg.

Geographical distribution. *Psylla betulae* is a Holarctic chorotype, widely distributed in central and northern Europe, but recorded also from many parts of Asia (Georgia, Mongolia, Far East Russia, Japan) and North America (Canada) (Hodkinson and White 1979, Hodkinson and MacLean 1980, Hodkinson 1988, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Labina 2006, Labina 2008, Gertsson 2010, Inoue 2010, O'Connor and Malumphy 2011, Serbina et al. 2015).

Biology. Strictly oligophagous on birches (*Betula* spp., Betulaceae), according to literature, it lives in Europe mainly on *B. pendula* Roth. or *B. pubescens* Ehrh. (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996); recorded in Asia also on many different *Betula* species, such as *B. middendorffii* Trautvetter & C.A. Meyer, *B. ermanii* Cham., *B. platyphyllea* Sukaczev and others (Hodkinson and MacLean 1980, Inoue 2010). It performs one generation per year and overwinters as eggs on its host plants.

***Psylla foersteri* Flor, 1861**

(Figs 3, 7, 11, 15)

Findings in Luxembourg. Personal field collection by the authors: Minette: Belvaux (2 ♂♂, 6 ♀♀, 19.VII.2019, on *Alnus glutinosa*); Moselle: Remerschen (3 ♂♂, 4 ♀♀, 28.V.2020, on *A. glutinosa*).

Material studied in the MNHN collection: Oesling: Sonlez, Pamer (1 ♂, 24.VIII - 22.IX. 2000, Malaise).

New record for Luxembourg.

Geographical distribution. The occurrence of *P. foersteri* is confirmed in the West Palaearctic Region, where it is present in most parts of Europe, North Africa (Algeria) and the Middle East (Caucasian Region, Lebanon, Turkey) (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988b, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1993, Burckhardt and Önuçar 1993, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Seljak 2006, Ripka 2008, Gertsson 2010, O'Connor and Malumphy 2011, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, den Bieman et al. 2019). Recorded also from New Zealand (Martoni et al. 2018), where it has been introduced.

Biology. Oligophagous on alder trees (*Alnus* Mill., Betulaceae); especially recorded on *Alnus glutinosa* (L.) Gaertn. and *A. incana* (L.) Moench (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Seljak 2006, Ripka 2008). Performs only one generation per year and overwinters in the egg stage on its host plants.

***Psylla hartigii* Flor, 1861**

(Figs 3, 7, 11, 15)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Strassen (4 ♂♂, 2 ♀♀, 27.IV.2020, on *Betula pendula*); Minette: Belvaux (2 ♂♂, 7.V.2020, on *B. pendula*; 1 ♀, 8.V.2020, on *B. pendula*).

New record for Luxembourg.

Geographical distribution. It probably has a wide distribution in the Holarctic Region; in addition to larger parts of Europe (especially its central and northern areas) (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, Gertsson 2010, O'Connor and Malumphy 2011, Kanturski and Drohojowska 2013, Serbina et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019), reports of *C. hartigii* concern the Far East (Japan and other nearby islands, such as Kuril and Sakhalin) and central Asia (Altai Province in Russia), the Caucasian Region and North America (USA and Canada) (Hodkinson 1988, Ossiannilsson 1992, Labina 2008, Inoue 2010).

Biology. Strictly oligophagous on birches (genus *Betula* L., Betulaceae), it has been found in Europe on *Betula alba* L., *B. pendula* Roth. and *B. pubescens* Ehrh., but records also originate from *B. platyphyllo* Sukaczev and *B. raddeana* Trautv. in Asia, as well as *B. populifolia* Marshall in North America (Hodkinson and White 1979, Hodkinson 1988,

Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Inoue 2010). It shows only one generation per year and overwinters in the egg (or perhaps in the nymphal) stage (Lauterer 1976).

***Spanioneura buxi* (Linnaeus, 1758)**

(Figs 3, 7, 11, 15)

Findings in Luxembourg. Galls produced by this psyllid on *Buxus sempervirens* are reported by Lambinon and Schneider (2004) (Oesling: Clervaux; Minette: Dudelange, Esch-sur-Alzette Mosel: Palmberg, Ahn; West Gutland: Bonnevoie, Bridel, Capellen; East Gutland: Diekirch, Ettelbruck, Stackels, Galgebierg, Betzdorf). Explicit mention on its finding in Luxembourg is provided by Eickermann et al. (2020).

Personal field collection by the authors: Oesling: Clervaux (1 ♀, 12.IX.2019, on *B. sempervirens*), Hosingen (1 ♀, 12.IX.2019, on *B. sempervirens*), Vianden (2 ♂♂, 4 ♀♀, 12.IX.2019, on *B. sempervirens*); West Gutland: Bettborn (3 ♀♀, 12.IX.2019, on *B. sempervirens*), Roodt-sur-Eisch (2 ♂♂, 5 ♀♀, 22.VIII.2019, on *B. sempervirens*), Strassen (4 ♂♂, 5 ♀♀, 22.VIII.2019, on *B. sempervirens*); East Gutland: Niederfeulen (1 ♂, 5 ♀♀, 12.IX.2019, on *B. sempervirens*), Reisdorf (1 ♂, 1 ♀, 12.IX.2019, on *B. sempervirens*); Moselle: Elvange/Burmerange (1 ♂, 1.VI.2020, by sweeping with net on a mix with *Acer platanoides* + *Populus cinerea*, occasional plants). Galls produced by *P. buxi* on *B. sempervirens* have been also detected by us in Weiler (Oesling), Bereldange, Colmar-Berg, Mersch and Steinsel (West Gutland), Luxembourg Ville (Minette).

Geographical distribution. Widely diffused in Europe, from where it seems to be native (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Seljak 2006, Ripka 2008, O'Connor and Malumphy 2011, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, den Bieman et al. 2019, Eickermann et al. 2020). Reported also from the Nearctic Region (Canada, USA, including Hawaii), where it has been probably introduced (Wheeler and Hoebeke 2005, Percy et al. 2012, Castillo Carrillo et al. 2016).

Biology. Strictly oligophagous on *Buxus* spp. (Buxaceae), this psyllid species has been reported on *Buxus balearica* Lam., *B. macrophylla* (Britton) Fawc. & Rendle and *B. sempervirens* L. (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt 2005, Percy et al. 2012). It performs one generation per year, always remaining on the host plants, on which it overwinters in the egg stage (or even as first instar nymph, according to environmental conditions) (Eickermann et al. 2020). *P. buxi* causes singular galls to box plants, creating "spoon-like" folds of the apical leaves, which are often close together for the shortening of the internodes, generating the typical "artichoke" appearance of the most complex galls.

***Spanioneura fonscolombii* Foerster, 1848**

(Figs 3, 7, 11, 15)

Findings in Luxembourg. First reported from Luxembourg by O'Connor and Malumphy (2011).

Personal field collection by the authors: Oesling: Clervaux (6 ♂♂, 8 ♀♀, 12.IX.2019, on *Buxus sempervirens*), Hosingen (1 ♀, 12.IX.2019, on *B. sempervirens*), Vianden (1 ♂, 2 ♀♀, 12.IX.2019, on *B. sempervirens*), Weller (6 ♂♂, 1 ♀, 12.IX.2019, on *B. sempervirens*); West Gutland: Roodt-sur-Eisch (20 ♂♂, 20 ♀♀, 22.VIII.2019, on *B. sempervirens*); East Gutland: Niederfeulen (5 ♂♂, 4 ♀♀, 12.IX.2019, on *B. sempervirens*), Reisdorf (5 ♂♂, 12 ♀♀, 12.IX.2019, on *B. sempervirens*).

Geographical distribution. As far as presently inferred from literature, *S. fonscolombii* has a predominantly western European distribution, with records originating from Belgium, France, Great Britain, Ireland, Italy, Luxembourg, Slovenia, Spain, Sweden and Switzerland (Hodkinson and White 1979, Conci et al. 1993, Conci et al. 1996, Burckhardt and Mühlethaler 2003, Seljak 2006, Gertsson 2010, O'Connor and Malumphy 2011, Ouvrard et al. 2015, den Bieman et al. 2019); it is also reported from the Caucasian Region (Azerbaijan; Conci et al. 1996) and from USA (Hodkinson 1988), where it has been probably introduced.

Biology. Strictly oligophagous on *Buxus* spp., especially found on *B. sempervirens* L. (Buxaceae) (Hodkinson and White 1979, Burckhardt 1983, Hodkinson 1988, Conci et al. 1993, Conci et al. 1996, Burckhardt and Mühlethaler 2003, Seljak 2006, Hodkinson 2009). It performs one generation per year and overwinters in the adult stage on its host plants.

Triozaeidae

***Bactericera albiventris* (Foerster, 1848)**

(Figs 4, 8, 12)

Findings in Luxembourg. Personal field collection by the authors: Minette: Belvaux (2 ♀♀, 19.VII.2019, on *Salix alba*; 1 ♀, 8.VIII.2019, on *S. alba*; 1 ♀♀, 8.VIII.2019, on *Crataegus monogyna*, occasional plant).

New record for Luxembourg.

Geographical distribution. *Bactericera albiventris* is an Eurasian chorotype, widely distributed in nearly all of Europe and eastwards to the Middle East (Iran, Israel, Lebanon, Turkey), Central Asia (Afghanistan, Caucasian Region, Mongolia, Tadzhikistan, Turkmenistan, Uzbekistan) to East Asia (Far East Russia) (Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1988b, Halperin et al. 1988, Ossiannilsson 1992, Burckhardt and Önuçar 1993, Conci et al. 1996, Burckhardt and Lauterer 1997a, Zeidan-Gèze and Burckhardt 1998, Burckhardt 2005, Seljak 2006, Ripka 2008, Hodkinson 2009, Gertsson 2010, O'Connor and Malumphy 2011, Malenovský et al. 2012, Kanturski and Drohojowska 2013, Zendedel et al. 2016, Spodek et al. 2017, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Strictly oligophagous on *Salix* spp. (Salicaceae), with records reported in literature on many species of willows, as *S. acmophylla* Boiss., *S. alba* L., *S. amygdalina* L., *S. aurita* L., *S. babylonica* L., *S. elaeagnos* Scop., *S. fragilis* L., *S. pentandra* L., *S. purpurea* L., *S. triandra* L. and *S. viminalis* L. (Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1988b, Ossiannilsson 1992, Conci et al. 1996, Burckhardt and Lauterer 1997a, Zeidan-Gèze and Burckhardt 1998, Burckhardt 2005, Seljak 2006, Ripka 2008, Hodkinson 2009, Malenovský et al. 2012, Spodek et al. 2017). It performs one or two generations per year and overwinters as an adult on shelter plants (conifers).

***Bactericera curvatinervis* (Foerster, 1848)**

(Figs 4, 8, 12)

Findings in Luxembourg. Personal field collection by the authors: West Gutland: Brouch (4 ♀♀, 15.V.2020, on *Salix viminalis*).

New record for Luxembourg.

Geographical distribution. Known from most of central and northern Europe, it is reported also from Lebanon, the Caucasus and far east Asia (Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt and Lauterer 1997a, Zeidan-Gèze and Burckhardt 1998, Seljak 2006, Ripka 2008, Labina and Evdogarova 2009, Gertsson 2010, O'Connor and Malumphy 2011, Ouvrard et al. 2015, Serbina et al. 2015, den Bieman et al. 2019).

Biology. Strictly oligophagous on willows (*Salix* spp., Salicaceae), reported in literature from *S. alba* L., *S. appendiculata* Vill., *S. aurita* L., *S. caprea* L., *S. cinerea* L., *S. elaeagnos* Scop., *S. glabra* Scop., *S. purpurea* L., *S. repens* L. and *S. viminalis* L. (Ossiannilsson 1992, Conci et al. 1993, Conci et al. 1996, Burckhardt and Lauterer 1997a, Ripka 2008, Labina and Evdogarova 2009). *Bactericera curvatinervis* performs only one generation per year and overwinters as an adult on shelter plants (conifers).

***Bactericera substriola* Ossiannilsson, 1992**

(Figs 4, 8, 12, 16)

Findings in Luxembourg. Material studied in the MNHN collection: East Gutland: Wilferdange, Conzefenn (1 ♂, 22.IX - 19.X.2000, Malaise); Minette: Kockelscheuer, Conter Jans Boesch (1 ♀, 22.VII - 5.VIII.1999; 2 ♀♀, 2.IX - 27.IX.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Known till now only from a few countries in central and northern Europe (Austria, Belarus, Czech Republic, Germany, Great Britain, Sweden, Switzerland, The Netherlands) (Ossiannilsson 1992, Burckhardt 1994b, Burckhardt and Lauterer 1997a, Malenovský and Lauterer 2012, Serbina et al. 2015, den Bieman et al. 2019), but the species is probably more widespread on the European continent.

Biology. Reported till now only from *Salix elaeagnos* Scop. and *S. lapponum* L. (Salicaceae) (Ossiannilsson 1992, Burckhardt and Lauterer 1997a), but probably living also on other willows (*Salix* spp.) (Malenovský and Lauterer 2012, den Bieman et al. 2019). It spends the winter as an adult on shelter plants (conifers), on which it has been collected during many of its findings; probably performs only one generation per year.

***Eryngiofaga lautereri* Loginova, 1977**

(Figs 4, 8, 16)

Findings in Luxembourg. Material studied in the MNHN collection: East Gutland: Niederanven, Aarnescht (2 ♂♂, 27.V - 9.VI.1999, Malaise).

New record for Luxembourg.

Geographical distribution. Known till now only from a few countries in central-eastern Europe, *E. lautereri* has been described on material from the Czech Republic (Moravia) and Germany (Thuringia) (Loginova 1977), although the German material has been erroneously indicated in the original description as originating from Austria (according to Lauterer 1979). Later, it was reported from other localities in the Czech Republic and Germany, as well as from Slovakia (Lauterer 1979, Malenovský and Lauterer 2012). It seems to be a rare species.

Biology. Trophically linked to plants of the genus *Bupleurum* L. (Apiaceae) (Hodkinson, 2009), it is reported exclusively to live on *B. falcatum* L. (Loginova 1977, Lauterer 1979, Malenovský et al. 2011, Malenovský and Lauterer 2012). Life cycle of this psyllid has been studied by Lauterer (1979), who reports two generations per year, with adults emerging in May-June and in August-September, respectively (only one generation may be shown by part of the population, which spends summer as diapausing nymphs of the first offspring); winter is spent by the nymphs (of both generations) on the host plant.

***Lauritrioza alacris* (Flor, 1861)**

(Figs 4, 8, 12)

Findings in Luxembourg. No explicit mention exists in literature on findings of this insect in Luxembourg, though its galls produced on *Laurus nobilis* are reported by Lambinon and Schneider (2004) and by Schneider (2016) (West Gutland: Bonnevoie; Moselle: Remich), who did not report, however, any finding of this psyllid in or nearby the galls. Therefore, the present record is the first direct finding of *L. alacris* in Luxembourg.

Personal collection by the authors in: Oesling: West Gutland: Bertrange (1 ♀, 13 nymphs, 7.VI.2020, on *L. nobilis*).

Geographical distribution. Most probably native from the Mediterranean Region on wild laurel and widely distributed in this area (Halperin et al. 1982, Burckhardt 1988a, Burckhardt 1989, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Spodek et al. 2017), it has spread (and has been introduced) on cultivated *L. nobilis* in nurseries, gardens and

parks throughout nearly all of Europe, including central and northern parts of the continent, up to Scandinavia (Hodkinson and White 1979, Hodkinson and White 1981, Burckhardt 1988c, Ossiannilsson 1992, Conci et al. 1996, O'Connor et al. 1997, Hollis and Martin 1997, Burckhardt and Mühlethaler 2003, Burckhardt 2005, Seljak 2006, Cocquempot 2008, O'Connor and Malumphy 2011, den Bieman et al. 2019), as well as eastwards to Anatolia (Turkey), the Caucasian Region, the Crimean Peninsula (Burckhardt 1989, Ossiannilsson 1992, Zeidan-Gèze and Burckhardt 1998, Burckhardt and Mühlethaler 2003) and westwards to North (USA) and South America (Argentina, Brazil, Chile) (Hodkinson and White 1981, Burckhardt 1988c, Hodkinson 1988, Mifsud et al. 2010, Burckhardt and Queiroz 2012, Percy et al. 2012).

Biology. According to the large body of literature on this species (Hodkinson and White 1979, Hodkinson and White 1981, Burckhardt 1983, Burckhardt 1988a, Burckhardt 1988b, Hodkinson 1988, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1996, Hollis and Martin 1997, O'Connor et al. 1997, Zeidan-Gèze and Burckhardt 1998, Burckhardt and Mühlethaler 2003, Burckhardt 2005, Seljak 2006, Cocquempot 2008, Hodkinson 2009, Marchini et al. 2012, Percy et al. 2012, Burckhardt and Queiroz 2012, Spodek et al. 2017), *L. alacris* is oligophagous on plants of the genus *Laurus* L. (Lauraceae), with *L. nobilis* L. being its most frequent host plant, but likely also living on other congeneric plant species. During spring and summer, adults are extremely active, flying intensively and spreading around from their host plants; this is why they can be found (and have been recorded in the past literature) on a great number of occasional plants. *Lauritrioza alacris* performs 1-2 generations per year, overwintering as an adult on the host plants. It causes leaf galls, by rolling their margins down to the lower surface (Houard 1908, Houard 1909, Houard 1913, Ross and Hedicke 1927, Buhr 1964, Buhr 1965, Sampò 1977).

Economic significance. In case of a strong infestation, leaf deformations and galls caused by this psyllid, which dry up and become black as soon as they are abandoned by the nymphs, may cause serious aesthetic damage to cultivated laurel in gardens and parks and direct control of this insect may be occasionally necessary, especially in nurseries and on young plants.

***Trichochermes walkeri* (Foerster, 1848)**

Findings in Luxembourg. *Trichochermes walkeri* was reported from Luxembourg by Baugnée (2001). In addition, the galls it causes on *Rhamnus cathartica* L. were reported for the country by Lambinon and Schneider (2004) (Moselle: Ahn, Wormeldange; Minette: Dudelange, Reckange-sur-Mess, Rumelange; West Gutland: Hamm, Steinfort, East Gutland: Rosport).

Geographical distribution. European chorotype, with a distribution almost limited to the central and northern part of the continent (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1996, Seljak 2006, Ripka 2008, Ripka 2009, Malenovský et al. 2011, Ouvrard et al. 2015, Serbina et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Strictly oligophagous on *Rhamnus* spp. (Rhamnaceae), with records especially on *R. cathartica* L. (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1996, Seljak 2006, Ripka 2008, Ripka 2009, Malenovský et al. 2011), but also frequently recorded on *R. saxatilis* Jacq. (Cobau 1912, Ripka 2008). It performs one generation per year and overwinters as an egg on the host plants.

***Trioza abdominalis* Flor, 1861**

(Figs 4, 8, 12, 16)

Findings in Luxembourg. Material studied in the MNHN collection: East Gutland: Wilferdange, Conzefenn (2 ♂♂, 4 ♀♀, 22.IX - 19.X.2000, Malaise).

New record for Luxembourg.

Geographical distribution. Widely distributed in Europe, especially in the north and central part of the continent (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1996, Seljak 2006, Gertsson 2010, Malenovský and Lauterer 2012, Malenovský et al. 2014, Ouvrard et al. 2015, Serbina et al. 2015, Drohojowska and Klasa 2019). Reports from central and east Asia, such as from the Korean Peninsula or Mongolia (Kwon 1983) need to be confirmed.

Biology. Based on data from the literature, *T. abdominalis* is widely oligophagous on plants belonging to the genera *Achillea* L., *Anthemis* L. and *Chrysanthemum* L. (Asteraceae) (Hodkinson and White 1979, Burckhardt 1983, Ossiannilsson 1992, Conci et al. 1996, Seljak 2006, Hodkinson 2009, Gertsson 2010, Malenovský and Lauterer 2012, Malenovský et al. 2014), with *Achillea millefolium* L. being the most common host plant. *Trioza abdominalis* performs one generation per year and spends winter as an adult on shelter plants (conifers).

***Trioza centranthi* (Vallot, 1829)**

Findings in Luxembourg. No explicit mention exists in literature till now on findings of this insect in Luxembourg, though its galls produced on *Centranthus ruber* are reported by Lambinon et al. (2012) (West Gutland: Bonnevoie, Buschdorf, Kirchberg; Minette: Dudelange, Tétange; Moselle: Mertert) and by Schneider (2016) (Oesling: Stolzembourg; West Gutland: Howald, Luxembourg; East Gutland: Schoos), who did not report, however, any finding of this psyllid in or nearby the galls. Therefore, the present record is the first direct finding of *T. centranthi* in Luxembourg.

Personal collection by the authors in: West Gutland, Strassen (1 ♀, 27.IV.2020, from general sweeping with net); Moselle: Remerschen (1 leaf gall with 3 nymphs inside, 28.V. 2020, on *Centranthus ruber*).

Geographical distribution. Euro-Mediterranean chorotype, distributed in central and eastern Europe (northwards up to Denmark and Great Britain; Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1996, Seljak 2006, Ripka 2008, Cocquempot 2008, Ouvrard et al. 2015, Spooner 2016, Badmin 2017,

Drohojowska and Klasa 2019, den Bieman et al. 2019) and the Mediterranean Basin (Algeria, Morocco, Israel; Halperin et al. 1982, Burckhardt 1989, Spodek et al. 2017); distributed eastwards to Turkey and the Caucasian Region (Burckhardt 1989, Ossiannilsson 1992, Burckhardt and Önuçar 1993).

Biology. Widely oligophagous on several species of the genera *Centranthus* DC. [*C. angustifolius* (Mill.) DC., *C. calcitrapae* (L.) Dufr., *C. ruber* (L.) DC.], *Fedia* Gaertn. (*F. cornucopiae* Gaertn.) and *Valerianella* Mill. [*V. carinata* Loisel., *V. coronata* (L.) DC., *V. dentata* (L.) Pollich, *V. locusta* (L.) Laterr. (= *V. olitoria* (L.) Pollich), *V. rimosa* Bast.] (Caprifoliaceae) (Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1996, Seljak 2006, Cocquempot 2008, Ripka 2008, Hodkinson 2009, Spooner 2016, Badmin 2017). It usually performs one generation per year, overwintering as an adult on shelter plants (conifers), though in warmer areas, it likely shows an almost continuous life-cycle (with an undefined number of yearly generations), spending the winter in all developmental stages.

On its host plants, *T. centranthi* causes showy leaf deformations, by rolling up the margins and forming irregular and turgid galls, which are initially pale green and later turn red; flowers and inflorescences may be also deformed by this psyllid, through hypertrophies or atrophies of the stamens or distortion of apical flowers (which are reduced to subglobular processes) (Houard 1908, Houard 1909, Houard 1913, Sampò 1975, Conci et al. 1996).

***Trioza cirsii* Löw, 1881**

(Figs 4, 8, 12, 16)

Findings in Luxembourg. Material studied in the MNHN collection: East Gutland: Koedange, Poenn (8 ♂♂, 13 ♀♀, 8.VII - 22.VII.1999; 5 ♂♂, 12 ♀♀, 22.VII - 5.VIII.1999; 1 ♀, 19.VIII - 2.IX.1999; 8 ♂♂, 6 ♀♀, 2.IX - 27.IX.1999, Malaise).

New record for Luxembourg.

Geographical distribution. With the exception of the Balkan Peninsula (Bulgaria, Greece, Romania), findings of *T. cirsii* concern exclusively central-northern Europe (Burckhardt 1988a, Ossiannilsson 1992, Seljak 2006, Gertsson 2010, Malenovský et al. 2011, Ouvrard et al. 2015, Holzinger et al. 2017, Drohojowska and Klasa 2019).

Biology. *Trioza cirsii* is oligophagous on Asteraceae of the genus *Cirsium* Mill., with records confirmed from *C. arvense* (L.) Scop., *C. erisithales* Scop., *C. heterophyllum* (L.) Hill, *C. oleraceum* Scop., *C. palustre* (L.) Scop. (Burckhardt 1988a, Ossiannilsson 1992, Seljak 2006, Hodkinson 2009, Malenovský et al. 2011). The number of yearly generations is unknown. It overwinters as adult on shelter plants (conifers).

***Trioza flavipennis* Foerster, 1848**

Findings in Luxembourg. This species has not been found by the authors in Luxembourg so far. Nevertheless, findings of its leaf deformations produced on *Aegopodium podagraria* are reported by Lambinon and Schneider (2004) and by Burton et al. (2019) (West

Gutland: Hollenfels, East Gutland: Manternach), who also mentioned the occurrence of eggs and nymphs of this psyllid species in the above galls, thus explicitly recording its occurrence in Luxembourg.

Geographical distribution. *Trioza flavipennis* is a typical European chorotype, being distributed mainly in the central and northern areas of the continent (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1996, Hansen and Greve 1999, Seljak 2006, Ripka 2008, Malumphy et al. 2009, Gertsson 2010, Ripka 2012, Serbina et al. 2015, Drohojowska and Klasa 2019, den Bieman et al. 2019).

Biology. Monophagous on *Aegopodium podagraria* L. (Apiaceae) (Hodkinson and White 1979, Ossiannilsson 1992, Conci et al. 1996, Hansen and Greve 1999, Seljak 2006, Ripka 2008, Malumphy et al. 2009). It performs likely only one generation per year and overwinters as an adult on shelter plants (conifers). On leaves of its host plant, this psyllid produces small pit-galls, protruding to the upper face (Houard 1908, Houard 1909, Cobau 1912, Houard 1913, Ross and Hedicke 1927, Buhr 1964, Buhr 1965, Lauterer 1991).

***Trioza galii* Foerster, 1848**

(Figs 4, 8, 12, 16)

Findings in Luxembourg. Personal field collection by the authors: Moselle: Bech/Kleinmacher (1 ♀, 30.VII.2019, on *Fraxinus excelsior*, occasional plant; 3 ♀♀, 8.V.2020, from general sweeping with net).

Material studied in the MNHN collection: East Gutland: Koedange, Poenn (1 ♂, 8.VII - 22.VII.1999, Malaise); Moselle: Canach, Wéngertsbierg (1 ♂, 25.VI - 8.VII.1999, Malaise).

New record for Luxembourg.

Geographical distribution. A large number of literature records is available for this species for nearly the entire Palaearctic Region; nevertheless, due to a recent assessment of the *T. galii* species group (Burckhardt and Lauterer 2006), many reports from the East Palaearctic Region may refer to other taxa. Therefore, based on the presently reliable information, a Western Palaearctic distribution extending to central Asia should be referred to for this psyllid.

Biology. Widely oligophagous on numerous Rubiaceae of the genera *Asperula* L., *Cruciata* Mill., *Galium* L., *Rubia* L. and *Sherardia* L. (Hodkinson and White 1979, Hodkinson and Hollis 1981, Hodkinson 1983, Burckhardt 1988a, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1996, Burckhardt 2005, Burckhardt and Lauterer 2006, Seljak 2006, Ripka 2008, Hodkinson 2009, Malenovský et al. 2011, Newbould 2012, Spodek et al. 2017, den Bieman et al. 2019). Literature offers numerous different descriptions of its life-cycle, probably deriving from the taxonomic confusion that has only recently been unravelled; based on the reliable information available today, *T. galii* very likely overwinters as an adult, but it is not possible to indicate details on its biology, including the number of yearly generations (Burckhardt and Lauterer 2006). Similarly, the gall-forming activity of this

species being confirmed, the considerable quantity of cecidia described and attributed to *T. galii* (deformation and redness of terminal leaves, shortening of internodes, rolling up of the leaf margins, hypogean cecidia on rhizomes etc.) (Houard 1908, Houard 1909, Houard 1913, Ross and Hedicke 1927, Docters van Leeuwen 1937, Buhr 1964, Buhr 1965) is most likely to be referred to different species of the group.

***Trioza remota* Foerster, 1848**

(Figs 4, 8, 12, 16)

Findings in Luxembourg. Galls produced by this psyllid on *Quercus robur* L. are reported by Lambinon and Schneider (2004) and by Schneider (2016) (Oesling: Troisvierges; West Gutland: Bonnevoie, Biergerkräiz, Howald, Kirchberg, Luxembourg, Steinfort; East Gutland: Oberanven; Minette: between Bergem and Schiffange, Dudelange, Esch-sur-Alzette, Frisange, Kayl, Schiffange; Moselle: between Gare d'Ellange and Altwies, Mondorf-les-Bains). The latter author also refers to the occurrence of an insect nymph within each pit gall, thus explicitly mentioning the finding of this insect in Luxembourg.

Personal collection by the authors in: Oesling: Arsdorf (21 nymphs, 22.VIII.2019, on *Quercus robur*); West Gutland: Strassen (2 pit-galls with nymphs, 16.VIII.2020, on *Quercus* sp.).

Material studied in the MNHN collection: Oesling: Basbellain, Klengelbaach (3 ♂♂, 7 ♀♀, 22.IX - 19.X.2000, Malaise), Eiselborn, Bréichen (2 ♂♂, 2 ♀♀, 19.X - 7.XI.2000, Malaise), Goebelsmühle (5 ♂♂, 11 ♀♀, 22.IX - 19.X.2000, Malaise), Hoscheid, Molberlay (2 ♂♂, 1 ♀, 22.IX - 19.X.2000, Malaise), Lellingen, Op Baerel (2 ♂♂, 2 ♀♀, 22.IX - 19.X.2000, Malaise); West Gutland: Capellen, Werwelslach (1 ♀, 13.IV - 22.IV.1999, Malaise); East Gutland: Godbrange, Schléidelbierg (1 ♂, 1 ♀, 6.IV - 22.IV.1999; 1 ♀, 22.IV - 11.V.1999, Malaise), Wilferdange, Conzefenn (2 ♀♀, 22.IX - 19.X.2000, Malaise); Minette: Kockelscheuer, Conter Jans Boesch (1 ♀, 13.IV - 22.IV.1999; 1 ♂, 2.IX - 27.IX.1999, Malaise), Schiffange, Kayl, Brucherbierg (1 ♀, 22.IV - 11.V.1999, Malaise).

Geographical distribution. *Trioza remota* shows a West Palaearctic distribution, with records from most parts of Europe and the Mediterranean Region (Hodkinson and White 1979, Burckhardt 1989, Ossiannilsson 1992, Conci et al. 1996, Seljak 2006, Ripka 2008, Malumphy et al. 2009, Gertsson 2010, Malenovský et al. 2011, O'Connor and Malumphy 2011, Serbina et al. 2015, Spodek et al. 2017, Drohojowska and Klasa 2019, den Bieman et al. 2019), eastwards to Anatolia, the Caucasian Region and Iran (Burckhardt 1989, Önuçar and Ulu 1991, Ossiannilsson 1992, Burckhardt and Lauterer 1993, Burckhardt and Önuçar 1993, Malumphy et al. 2009). Records from Japan (Burckhardt 1989, Önuçar and Ulu 1991, Ossiannilsson 1992, Burckhardt and Lauterer 1993, Burckhardt and Önuçar 1993, Malumphy et al. 2009) would require further verification.

Biology. Strictly oligophagous on many deciduous species of the genus *Quercus* L. (Fagaceae), with most frequent records on *Q. petraea* (Matt.) Liebl. and *Q. robur* L. (Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1989, Önuçar and Ulu 1991, Ossiannilsson 1992, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Seljak 2006,

Ripka 2008, Hodkinson 2009, Malumphy et al. 2009, Malenovský et al. 2011, Spodek et al. 2017). According to Lauterer (1991), it performs only one generation per year, overwintering as an adult on shelter plants (conifers) and with a very slow nymphal development during summer (from May to late August), which is spent as a 2nd instar nymph in a sort of “parapause”; aestivating nymphs soon resume their development in September and produce adults already in October (the latter overwinter).

This psyllid produces little pit galls on the leaves of its host plants, protruding to the upper face; the nymph causing each gall settles on the corresponding concavity occurring on the lower face (Houard 1908, Houard 1909, Houard 1913, Ross and Hedicke 1927, Buhr 1964, Buhr 1965).

***Trioza rhamni* (Schrank, 1801)**

Findings in Luxembourg. No explicit mention exists in literature on findings of this insect in Luxembourg, though its galls produced on *Rhamnus cathartica* L. are reported by Lambinon and Schneider (2004) (Moselle: Ahn), who did not report, however, any finding of this psyllid in or nearby the galls.

This species has not been found by us during our research in 2019-2020 nor in the investigated Malaise and Möricker trap samples; therefore, based on finding of its galls, it likely occurs in Luxembourg, but its presence needs to be confirmed by direct observations.

Geographical distribution. *Trioza rhamni* is a European chorotype, distributed and recorded over nearly the whole continent eastwards to Turkey and the Caucasian Region (Hodkinson and White 1979, Burckhardt 1988b, Ossiannilsson 1992, Burckhardt and Önuçar 1993, Conci et al. 1996, Hansen and Greve 1999, Seljak 2006, Ripka 2008, Ripka 2009, Gertsson 2010, Malenovský et al. 2011, Kanturski and Drohojowska 2013, Ouvrard et al. 2015).

Biology. Strictly oligophagous on *Rhamnus* spp., with *R. cathartica* L. being its most frequent host plant, but also reported on *Rhamnus pallasii* Fisch. & C.A. Mey. (Hodkinson and White 1979, Burckhardt 1983, Burckhardt 1988b, Ossiannilsson 1992, Conci et al. 1996, Hansen and Greve 1999, Seljak 2006, Ripka 2008, Ripka 2009, Malenovský et al. 2011, Kanturski and Drohojowska 2013). It probably performs two generations per year (Lauterer 1991) and overwinters as an adult on shelter plants (conifers). As a result of egg deposition, *T. rhamni* causes small pit-galls on its host plant leaves (Houard 1908, Houard 1909, Cobau 1912, Houard 1913, Ross and Hedicke 1927, Buhr 1964, Buhr 1965).

***Trioza urticae* (Linnaeus, 1758)**

(Figs 4, 8, 12, 16)

Findings in Luxembourg. Galls produced by this species on *Urtica dioica* are reported by Lambinon et al. (2012) (Oesling: between Perlé and Holtz, Troisvierges; Minette: Dudelange; Moselle: Remich; West Gutland: Bonnevoie, Gantenbeinsmühle, Schleifmühle,

Steinfort; East Gutland: Bleesbréck, Gare de Sandweiler-Contern), who did not report, however, any finding of this psyllid in or nearby the galls. Therefore, the present record is the first direct finding of *T. urticae* in Luxembourg.

Personal collection by the authors in: Oesling: West Gutland: Arsdorf (1♀, 20.V.2020, from general sweeping with net; 1♂, 1♀, 8.VI.2020, on *Urtica dioica* L.), Brouch (1♀, 6.IX.2019, from general sweeping with net); Minette: Kayl/Tetange (13♂, 16♀, 23.VII.2019, on *U. dioica*).

Material studied in the MNHN collection: Oesling: Eselborn, Bréichen (1♂, 19.X - 7.XI.2000, Malaise); West Gutland: Capellen, Engelsratt (4♂, 3♀, 9.VI - 25.VI.1999; 15♂, 10♀, 25.VI - 8.VII.1999; 6♂, 8♀, 8.VII - 22.VII.1999; 5♂, 4♀, 22.VII - 5.VIII.1999, Malaise); East Gutland: Niederanven, Aarnesch (1♀, 9.VI - 25.VI.1999, Malaise), Waldhausenerdickt, Sauerwisen (5♂, 15♀, 22.IX - 19.X.2000, Malaise), Wilferdange, Conzefenn (1♀, 3.VIII - 24.VIII.2000, Malaise); Minette: Kockelscheuer, Conter Jans Boesch (1♂, 19.VIII - 2.IX.1999, Malaise); Moselle: Canach, Wéngertsbierg (1♂, 9.VI - 25.VI.1999, Malaise).

Geographical distribution. Widely distributed in the Palaearctic Region, from the Azores to Japan, being one of the psyllid species with the highest number of records reported in literature (amongst many others: Mathur 1975, Hodkinson and White 1979, Burckhardt 1988a, Burckhardt 1989, Halperin et al. 1988, Ossiannilsson 1992, Burckhardt and Lauterer 1993, Burckhardt and Önuçar 1993, Conci et al. 1996, Zeidan-Gèze and Burckhardt 1998, Aguiar and Martin 2001, Burckhardt 2005, Seljak 2006, Tishetshkin 2007, Labina 2008, Ripka 2008, Ripka and Kiss 2008, Malumphy et al. 2009, Gertsson 2010, Lauterer 2011, O'Connor and Malumphy 2011, Malenovský et al. 2011, Malenovský et al. 2012, Kanturski and Drohojowska 2013, Ouvrard et al. 2015, Serbina et al. 2015, Zendedel et al. 2016, Holzinger et al. 2017, Spodek et al. 2017, Haapalainen et al. 2018, Drohojowska and Klasa 2019, den Bieman et al. 2019

Biology. Strictly oligophagous on *Urtica* spp. (Urticaceae), with many records especially from *U. dioica* L. and *U. urens* L., but also from other plant species having a more restricted geographical distribution within the wide distribution area of this psyllid. *Trioza urticae* has a relatively rapid life cycle (depending on environmental conditions) and performs various generations per year (even more than four) on its host plants; it overwinters as an adult on various shelter plants, especially conifers.

This psyllid species may sometimes cause deformations to plants, by wrinkling or bubbling their leaves (Houard 1908, Houard 1909, Houard 1913, Ross and Hedicke 1927, Zangheri 1954, Buhr 1964, Buhr 1965, Sampò 1975).

Final remarks

The research here presented, realised through field collections and the study of historical collection material stored at the MNHN, allowed us to significantly increase the number of species of Psylloidea known for Luxembourg. Based on data available in literature so far

and considering also the species already known in the country only by the records of their galls or deformations, the Luxembourg psyllid fauna has more than tripled as a result of the present work. The increase is even greater (almost 7-fold) if the number of species previously known for the territory is strictly limited to only those reported through the direct finding of specimens. As to methodological aspects, it is interesting to note that the contribution of passive collection - in particular Malaise trapping - added seven new species to our assessment.

Despite the new records here presented, our knowledge of the psyllid fauna of Luxembourg is still incomplete. Further sampling is required on potential host plants on which psyllids have not been collected so far. Thus, for example, research on plants of the genus *Sorbus* L. must be intensified, as well as on numerous herbaceous plants that host in Europe psyllid species of the genera *Craspedolepta* Enderlein, *Bactericera* Puton and *Trioza* Foerster; on willows (*Salix* spp.), further research could also allow us to find additional psyllid species. Considering the floristic richness occurring in Luxembourg and also the psyllid fauna of neighbouring or close countries which are better explored, despite its small size, it is likely that further research could lead to a significant increase of up to about 20% of the psyllid species found so far in Luxembourg.

Finally, on the applied level, all species living on agricultural crops deserve appropriate future attention, especially those belonging to species complexes recognised as vectors of phytopathogenic organisms and whose real distribution and harmfulness in Luxembourg should be suitably monitored. This can be realised by involving local plant protection services, thus to further investigate possible impacts of changing environments on biology, vectoring activity, pest importance and spread of individual species.

Funding program

The work was carried out within the project “Risk assessment for psyllids and whiteflies under current and future climate conditions in Luxembourg”, financed by the Fonds National de la Recherche de Luxembourg (FNR) in the framework of the INTER MOBILITY programme (INTER/Mobility/18/12958064/RAPID-IN/).

Conflicts of interest

The authors have no conflicts of interest.

References

- Aguiar AM, Martin JH (2001) Psyllids (Homoptera: Psylloidea) from Madeira Island: an updated checklist including new records. Suplemento ao Boletim da Sociedade Portuguesa de Entomologia 6: 335-342.

- Agusti N, Unruh TR, Welter SC (2003) Detecting *Cacopsylla pyricola* (Hemiptera: Psyllidae) in predator guts using COI mitochondrial markers. Bulletin of Entomological Research 93: 179-185. <https://doi.org/10.1079/BER2003236>
- Allahverdi T, Rahimian H, Babaeizad V (2014) Association of *Candidatus Phytoplasma prunorum* associated with plum yellow leaf stunt in Iran. Iranian Journal of Plant Pathology 50 (2): 185.
- Andrianjaka-Camps ZN, Schaefer S, Kuske S, Bünter M, Sauvion N, Christen D (2018) Influence of the distribution and infection rates of psyllids on the vectoring ability of European stone fruit yellows in Switzerland. Acta Horticulturae 1214: 239-242. <https://doi.org/10.17660/ActaHortic.2018.1214.43>
- Badmin J (2017) Red Valerian psyllid *Trioza centranthi* (Vallot) and its potential predator *Anthocoris confusus* Reuter in Kent. British Journal of Entomology and Natural History 30 (2): 78.
- Baeva VG, Kankina VK (1971) Псиллиды (Homoptera, Psylloidea) Памира. Bulletin of the Academy of Sciences of the Tajik SSR, Department of Biological Sciences 3 (44): 80-86.
- Baugnée JY (2001) Observations of some interesting psyllids in Belgium and the Great Duchy of Luxembourg (Homoptera Psylloidea). Bulletin de la Société Royale Belge d'Entomologie 137 (1-6): 20-22.
- Baugnée JY, Burckhardt D, Fassotte C (2002) Les hémiptères Psylloidea de Belgique: état des connaissances et liste actualisée. Bulletin de l'Institut royal des Sciences naturelles de Belgique, Biologie 72 (Suppl.): 125-127.
- Ben Khalifa M, Aldaghi M, Hacheche H, Kummert J, Marrakchi M, Fakhfakh H (2011) First report of 'Candidatus Phytoplasma prunorum' infecting apricots in Tunisia. Journal of Plant Pathology 93 (2): 517-519.
- Ben Othmen S, Abbes K, El Imem M, Marroquín C, Ouvrard D, Rapisarda C, Chermiti B (2019) *Bactericera trigonica* and *B. nigricornis* (Hemiptera: Psylloidea) in Tunisia as potential vectors of "Candidatus Liberibacter solanacearum" on Apiaceae. Oriental Insects 53 (4): 497-509. <https://doi.org/10.1080/00305316.2018.1536003>
- Bin F (1970) *Lygus viscicola* Put. (Miridae) e *Psylla visci* Curt. (Psyllidae), Rincoti delVischio nuovi per la fauna italiana. Bollettino Zoologia agraria Bachicoltura, Serie II 10: 133-144.
- Bleach C (2019) Plant health surveillance and incursion investigation report: July to September 2019. Surveillance 46 (4): 27-35.
- Bufaur M, Harizanova V (2015) *Trechnites psyllae* (Dalm.) and *Prionomitus mitratus* (Rusch.) as Hymenopterous Parasitoids on the Pear Sucker *Cacopsylla pyri* in Bulgaria. Jordan Journal of Agricultural Sciences 11 (4): 1160-1168.
- Buhr H (1964) Bestimmungstabellen der Gallen (Zoo- und Phytoceciden) an Pflanzen Mittel- und Nordeuropas. 1. Gustav Fischer Verlag, Jena (Germany), 761 pp.
- Buhr H (1965) Bestimmungstabellen der Gallen (Zoo- und Phytoceciden) an Pflanzen Mittel- und Nordeuropas. 2. Gustav Fischer Verlag, Jena (Germany), 811 pp.
- Burckhardt D (1983) Beiträge zur Systematik und Faunistik der schweizerischen Psylloidea (Sternorrhyncha). Entomologica Basiliensis 8: 43-83.
- Burckhardt D (1984) Revision der Gattung *Rhinocola* Förster (Homoptera: Psylloidea). Mitteilungen der Entomologischen Gesellschaft Basel 34 (4): 115-120.

- Burckhardt D, Hodkinson ID (1986) A revision of the west Palaearctic pear psyllids. Bulletin of entomological Research 76: 119-132. <https://doi.org/10.1017/S0007485300015340>
- Burckhardt D (1988a) The jumping plant lice or psyllids (Homoptera: Psylloidea) from Greece. Biologia Gallo-hellenica 13: 107-109.
- Burckhardt D (1988b) Angaben zur Psyllidenfauna der Nordosttürkei (Homoptera: Psylloidea). Mitteilungen der Entomologischen Gesellschaft Basel 38 (1): 31-44.
- Burckhardt D (1988c) Jumping plant lice (Homoptera: Psylloidea) of the temperate neotropical region. Part 3: Calophyidae and Trioziidae. Zoological Journal of the Linnean Society 92: 115-191. <https://doi.org/10.1111/j.1096-3642.1988.tb00101.x>
- Burckhardt D (1989) Les psylles (Insecta, Homoptera, Psylloidea) de l'Algérie. Archives des Sciences, Genève 42 (2): 367-424.
- Burckhardt D, Lauterer P (1989) Systematics and biology of the Rhinocolinae (Homoptera, Psylloidea). Journal of Natural History 23 (3): 643-712. <https://doi.org/10.1080/00222938900770371>
- Burckhardt D, Kofler A (1991) Blattflöhe (Homoptera, Psylloidea) aus Osttirol (Österreich). Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck 78: 103-109.
- Burckhardt D, Halperin J (1992) Additions to the psyllid fauna of Israel (Homoptera: Psylloidea). Israel Journal of Entomology 25-26: 41-50.
- Burckhardt D, Lauterer P (1993) The jumping plant-lice of Iran (Homoptera, Psylloidea). Revue Suisse de Zoologie 100 (4): 829-898. <https://doi.org/10.5962/bhl.part.79887>
- Burckhardt D, Önuçar A (1993) A review of Turkish jumping plant-lice (Homoptera, Psylloidea). Revue Suisse de Zoologie 100 (3): 547-574. <https://doi.org/10.5962/bhl.part.79887>
- Burckhardt D (1994a) Psylloid pests of temperate and subtropical crop and ornamental plants (Hemiptera, Psylloidea): A review. Entomology (Trends in Agricultural Sciences) 2: 173-186.
- Burckhardt D (1994b) Notes sur la répartition en Suisse de quelques espèces de psylles (Hemiptera, Psylloidea). Bulletin Romand d'Entomologie 12 (2): 109-115.
- Burckhardt D, Lauterer P (1997a) A taxonomic reassessment of the trioziid genus *Bactericera* (Hemiptera: Psylloidea). Journal of Natural History 31 (1): 99-153. <https://doi.org/10.1080/00222939700770081>
- Burckhardt D, Lauterer P (1997b) Systematics and biology of the *Aphalara exilis* (Weber & Mohr) species assemblage (Hemiptera: Psyllidae). Entomologica Scandinavica 28: 271-305.
- Burckhardt D (2002) Vorläufiges Verzeichnis der Blattflöhe Mitteleuropas mit Wirtspflanzenangaben. Beiträge zur Zikadenkunde 5: 1-9.
- Burckhardt D, Mifsud D (2003) Jumping plant-lice of the Paurocephalinae (Insecta, Hemiptera, Psylloidea): systematics and phylogeny. Contributions to Natural History 2: 3-34.
- Burckhardt D, Mühlenthaler R (2003) Exotische Elemente des Schweizer Blattflohfauna (Hemiptera, Psylloidea) mit einer Liste weiterer potentieller Arten. Mitteilungen der Entomologischen Gesellschaft Basel 53 (4): 98-110.
- Burckhardt D (2005) Les psyllidés, un groupe d'insectes peu connus, jamais recensés dans le Finistère. Penn ar Bed 195: 32-35.

- Burckhardt D, Lauterer P (2006) The Palaearctic triozids associated with Rubiaceae (Hemiptera, Psylloidea): a taxonomic re-evaluation of the *Trioza galii* Foerster complex. Revue Suisse de Zoologie 113 (2): 269-286. <https://doi.org/10.5962/bhl.part.80350>
- Burckhardt D, Queiroz DL (2012) Checklist and comments on the jumping plant-lice (Hemiptera: Psylloidea) from Brazil. Zootaxa 3571: 26-48. <https://doi.org/10.11646/zootaxa.3571.1.2>
- Burckhardt D, Díaz F, Queiroz DL (2017) Four new neotropical *Trioza* species associated with Loranthaceae (Santalales) and comments on mistletoe inhabiting psyllids (Hemiptera, Psylloidea). Alpine Entomology 1: 91-108. <https://doi.org/10.3897/alpineentomology.1.20905>
- Burckhardt D, Sharma A, Raman A (2018) Checklist and comments on the jumping plant-lice (Hemiptera: Psylloidea) from the Indian subcontinent. Zootaxa 4457 (1): 1-38. <https://doi.org/10.11646/zootaxa.4457.1.1>
- Burckhardt D, Ouvrard D, Percy DM (2021) An updated classification of the jumping plant-lice (Hemiptera: Psylloidea) integrating molecular and morphological evidence. European Journal of Taxonomy 736: 137-182. <https://doi.org/10.5852/ejt.2021.736.1257>
- Burrows M (2012) Jumping mechanisms in jumping plant lice (Hemiptera, Sternorrhyncha, Psyllidae). Journal of Experimental Biology 215 (20): 3612-3621. <https://doi.org/10.1242/jeb.074682>
- Burton L, Carbonnelle S, Ellis WN, Schneider N (2019) Notules cécidologiques. Observations faites au Luxembourg (2015-2018). Bulletin de la Société des Naturalistes Luxembourgeois 121: 135-151.
- Camerota C, Raddadi N, Pizzinat A, Gonella E, Crotti E, Tedeschi R, Mozes-Daube N, Ember I, Acs Z, Kolber M, Zchori-Fein E, Daffonchio D, Alma A (2012) Incidence of '*Candidatus Liberibacter europaeus*' and phytoplasmas in *Cacopsylla* species (Hemiptera: Psyllidae) and their host/shelter plants. Phytoparasitica 40 (3): 213-221. <https://doi.org/10.1007/s12600-012-0225-5>
- Carraro L, Osler R, Loi N, Ermacora P, Refatté E (1998) Transmission of European stone fruit yellows phytoplasma by *Cacopsylla pruni*. Journal of Plant Pathology 80: 233-239.
- Carraro L, Ferrini F, Labonne G, Ermacora P, Loi N (2004) Seasonal infectivity of *Cacopsylla pruni*, vector of European stone fruit yellows phytoplasma. Annals of Applied Biology 144: 191-195. <https://doi.org/10.1111/j.1744-7348.2004.tb00333.x>
- Carrières E (2003) Espèces nouvelles et remarquables de syrphes (Diptera, Syrphidae) au Luxembourg. Bulletin de la Société des Aturalistes luxembourgeois 103: 79-82.
- Castillo Carrillo CI, Jensen AS, Snyder WE (2016) Checklist of the Psylloidea (Hemiptera) of the U.S. Pacific Northwest. Proceedings of the Entomological Society of Washington 118 (4): 498-509. <https://doi.org/10.4289/0013-8797.118.4.498>
- Chireceanu C, Fătu V (2012) Data on the Hawthorn Psyllid *Cacopsylla melanoneura* (Förster) Populations in Southeast Romania. Ecologia Balkanica 4 (2): 43-49.
- Cho G, Burckhardt D, Lee S (2017a) On the taxonomy of Korean jumping plant-lice (Hemiptera: Psylloidea). Zootaxa 4238 (4): 531-561. <https://doi.org/10.11646/zootaxa.4238.4.3>
- Cho G, Burckhardt D, Inoue H, Luo X, Lee S (2017b) Systematics of the east Palaearctic pear psyllids (Hemiptera: Psylloidea) with particular focus on the Japanese and Korean fauna. Zootaxa 4362 (1): 75-98. <https://doi.org/10.11646/zootaxa.4362.1.4>

- Cho G, Malenovsky I, Burckhardt D, Inoue H, Lee S (2020) DNA barcoding of pear psyllids (Hemiptera: Psylloidea: Psyllidae), a tale of continued misidentifications. Bulletin of Entomological Research 110 (4): 521-530. <https://doi.org/10.1017/S0007485320000012>
- Cieślińska M (2011) European stone fruit yellows disease and its causal agent 'Candidatus Phytoplasma prunorum'. Journal of Plant Protection Research 51 (4): 441-447. <https://doi.org/10.2478/v10045-011-0073-1>
- Civolani S, Grandi G, Chicca M, Pasqualini E, Fano EA, Musacchi S (2013) Probing behaviour of *Cacopsylla pyri* on a resistant pear selection. Journal of Applied Entomology 137 (5): 365-375. <https://doi.org/10.1111/jen.12003>
- Cobau R (1912) Altri cecidi della valle del Brenta. Atti della Società Italiana di Scienze Naturali 51: 31-67.
- Cocquempot C (2008) Un nouveau psylle sur pittospore du Japon en France. PHM-Revue Horticole 498: 33-36.
- Conci C, Rapisarda C, Tamanini L (1993) Annotated catalogue of the Italian Psylloidea (Insecta Homoptera) - First part. Atti Accademia roveretana degli Agiati, ser. VII 2 (B): 33-135.
- Conci C, Rapisarda C, Tamanini L (1996) Annotated catalogue of the Italian Psylloidea (Insecta Homoptera) - Second part. Atti Accademia roveretana degli Agiati, ser. VII 5 (B): 5-207.
- Cooper WR, Horton DR (2015) Effects of elicitors of host plant defenses on pear psylla, *Cacopsylla pyricola*. Entomologia Experimentalis et Applicata 157 (3): 300-306. <https://doi.org/10.1111/eea.12360>
- Cravedi P, Mazzoni E, Pasqualini E, Pellizzari G, Rapisarda C, Russo A, Suma P, Tranfaglia A (2008) Psille, cocciniglie e aleirodi – Fruttiferi, agrumi, vite, olivo e orticole. Ed. L'Informatore Agrario, Milano, 184 pp. [In Italian]. [ISBN 978-88-7220-255-5]
- Cruz M, Cooper WR, Horton DR, Barcenas NM (2018) "Candidatus Phytoplasma pyri" affects behavior of *Cacopsylla pyricola* (Hemiptera: Psyllidae). Journal of Entomological Science 53 (3): 361-371. <https://doi.org/10.18474/JES17-115.1>
- Darboux G, Houard C (1901) Catalogue systématique du Zoocécidies de l'Europe et du Bassin Méditerranéen. Bulletin scientifique de la France et de la Belgique 34 (Bis): XII+1-544.
- den Bieman K, Malenovský I, Burckhardt D, Heijerman T (2019) First checklist of the Dutch jumping plant lice since 93 years (Hemiptera: Psylloidea). Nederlandse Faunistische Mededelingen 53: 55-118.
- Dietz M, Pir JB (2009) Distribution and habitat selection of *Myotis bechsteinii* in Luxembourg: Implications for forest management and conservation. Folia Zoologica 58: 327-340.
- Docters van Leeuwen WM (1937) New and noteworthy zoocecidia from the Netherlands. Marcellia 29: 73-86.
- Dohet A, Ferréol M, Cauchie H-M, Hoffmann L (2008) Caddisfly assemblages characterizing different ecological areas in Luxembourg: from geographical distribution to bioindication. Ferrantia 55: 33-56.
- Drees M (2005) Ergänzungen zur Entomofauna Germanica (Bd. 6) (Insecta: Homoptera, Rhaphidoptera, Mecoptera). Decheniana 158: 101-102.
- Drohojowska J (2009a) General information on thorax morphology of selected species of psyllids (Hemiptera, Psylloidea). In: Cichocka E (Ed.) Aphids and other hemipterous

- insects. 15. Deaprtment of Nature Preservation, The John Paul II Catholic University of Lublin, Lublin, Poland, 5-16 pp.
- Drohojowska J (2009b) Structure of head and thorax of *Livia juncorum* (Latreille, 1978) (Hemiptera, Psylloidea). In: Cichocka E (Ed.) Aphids and other hemipterous insects. 15. Deaprtment of Nature Preservation, The John Paul II Catholic University of Lublin, Lublin, Poland, 17-30 pp.
 - Drohojowska J, Burckhardt D (2014) The jumping plant-lice (Hemiptera: Psylloidea) of Turkey: a checklist and new records. Turkish Journal of Zoology 38: 1-10. <https://doi.org/10.3906/zoo-1307-15>
 - Drohojowska J, Klasa A (2019) Jumping plant lice (Hemiptera: Psylloidea) of the Ojców National Park (Southern Poland). Fragmenta Faunistica 62 (1): 27-37.
 - Eickermann M, Beyer M, Goergen K, Hoffmann L, Junk J (2014) Shifted migration of the rape stem weevil (*Ceutorhynchus napo*) linked to climate change. European Journal of Entomology 111: 243-250. <https://doi.org/10.14411/eje.2014.018>
 - Eickermann M, Ronellenfitsch F-K, Rapisarda C (2020) First record of *Psylla buxi* (L., 1758) (Hemiptera, Psylloidea) from Luxembourg. Bulletin OEPP / EPPO Bulletin 50 (2): 271-276. <https://doi.org/10.1111/epp.12636>
 - Erler F, Tosun HS (2017) Plant oils as oviposition deterrents against winterform females of pear psylla, *Cacopsylla pyri* (L.) (Hemiptera: Psyllidae). Phytoparasitica 45: 509-515. <https://doi.org/10.1007/s12600-017-0609-7>
 - Facundo R, Quiroga N, Méndez P, Zamorano A, Fiore N (2017) First report of 'Candidatus Phytoplasma pyri' on Pear in Chile. *Plant Disease* 101 (5): 830. <https://doi.org/10.1094/PDIS-08-16-1144-PDN>
 - Gallinger J, Gross J (2018) Unraveling the host plant alternation of *Cacopsylla pruni* – Adults but not nymphs can survive on conifers due to phloem/xylem composition. *Frontiers in Plant Science* 9: 484. <https://doi.org/10.3389/fpls.2018.00484>
 - Gertsson CA (2010) Fångst av bladloppor (Hemiptera: Psylloidea) från Svenska Malaisefälleprojektet - nya provinsfynd samt sällsynt förekommande arter i Sverige. *Entomologisk Tidskrift* 131 (1): 55-65.
 - Goergen K, Beersma J, Hoffmann L, Junk J (2013) ENSEMBLES-based assessment of regional climate effects in Luxembourg and their impact on vegetation. *Climatic Change* 119: 761-773. <https://doi.org/10.1007/s10584-013-0756-x>
 - Guédot C, Horton DR, Landolt PJ (2009) Attraction of male winterform pear psylla to female-produced volatiles and to female extracts and evidence of male-male repellency. *Entomologia Experimentalis et Applicata* 130 (2): 191-197. <https://doi.org/10.1111/j.1570-7458.2008.00807.x>
 - Haapalainen M (2014) Biology and epidemics of *Candidatus Liberibacter* species, psyllid transmitted plant-pathogenic bacteria. *Annals of Applied Biology* 165: 172-198. <https://doi.org/10.1111/aab.12149>
 - Haapalainen M, Wang J, Latvala S, Lehtonen MT, Pirhonen M, Nissinen AI (2018) Genetic variation of 'Candidatus Liberibacter solanacearum' haplotype C and identification of a novel haplotype from *Trioza urticae* and stinging nettle. *Phytopathology* 108 (8): 925-934. <https://doi.org/10.1094/PHYTO-12-17-0410-R>
 - Halperin J, Hodgkinson ID, Russell LM, Berlinger MJ (1982) A contribution to the knowledge of the psyllids of Israel (Homoptera: Psylloidea). *Israel Journal of Entomology* 16: 27-44.

- Halperin J, Hodkinson ID, Burckhardt D (1988) Six local psyllids (Homoptera: Psylloidea) new to the Israeli fauna. *Phytoparasitica* 16 (3): 283-284. <https://doi.org/10.1007/BF02979531>
- Hansen LO (1996) Two species of Psyllidae (Homoptera) new to Norway. *Fauna Norvegica*, Ser. B 43 (2): 98.
- Hansen LO, Greve L (1999) New records of Triozidae (Hem., Psylloidea) from Norway. *Norwegian Journal of Entomology* 46 (1): 47-48.
- Hansen LO, Hodkinson ID (2006) The mistletoe associated psyllid *Cacopsylla visci* (Curtis, 1835) (Homoptera, Psyllidae) in Norway. *Norwegian Journal of Entomology* 53 (1): 89-91.
- Hodkinson ID (1973a) The biology of *Strophingia ericae* (Curtis) (Homoptera, Psylloidea) with notes on its primary parasite *Tetrastichus actis* (Walker) (Hym., Eulophidae). *Norsk Entomologisk Tidsskrift* 20: 237-243.
- Hodkinson ID (1973b) The population dynamics and host plant interactions of *Strophingia ericae* (Curt.) (Homoptera: Psylloidea). *Journal of Animal Ecology* 42: 565-583. <https://doi.org/10.2307/3124>
- Hodkinson ID (1974) The biology of the Psylloidea (Homoptera): a review. *Bulletin of Entomological Research* 64: 325-338. <https://doi.org/10.1017/S0007485300031217>
- Hodkinson ID, White IM (1979) Homoptera Psylloidea. *Handbooks for the Identification of British Insects*. 2 (5a). Royal Entomological Society of London, London (UK), 98 pp.
- Hodkinson ID, MacLean SFJ (1980) The psyllids (Homoptera: Psylloidea) of Chukotka, Northeast USSR. *Arctic and Alpine Research* 12 (3): 377-380. <https://doi.org/10.2307/1550724>
- Hodkinson ID (1981) Heather-feeding psyllids of the genus *Strophingia* (Homoptera). *Systematic Entomology* 6 (1): 77-90. <https://doi.org/10.1111/j.1365-3113.1981.tb00017.x>
- Hodkinson ID, Hollis D (1981) The psyllids (Homoptera: Psylloidea) of Mallorca. *Entomologica Scandinavica* 12 (1): 65-77. <https://doi.org/10.1163/187631281X00382>
- Hodkinson ID, White IM (1981) The Neotropical Psylloidea (Homoptera: Insecta): an annotated checklist. *Journal of Natural History* 15 (3): 491-523. <https://doi.org/10.1080/00222938100770361>
- Hodkinson ID (1983) The psyllids (Homoptera: Psylloidea) of the Austro-Oriental, Pacific and Hawaiian zoogeographical realms: an annotated check list. *Journal of Natural History* 17: 341-377. <https://doi.org/10.1080/00222938300770251>
- Hodkinson ID (1984) The biology and ecology of the gall-forming Psylloidea (Homoptera). In: Ananthakrishnan TN (Ed.) *The biology of gall forming insects*. Oxford and IBH Publishing Co., New Delhi, Bombay, Calcutta (India), 59-77 pp.
- Hodkinson ID (1986) The psyllids (Homoptera: Psylloidea) of the Oriental Zoogeographical Region: an annotated check-list. *Journal of Natural History* 20: 299-357. <https://doi.org/10.1080/00222938600770251>
- Hodkinson ID, Hollis D (1987) The legume-feeding psyllids (Homoptera) of the west Palaearctic Region. *Bulletin of the British Museum (Natural History) - Entomology* 56 (1): 1-86.
- Hodkinson ID (1988) The Nearctic Psylloidea (Insecta: Homoptera): an annotated check list. *Journal of Natural History* 22 (5): 1179-1243. <https://doi.org/10.1080/00222938800770751>

- Hodkinson ID, Bird J (2000) Sedge and rush-feeding psyllids of the subfamily Liviinae (Insecta: Hemiptera: Psylloidea): a review. *Zoological Journal of the Linnean Society* 128 (1): 1-49. <https://doi.org/10.1111/j.1096-3642.2000.tb00648.x>
- Hodkinson ID (2009) Life cycle variation and adaptation in jumping plant lice (Insecta: Hemiptera: Psylloidea): a global synthesis. *Journal of Natural History* 43 (1): 65-179. <https://doi.org/10.1080/00222930802354167>
- Hogg BN, Smith L, Daane KM (2015) Impacts of the adventive psyllid *Arytainilla spartiophila* (Hemiptera: Psyllidae) on growth of the invasive weed *Cytisus scoparius* under controlled and field conditions in California. *Environmental Entomology* 45 (1): 109-116. <https://doi.org/10.1093/ee/nvv179>
- Hollis D, Martin JH (1997) Jumping plantlice (Hemiptera: Psylloidea) attacking avocado pear trees, *Persea americana*, in the New World, with a review of Lauraceae-feeding among psylloids. *Bulletin of Entomological Research* 87 (5): 471-480. <https://doi.org/10.1017/S000748530004133X>
- Hollis D (2004) Australian Psylloidea. Jumping plantlice and lerp insects. Australian Biological Resources Study, Canberra (Australia), 216 pp.
- Holzinger W, Aukema B, Den Bieman K, Bourgoin T, Burckhardt D, Carapezza A, Cianferoni F, Chen P-P, Faraci F, Goula M, Helden AJ, Hemala V, Huber E, Jerinić-Prodanović D, Kment P, Kunz G, Nickel H, Morkel C, Rabitsch W, Ramsay AJ, Rakauskas R, Roca-Cusachs M, Schlosser L, Seljak G, Serbina L, Soulier-Perkins A, Spodek M, Friess T (2017) Hemiptera records from Lake Spechtensee and from Southern Styria (Austria). *Entomologica Austriaca* 24: 67-82.
- Houard C (1908) Les Zoocécidies des Plantes d'Europe et du Bassin de la Méditerranée. 1. A. Hermann et fils, Paris (France), 570 pp.
- Houard C (1909) Les Zoocécidies des Plantes d'Europe et du Bassin de la Méditerranée. 2. A. Hermann et fils, Paris (France), 678 pp.
- Houard C (1913) Les Zoocécidies des Plantes d'Europe et du Bassin de la Méditerranée. 3. A. Hermann et fils, Paris (France), 312 pp.
- Inoue H (2010) The generic affiliation of Japanese species of the subfamily Psyllinae (Hemiptera: Psyllidae) with a revised checklist. *Journal of Natural History* 44 (5): 333-360. <https://doi.org/10.1080/00222930903437325>
- Jarausch B, Schwind N, Fuchs A, Jarausch W (2012) Characteristics of the spread of apple proliferation by its vector *Cacopsylla picta*. *Phytopathology* 101 (12): 1471-1480. <https://doi.org/10.1094/PHYTO-01-11-0012>
- Jarausch W, Danet JL, Labonne G, Dosba F, Broquaire JM, Saillard C, Garnier M (2001) Mapping the spread of apricot chlorotic leaf roll (ACLR) in southern France and implication of *Cacopsylla pruni* as a vector of European stone fruit yellows (ESFY) phytoplasmas. *Plant Pathology* 50: 782-790. <https://doi.org/10.1046/j.1365-3059.2001.00645.x>
- Jarausch W, Jarausch B (2016) A permanent rearing system for *Cacopsylla pruni*, the vector of 'Candidatus Phytoplasma prunorum'. *Entomologia Experimentalis et Applicata* 159 (1): 112-116. <https://doi.org/10.1111/eea.12427>
- Jarausch W, Jarausch B, Fritz M, Runne M, Etropsolska A, Pfeilstetter E (2019) Epidemiology of European stone fruit yellows in Germany: the role of wild *Prunus spinosa*. *European Journal of Plant Pathology* 154 (2): 463-476. <https://doi.org/10.1007/s10658-019-01669-3>

- Jerinić-Prodanović D, Protić L (2013) True bugs (Hemiptera, Heteroptera) as psyllid predators (Hemiptera, Psylloidea). ZooKeys 319: 169-189. <https://doi.org/10.3897/zookeys.319.4316>
- Kanturski M, Drohojowska J (2013) Koliszki (Hemiptera: Psylloidea) Garbu Tarnogórskiego. Wiadomości Entomologiczne 32 (4): 245-254.
- Kaya K, Serce CS, Gazel M, Caglayan K, Sauvion N (2016) Potential psyllid vectors of *Candidatus Phytoplasma mali* and *Candidatus Phytoplasma pyri* in Turkey. Pakistan Journal of Agricultural Science 53 (2): 383-392. <https://doi.org/10.21162/PAKJAS/16.3804>
- Kieffer JJ (1901) Synopsis des Zoocécidies d'Europe. Annales de la Société entomologique de France 70: 233-579.
- Kwon YJ (1983) Psylloidea of Korea. Series 2. Insecta Koreana, Seoul (South Korea), 181 pp.
- Labina ES (2006) Psyllids of the Kuril Islands (Homoptera: Psylloidea). Zoosystematica Rossica 15 (1): 51-55. <https://doi.org/10.31610/zsr/2006.15.1.51>
- Labina ES (2008) The jumping plant lice (Homoptera, Psyllinea) fauna of the Republic of Altai. Entomological Review 88 (1): 277-285. <https://doi.org/10.1134/S0013873808030032>
- Labina ES, Evdogarova T (2009) Jumping plant lice (Homoptera, Psyllinea) from the relict steppes of Central Yakutia. Entomological Review 89 (9): 1069-1071. <https://doi.org/10.1134/S0013873809090073>
- Lambinon J, Schneider N (2004) Contribution à la connaissance des galles d'hémiptères (Insecta, Hemiptera) du Luxembourg. Bulletin de la Société des naturalistes luxembourgeois 105: 109-132.
- Lambinon J, Verlooove F (2012) Nouvelle Flore de la Belgique, du Grande-Duché de Luxembourg, du Nord de la France et des Régions Voisines: Ptéridophytes et Spermatophytes. 6th Edition. National Botanic Garden of Belgium, 1195 pp.
- Lambinon J, Schneider N, Feitz F (2012) Contribution complémentaire à la connaissance des zoocécidies du Luxembourg. Bulletin de la Société des Naturalistes Luxembourgeois 113: 167-187.
- Lauterer P, Baudys E (1968) Description of a new gall on *Chamaenerion angustifolium* (L.) Scop, produced by the larva of *Craspedolepta subpunctata* (Först.), with notes on the bionomics of this Psyllid. Acta Musei Moraviae, Scientiae Naturales 53: 243-248.
- Lauterer P (1976) Psyllids of wetland nature reserves of the German Democratic Republic, with notes on their biology, taxonomy and zoogeography (Homoptera, Psylloidea). Faunistische Abhandlungen 6 (10): 111-122.
- Lauterer P (1979) New and interesting records of psyllids from Czechoslovakia (Homoptera, Psylloidea). Acta Musei Moraviae, Scientiae Biologicae 64: 93-102.
- Lauterer P (1991) Psyllids (Homoptera, Psylloidea) of the limestone cliff zone of the Pavlovské vrchy Hills (Czechoslovakia). Acta Musei Moraviae, Scientiae Biologicae 76 (1-2): 241-263.
- Lauterer P (1994) Psyllids (Homoptera, Psylloidea) from the area flooded by the Nové Mlýny reservoir system and its environs in southern Moravia. Acta Musei Moraviae, Scientiae Naturales 78 (1-2): 165-200.
- Lauterer P, Burckhardt D (1997) Central and West European willow-feeding jumping plant-lice of the genus *Cacopsylla* (Hemiptera: Psylloidea). Entomological Problems 28 (2): 81-94.

- Lauterer P (1999) Results of the investigations on Hemiptera in Moravia, made by the Moravian Museum (Psylloidea 2). *Acta Musei Moraviae, Scientiae Biologicae* 84 (1-2): 71-151.
- Lauterer P (2001) Mery (Sternorrhyncha, Psylloidea) Jizerských hor. Sborník Severočeského Muzea, Přírodní Vědy 22: 85-99.
- Lauterer P (2011) Psyllids (Hemiptera, Psylloidea) inhabiting the canopy and shrub layer of an oak and hornbeam forest in Báb near Nitra (Slovakia). *Folia Faunistica Slovaca* 16 (2): 97-101.
- Li F (2011) Psyllidomorpha of China (Insecta: Hemiptera). 1-2. Huayu Nature Book Trade Co. Ltd, Beijing (China), 1976 pp.
- Loginova MM (1968) New data on the fauna and biology of the Caucasian Psylloidea (Homoptera). Trudy Vsesoyuznogo Entomologicheskogo Obshchestva, Akademiya Nauk SSSR 52: 275-328.
- Loginova MM (1974) The psyllids (Psylloidea, Homoptera) of the Mongolian People's Republic. II. Nasekomye Mongoli 4 (2): 51-66.
- Loginova MM (1977) [A review of the genus *Eryngiofaga* Klimasz. (Homoptera, Triozidae)]. *Zoologicheskii Zhurnal* 56 (4): 510-521.
- Malenovský I, Baňař P, Kment P (2011) A contribution to the faunistics of the Hemiptera (Cicadomorpha, Fulgoromorpha, Heteroptera, and Psylloidea) associated with dry grassland sites in southern Moravia (Czech Republic). *Acta Musei Moraviae, Scientiae Biologicae* 96 (1): 41-187.
- Malenovský I, Lauterer P (2012) Jumping plant-lice (Hemiptera: Psylloidea) of the Bílé Karpaty Protected Landscape Area and Biosphere Reserve (Czech Republic). *Acta Musei Moraviae, Scientiae Biologicae* 96 (2): 105-154.
- Malenovský I, Lauterer P, Labina ES, Burckhardt D (2012) Jumping plant-lice (Hemiptera: Psylloidea) of Afghanistan. *Acta Entomologica Musei Nationalis Pragae* 52 (1): 1-22.
- Malenovský I, Kment P, Sychra J (2014) [True bugs, leafhoppers, planthoppers and psyllids (Hemiptera: Heteroptera, Auchenorrhyncha, Psylloidea) of the environs of Prebuž in the Krusne hory Mountains (Czech Republic)]. *Klapalekiana* 50 (3-4): 181-234.
- Malumphy C, Ostrauskas H, Pye D (2009) Contribution to the knowledge of jumping plant-lice (Hemiptera, Psylloidea) of Lithuania. *Acta Zoologica Lituanica* 19 (2): 128-131. <https://doi.org/10.2478/v10043-009-0015-6>
- Marchini D, Del Bene G, Viscuso R, Dallai R (2012) Sperm Storage by Spermatodoses in the Spermatheca of *Trioza alacris* (Flor, 1861) Hemiptera, Psylloidea, Triozidae: A Structural and Ultrastructural Study. *Journal of Morphology* 273 (2): 195-210. <https://doi.org/10.1002/jmor.11017>
- Marcone C, Jarausch B, Jarausch W (2010) 'Candidatus Phytoplasma prunorum', the causal agent of European stone fruit yellows: An overview. *Journal of Plant Pathology* 92: 19-34.
- Marie-Jeanne V, Bonnot F, Thébaud G, Peccoud J, Labonne G, Sauvion N (2020) Multi-scale spatial genetic structure of the vector-borne pathogen 'Candidatus Phytoplasma prunorum' in orchards and in wild habitats. *Scientific Reports* 10: 5002. <https://doi.org/10.1038/s41598-020-61908-0>

- Martoni F, Bulman S, Pitman A, Taylor G, Armstrong K (2018) DNA Barcoding highlights cryptic diversity in the New Zealand Psylloidea (Hemiptera: Sternorrhyncha). *Diversity* 10 (3): 50. <https://doi.org/10.3390/d10030050>
- Mathur RN (1975) Psyllidae of the Indian Subcontinent. Indian Council of Agricultural Research, New Delhi (India), 429 pp.
- Mayer CJ, Jarausch B, Jarausch W, Jelkmann W, Vilcinskas A, Gross J (2009) *Cacopsylla melanoneura* has no relevance as vector of apple proliferation in Germany. *Phytopathology* 99 (6): 729-738. <https://doi.org/10.1094/PHYTO-99-6-0729>
- Mifsud D, Cocquempot C, Mühlenthaler R, Wilson M, Streito JC (2010) Other Hemiptera Sternorrhyncha (Aleyrodidae, Phylloxeroidea, and Psylloidea) and Hemiptera Auchenorrhyncha. In: Roques A, et al. (Ed.) Alien terrestrial arthropods of Europe. Chapter 9.4. BioRisk 4 (1), 511-552 pp. <https://doi.org/10.3897/biorisk.4.63>
- Miñarro M, Somoano A, Moreno A, García RR (2016) Candidate insect vectors of apple proliferation in Northwest Spain. *SpringerPlus* 5: 1240. <https://doi.org/10.1186/s40064-016-2907-9>
- Munyaneza JE (2010) Psyllids as vectors of emerging bacterial diseases of annual crops. *Southwestern Entomologist* 35: 471-477. <https://doi.org/10.3958/059.035.0335>
- Mustafa SA, Zubair SM, Zandi ZA, Al-Maroof IN, Kidir A, Anwer AM (2014) Preliminary survey of economic insects and their insect predators in northern Iraq. *Munis Entomology & Zoology* 9 (1): 150-160.
- Newbould J (2012) A selection of interesting plant galls in Yorkshire in 2011. *Naturalist* 137 (1079): 48-50.
- Nokkala S (1995) *Camarotoscena speciosa* (Flor, 1861) (Homoptera, Psylloidea), a new species to Finland. *Sahlbergia* 2 (2): 126.
- O'Connor JP, Ashe P, Wistow S (1997) *Trioza alacris* Flor (Hemiptera), a gall-causing psyllid new to Ireland. *Entomologist's Record and Journal of Variation* 109 (5-6): 135-136.
- O'Connor JP, Malumphy C (2011) A review of the Irish jumping plant-lice (Hemiptera: Psylloidea). *Bulletin of the Irish Biogeographical Society* 35: 21-63.
- Oettl S, Schlink K (2015) Molecular identification of two vector species, *Cacopsylla melanoneura* and *Cacopsylla picta* (Hemiptera: Psyllidae), of apple proliferation disease and further common psyllids of Northern Italy. *Journal of Economic Entomology* 108 (5): 2174-2183. <https://doi.org/10.1093/jee/tov204>
- Önuçar A, Ulu O (1991) Five jumping plant lice species (Homoptera, Psylloidea) new for Turkish fauna. *Turkiye Entomoloji Dergisi* 15 (3): 153-155.
- Oppedisano T, Panassiti B, Pedrazzoli F, Mittelberger C, Bianchedi PL, Angeli G, De Cristofaro A, Janik K, Anfora G, Ioriatti C (2020) Importance of psyllids' life stage in the epidemiology of apple proliferation phytoplasma. *Journal of Pest Science* 93: 49-61. <https://doi.org/10.1007/s10340-019-01130-8>
- Ossiannilsson F (1992) The Psylloidea (Homoptera) of Fennoscandia and Denmark. 26. *Fauna Entomologica Scandinavica*, E.J. Brill, Leiden (The Netherlands), 346 pp.
- Ouvrard D, Burckhardt D, Cocquempot C (2015) An annotated checklist of the jumping plant-lice (Insecta: Hemiptera: Psylloidea) from the Mercantour National Park, with seven new records for France and one new synonymy. *Zoosystema* 37 (1): 251-271. <https://doi.org/10.5252/z2015n1a13>
- Ouvrard D (2021) Psyl'list - The World Psylloidea Database. <http://www.hemiptera-databases.com/psyllist>. Accessed on: 2021-2-27.

- Park HC (1996) Taxonomy of Korean psyllids (Homoptera: Psylloidea) 1. A revised checklist. *Korean Journal of Entomology* 26 (3): 267-278.
- Parkinson JD, Whittaker JB (1975) A study of two physiological races of the heather psyllid, *Strophingia ericae* (Curtis) (Homoptera: Psylloidea). *Biological Journal of the Linnean Society* 7 (1): 73-81. <https://doi.org/10.1111/j.1095-8312.1975.tb00734.x>
- Peccoud J, Labonne G, Sauvion N (2013) Molecular test to assign individuals within the *Cacopsylla pruni* complex. *PLOS One* 8: e72454. <https://doi.org/10.1371/journal.pone.0072454>
- Peccoud J, Pleydell DR, Sauvion N (2018) A framework for estimating the effects of sequential reproductive barriers: implementation using Bayesian models with field data from cryptic species. *Evolution* 72: 2503-2512. <https://doi.org/10.1111/evo.13595>
- Pekár S, Michalko R, Loverre P, Líznarová E, Černecká L' (2015) Biological control in winter: novel evidence for the importance of generalist predators. *Journal of Applied Ecology* 52 (1): 270-279. <https://doi.org/10.1111/1365-2664.12363>
- Percy D, Rung A, Hoddle MS (2012) An annotated checklist of the psyllids of California (Hemiptera: Psylloidea). *Zootaxa* 3193: 1-27. <https://doi.org/10.11646/zootaxa.3193.1.1>
- Pizzinat A, Tedeschi R, Alma A (2011) *Cacopsylla melanoneura* (Foerster): aestivation and overwintering habitats in Northwest Italy. *Bulletin of Insectology* 64 (Supplement): S135-S136.
- Raddadi N, Gonella E, Camerota C, Pizzinat A, Tedeschi R, Crotti E, Mandrioli M, Bianco PA, Daffonchio D, Alma A (2011) 'Candidatus Liberibacter europaeus' sp. nov. that is associated with and transmitted by the psyllid *Cacopsylla pyri* apparently behaves as an endophyte rather than a pathogen. *Environmental Microbiology* 13: 414-426. <https://doi.org/10.1111/j.1462-2920.2010.02347.x>
- Ramírez Gómez C (1956) Los Psilidos de España. *Boletín de la Real Sociedad Española de Historia Natural* (Secc. Biol.) 53: 151-217.
- Rapisarda C (1985) Notizie preliminari sulla psillidofauna della Sicilia. In: Mineo G (Ed.) Atti XIV Congresso nazionale italiano di Entomologia. Palermo-Erice-Bagheria (Italy), 28/5 - 1/6/1985. 111-117 pp.
- Rapisarda C (1994) Additions to the Sicilian psyllid fauna (Homoptera Psylloidea). *Naturalista siciliano* 18 (1-2): 161-173.
- Rapisarda C, Weigand A, Braun P, Eickermann M (2021) Inventory of the jumping plant lice of Luxembourg (Hemiptera; Sternorrhyncha; Psylloidea). Version 1.8. National Museum of Natural History, Luxembourg. URL: <https://doi.org/10.15468/svhf53>
- Riedle-Bauer M, Paleskic C, Brader G (2019a) Investigations on pear psyllids as vectors of 'Candidatus Phytoplasma pyri' in Austria. *Phytopathogenic Mollicutes* 9 (1): 99-100. <https://doi.org/10.5958/2249-4677.2019.00050.1>
- Riedle-Bauer M, Paleskic C, Schwanzer J, Kölber M, Bachinger K, Antonielli L, Schönhuber C, Elek R, Stradinger J, Emberger M, Engel C, Makay M, Zajcsek F, Brader G (2019b) Epidemiological and molecular study on 'Candidatus Phytoplasma prunorum' in Austria and Hungary. *Annals of Applied Biology* 175 (3): 400-414. <https://doi.org/10.1111/aab.12541>
- Riedle-Bauer M, Paleskic C, Schönhuber C, Staples M, Brader G (2022) Vector transmission and epidemiology of 'Candidatus Phytoplasma pyri' in Austria and identification of *Cacopsylla pyrisuga* as new pathogen vector. *Journal of Plant Diseases and Protection* 129: 375-386.

- Ripka G (2008) Checklist of the Psylloidea of Hungary (Hemiptera: Sternorrhyncha). *Acta Phytopathologica et Entomologica Hungarica* 43 (1): 121-142. <https://doi.org/10.1556/APhyt.43.2008.1.14>
- Ripka G, Kiss B (2008) Recent data to the knowledge on psyllid species (Hemiptera: Psylloidea) occurring on common ragweed in Hungary. *Növényvédelem* 44 (6): 257-261.
- Ripka G (2009) Additional data to the aphid and psylloid fauna of Hungary (Hemiptera: Sternorrhyncha). *Acta Phytopathologica et Entomologica Hungarica* 44 (2): 397-417. <https://doi.org/10.1556/APhyt.44.2009.2.16>
- Ripka G (2012) New psyllid records from Hungary (Hemiptera: Psylloidea). *Acta Phytopathologica et Entomologica Hungarica* 47 (1): 75-78. <https://doi.org/10.1556/APhyt.47.2012.1.9>
- Ross H, Hedicke H (1927) Die Pflanzengallen Mittel- und Nord-europas, ihre Erreger un Biologie un Bestimmungstabelle. G. Fischer, Jena (Germany), 348 pp.
- Sampò A (1975) Di alcuni Psillocecdidi nuovi o poco noti della Valle d'Aosta. *Revue Valdôtaine d'Histoire Naturelle* 29: 153-174.
- Sampò A (1977) La psilla del Lauro (*Trioza alacris* Flor). II *Floriculture* 14 (1): 79, 82-83.
- Sanchez JA, Ortín-Angulo MC (2011) Sampling of *Cacopsylla pyri* (Hemiptera: Psyllidae) and *Pilophorus gallicus* (Hemiptera: Miridae) in Pear Orchards. *Journal of Economic Entomology* 104 (5): 1742-1751. <https://doi.org/10.1603/EC11040>
- Sanchez JA, Ortín-Angulo MC (2012) Abundance and population dynamics of *Cacopsylla pyri* (Hemiptera: Psyllidae) and its potential natural enemies in pear orchards in southern Spain. *Crop Protection* 32: 24-29. <https://doi.org/10.1016/j.cropro.2011.11.003>
- Sauvion N, Lachenaud O, Genson G, Rasplus J-Y, Labonne G (2007) Are there several biotypes of *Cacopsylla pruni*? *Bulletin of Insectology* 60 (2): 185-186.
- Sauvion N, Peccoud J, Meynard CN, Ouvrard D (2021) Occurrence data for the two cryptic species of *Cacopsylla pruni* (Hemiptera: Psylloidea). *Biodiversity Data Journal* 9: e68860. <https://doi.org/10.3897/BDJ.9.e68860>
- Schneider N (2016) Notules cécidologiques. Observations faites au Luxembourg (2000-2015). *Bulletin de la Société des Naturalistes Luxembourgeois* 118: 135-143.
- Seljak G (2006) An overview of the current knowledge of jumping plant-lice of Slovenia (Hemiptera: Psylloidea). *Acta Entomologica Slovenica* 14 (1): 11-34.
- Seljak G, Malenovský I, Lauterer P (2008) New records of jumping plant-lice from Slovenia with description of *Bactericera lyrata* sp. n. (Hemiptera: Psylloidea). *Revue Suisse de Zoologie* 115 (3): 527-540. <https://doi.org/10.5962/bhl.part.80442>
- Serbina L, Burckhardt D, Borodin O (2015) The jumping plant-lice (Hemiptera: Psylloidea) of Belarus. *Revue Suisse de Zoologie* 122 (1): 7-44.
- Sertkaya G, Martini M, Ermacora P, Musetti R, Osler R (2005) Detection and characterization of phytoplasmas in diseased stone fruits and pear by PCR-RFLP analysis in Turkey. *Phytoparasitica* 33 (4): 380-390. <https://doi.org/10.1007/BF02981306>
- Spodek M, Burckhardt D, Freidberg A (2017) The Psylloidea (Hemiptera) of Israel. *Zootaxa* 4276 (3): 301-345. <https://doi.org/10.11646/zootaxa.4276.3.1>
- Spooner BM (2016) *Trioza centranthi* (Psylloidea: Triozidae) in Britain and a record from Surrey. *Cecidology* 31: 40-44.
- Steffek R, Follak S, Sauvion N, Labonne G, MacLeod A (2012) Distribution of 'Candidatus Phytoplasma prunorum' and its vector *Cacopsylla pruni* in European fruit-

- growing areas: a review. Bulletin OEPP/EPPO Bulletin 42 (2): 191-202. <https://doi.org/10.1111/epp.2567>
- Stevens A, Udelhoven Z, Denis A, Tychon B, Lioy R, Hoffmann L, van Wesemael B (2010) Measuring solid organic carbon in croplands at regional scale using airborne imaging spectroscopy. Geoderma 158: 32-45. <https://doi.org/10.1016/j.geoderma.2009.11.032>
 - Struwe I, Gertsson CA, Coulianos CC (2009) Insects monophagous on mistletoe (*Viscum album* L.) newly discovered in Sweden: *Cacopsylla visci* (Curtis, 1835) (Hemiptera, Psyllidae) and *Pinalitus viscidola* (Puton, 1888) (Hemiptera, Miridae). Entomologisk Tidskrift 130 (3-4): 155-160.
 - Syrett P, Fowler SV, Harman HM, Hayes LM, Memmott J, Sheat JJ (2007) Establishment of *Arytainilla spartiophila* Förster (Hemiptera: Psyllidae), a new biological control agent for broom, *Cytisus scoparius*, in New Zealand. New Zealand Entomologist 30: 53-62. <https://doi.org/10.1080/00779962.2007.9722151>
 - Tedeschi R, Lauterer P, Bertignono L, Alma A (2008) Hawthorn psyllid fauna in northwestern Italy. Bulletin of Insectology 61 (1): 143-144.
 - Tedeschi R, Lauterer P, Brusetti L, Tota F, Alma A (2009) Composition, abundance and phytoplasma infection in the hawthorn psyllid fauna of northwestern Italy. European Journal of Plant Pathology 123: 301-310. <https://doi.org/10.1007/s10658-008-9367-1>
 - Tedeschi R, Nardi F (2010) DNA-based discrimination and frequency of phytoplasma infection in the two hawthorn-feeding species, *Cacopsylla melanoneura* and *Cacopsylla affinis*, in northwestern Italy. Bulletin of Entomological Research 100 (6): 741-747. <https://doi.org/10.1017/S0007485310000118>
 - Tedeschi R, Baldessari M, Mazzoni V, Trona F, Angeli G (2012) Population dynamics of *Cacopsylla melanoneura* (Hemiptera: Psyllidae) in northeast Italy and its role in the apple proliferation epidemiology in apple orchards. Journal of Economic Entomology 105 (2): 322-328. <https://doi.org/10.1603/EC11237>
 - Tedeschi R, Jarausch B, Delic D, Weintraub P (2013) Actual distribution of fruit tree and grapevine phytoplasma diseases and their vectors in Europe and neighboring regions. Phytopathogenic Mollicutes 3 (1): 3-4. <https://doi.org/10.5958/j.2249-4677.3.1.002>
 - Tishetshkin DY (2007) The possibility to use bioacoustic characters in the taxonomy of the jumping plant lice with an example of the genus *Craspedolepta* (Homoptera, Psyllinea, Aphalaridae) and description of a new species from Transbaikalia. Entomological Review 87 (5): 561-570. <https://doi.org/10.1134/S0013873807050065>
 - Tomasi E (2003) Gall-making organisms in the area of Muggia and the Noghere lakes (Muggia, Trieste, noth north eastern Italy). Atti Museo civico di Storia naturale di Trieste 50: 287-30.
 - Valasevich N, Schneider B (2016) Detection, identification and molecular diversity of 'Candidatus Phytoplasma prunorum' in Belarus. Journal of Plant Pathology 98 (3): 625-629.
 - Varga I, Keresztes B, Poczai P (2012) Data to the Hungarian insect fauna of European mistletoe (*Viscum album*). Növényvédelem 48 (4): 153-164.
 - Wagner W, Franz H (1961) Unterordnung Homoptera, Überfamilie Sternorrhyncha (Psylloidea). In: Franz H (Ed.) Die Nordost-Alpen im Spiegel ihrer Landtierwelt. 2. Universitätsverlag Wagner, Innsbruck (Austria), 158-179 pp.

- Warabieda W, Soika G, Cieślińska M (2017) *Cacopsylla pruni* in Poland and its significance as a vector of ‘*Candidatus Phytoplasma prunorum*’. Zemdirbyste-Agriculture 105 (2): 177-182. <https://doi.org/10.13080/z-a.2018.105.023>
- Wheeler AGJ, Stoops CA (2001) *Cacopsylla peregrina* (Foerster) (Sternorrhyncha: Psylloidea: Psyllidae): First U.S. Records of an Old World specialist on hawthorns (*Crataegus* spp.). Proceedings of the Entomological Society of Washington 103 (1): 103-109.
- Wheeler AGJ, Hoebeke ER (2005) *Livilla variegata* (Löw) (Hemiptera: Sternorrhyncha: Psyllidae) new to North America, with records of three other palearctic psyllids new to Newfoundland. Proceedings of the Entomological Society of Washington 107 (4): 941-946.
- Wheeler AGJ (2017) New records of a European psyllid, *Arytainilla spartiophila* (Foerster) (Hemiptera: Psyllidae), in the Southeastern United States. Proceedings of the Entomological Society of Washington 119 (2): 264-270. <https://doi.org/10.4289/0013-8797.119.2.264>
- Yang CT (1984) Psyllidae of Taiwan. Series 3. Taiwan Museum Special Publication, 305 pp.
- Zangheri S (1954) Nota sulla *Trioza urticae* L. Bollettino Istituto Entomologia Università Bologna 20: 257-273.
- Zeidan-Gèze N, Burckhardt D (1998) The jumping plant-lice of Lebanon (Hemiptera: Psylloidea). Revue Suisse de Zoologie 105 (4): 797-812.
- Zendedel A, Burckhardt D, Fekrat L, Manzari S, Namaghi HS (2016) An annotated checklist of the jumping plant-lice (Hemiptera: Psylloidea) of Iran. Journal of the Entomological Research Society 18 (1): 37-55.