



# New records of amphibians from Ha Tinh Province, Vietnam

Vinh Quang Dau<sup>‡</sup>, Cuong The Pham<sup>§</sup>, Truong Quang Nguyen<sup>§</sup>, Toan Canh Thai<sup>‡</sup>, Anh Dinh Tran<sup>‡</sup>, Anh Van Pham<sup>¶</sup>

<sup>‡</sup> Hong Duc University, 565 Quang Trung Road, Dong Ve Ward, Thanh Hoa City, Vietnam

<sup>§</sup> Institute of Ecology and Biological Resources and Graduate University of Science and Technology, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet Road, Cau Giay, Hanoi, Vietnam

<sup>‡</sup> Vu Quang National Park, Vu Quang Town, Vu Quang District, Ha Tinh Province, Ha Tinh, Vietnam

<sup>¶</sup> Faculty of Environmental Sciences, University of Science, Vietnam National University, Hanoi, 334 Nguyen Trai Road, Hanoi City, Vietnam

Corresponding author: Anh Van Pham ([phamanh@hus.edu.vn](mailto:phamanh@hus.edu.vn))

Academic editor: Franco Andreone

Received: 08 Mar 2024 | Accepted: 18 Apr 2024 | Published: 08 May 2024

Citation: Dau V, Pham C, Nguyen TQ, Thai T, Tran A, Pham A (2024) New records of amphibians from Ha Tinh Province, Vietnam. Biodiversity Data Journal 12: e122598. <https://doi.org/10.3897/BDJ.12.e122598>

## Abstract

## Background

Since the establishment of the Vu Quang National Park in 2002 in Ha Tinh Province, central Vietnam, several surveys on the amphibian fauna have been undertaken in this protected area. In 2009, Nguyen et al. provided a list of 23 amphibian species from Vu Quang National Park. In addition, a new species was described in 2021 from the National Park, namely *Vietnamophryne vuquangensis*.

## New information

As a result of our field surveys in 2013 and 2023, a total of 29 species of amphibians was recorded from the Vu Quang National Park. Six of them are recorded for the first time from Ha Tinh Province, comprising three species of Megophryidae, one species of

Dicroglossidae, one species of Ranidae and one species of Rhacophoridae. In addition, we provide morphological data and ecological notes on the aforementioned species.

## Keywords

distribution, frogs, morphology, taxonomy, Vu Quang National Park

## Introduction

The Vu Quang National Park (NP), located in Ha Tinh Province of Vietnam, was established by Decision No. 102/TTg of the Prime Minister on 30 July 2002 with an area of 55,029 hectares (Tordoff et al. 2004). The Park was recognised as an ASEAN Heritage Park in October 2019. Vu Quang NP is situated in the northern part of the Truong Son Range at elevations between 100 and 2,200 m.

In terms of amphibian diversity, Ha Tinh Province is one of the most poorly-studied provinces in Vietnam. In their herpetofaunal list of Vietnam, Nguyen et al. (2009) recorded 26 species of amphibians from Ha Tinh Province and most of them were reported from Vu Quang NP and Ho Ke Go Nature Reserve (Suppl. material 1). Further new records of amphibians from Ha Tinh Province were published by Dau et al. (2015), Thai (2017), Ziegler et al. (2022) and Sheridan et al. (2023) (Suppl. material 1). Most recently, a new species was described from Vu Quang NP, viz. *Vietnamophryne vuquangensis* Hoang, Jiang, Nguyen, Orlov, Le, Nguyen, Nguyen, Nguyen, Nguyen and Ziegler (Hoang et al. 2021).

Based on our recent fieldwork in Vu Quang NP, Ha Tinh Province, Vietnam in 2013 and 2023, we herein report six new provincial records of amphibians for Ha Tinh Province.

## Materials and methods

### Sampling

Field surveys were conducted in Vu Quang NP, Ha Tinh Province, Vietnam from 15 to 30 June 2013 by Dau QV, Thai CT, Nguyen VH and Tran DA and from 18 to 27 December 2023 by Pham VA and Tran DA. The coordinates (WGS 84) and elevations were determined by using the GPS Garmin 62SX (Fig. 1).

Specimens were collected by hand between 19:00 and 22:00 h pm. After taking live photographs, specimens were euthanised in a closed vessel with a piece of cotton wool containing ethyl acetate (Simmons 2002), fixed in 80% ethanol for five hours and then transferred to 70% ethanol for permanent storage. Voucher specimens were subsequently deposited in the collections of the Faculty of Natural Sciences, Hong Duc University (HDU), Thanh Hoa Province; Faculty of Environmental Sciences, University of Science, Vietnam

National University, Hanoi (VNU) and Vu Quang National Park (VQNP), Ha Tinh Province, Vietnam.




Figure 1. doi

New distribution records of amphibians in Vu Quang National Park, Ha Tinh Province, Vietnam: 1 = *Leptobrachium masatakasatoj*; 2 = *Xenophrys* cf. *parva*; 3 = *Xenophrys lancangica*; 4 = *Quasipaa* cf. *verrucospinosa*; 5, 6 = *Amolops compotrix*; 7, 8 = *Kurixalus odontotarsus*.

### Morphological characters

Measurements were taken with a digital caliper to the nearest 0.1 mm. Abbreviations are as follows: (SVL) snout-vent length; (HL) head length, from posterior corner of mandible to tip of snout; (HW) maximum head width, at the angle of jaws; (RL) rostral length, from anterior corner of orbit to tip of snout; (IN) internarial distance; (NS) distance from anterior edge of nostril to tip of snout; (EN) distance from anterior corner of eye to posterior edge of nostril; (ED) eye length, from anterior corner to posterior corner of eye; (IOD) minimum distance between upper eyelids; (UEW) maximum width of upper eyelid; (TD) maximum tympanum diameter; (TYE) tympanum-eye distance, from anterior margin of tympanum to posterior corner of the eye; (HAL) hand length, from elbow to tip of third finger; (FL) thigh length, from vent to knee; (TbL) shank length. For webbing formula, we followed Glaw and Vences (2007). Sex was determined by gonadal inspection in these frogs are all adults. (JJLR) Jodi J. L. Rowley; DQV (Dau Quang Vinh); HT (Ha Tinh).

## Taxon treatments

### *Leptobrachium masatakasatoi* Matsui, 2013

#### Material

- a. scientificName: *Leptobrachium masatakasatoi*; scientificNameID: *Leptobrachium masatakasatoi*; class: Amphibia; order: Anura; family: Megophryidae ; genus: *Leptobrachium*; specificEpithet: *masatakasatoi*; scientificNameAuthorship: Matsui, 2013; country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 1294 m; verbatimLatitude: 18°14'13.67"N; verbatimLongitude: 105°25'54.23"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: HDU 03111; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: 2CE20B14-6A4E-5F9F-99AF-C88DC30BA948

#### Description

Morphological characters of the specimen from Ha Tinh Province matched descriptions of Matsui (2013) and Pham et al. (2016): Body elongate, SVL 58.6 mm in male (n = 1). Head wider than long (HL 25.6 mm, HW 26.1 mm); snout round, barely projecting beyond upper jaw (RL 10.5 mm), longer than horizontal diameter of eye (ED 8.2 mm); canthus sharp, straight; loreal region oblique, moderately concave; nostril lateral, closer to tip of snout than to eye (NS 4.8 mm, EN 5.7 mm); interorbital space flat, broader than upper eyelid and internarial distance (IOD 7.1 mm, UEW 6.4 mm, IN 4.4 mm); tympanum indistinct; vomerine teeth absent; tongue heart-shaped, notched posteriorly; vocal openings absent. Fore-limb long (HAL 33.7 mm); relative finger lengths II < IV < I < III; fingers without dermal fringe, free of webbing; tips of fingers round, slightly swollen; palmar tubercles two, oval; nuptial pads absent. Hind-limb slender, thigh longer than tibia (FL 26.7 mm, TbL 22.6 mm, TbL/SVL 38.5%); relative toe lengths I < II < V < III < IV; tips of toes slightly swollen; webbing formula I1–1II1–1III1–2IV2–1V; tibiotarsal articulation reaching to posterior margin of tympanum when limb adpressed along body; inner metatarsal tubercle distinct, as long as of toe I.

Skin: Dorsal surface with fine network of ridges, tubercles present in the posterior region of sacrum, more distinct in anterior part of vent; upper lip without spines in male; supratympanic fold present, from posterior edge of eye to axilla; flanks granular; throat and chest asperities; belly and ventral surfaces of limbs smooth, except for granular thigh.

Colouration in life: Dorsal surface of head reddish-brown, with some dark spots on upper eyelid, back with irregularly black spots; supratympanic fold edged in black below; flank light brown with large black spots; anterior part of thigh with large dark spots; dorsal limb light brown with dark crossbars; ventral surface with irregular brown and cream markings (Fig. 2).



Figure 2. [doi](#)

*Leptobrachium masatakasatoi* (adult male) from Ha Tinh Province, Vietnam. Photo by V. Q. Dau.

### Distribution

In Vietnam, this species was recorded from Son La and Nghe An Provinces (Pham et al. 2016, Do et al. 2017). Elsewhere, this species is known from Laos (Frost 2024).

### Ecology

The specimen was found on the ground near a small stream at 19:15 h pm. The surrounding habitat was evergreen forest of large hardwood and shrubs.

## *Xenophrys cf. parva* (Boulenger, 1893)

### Material

- a. scientificName: *Xenophrys cf. parva*; scientificNameID: *Xenophrys cf. parva*; class: Amphibia; order: Anura; family: Megophryidae; genus: *Xenophrys*; specificEpithet: *parva*; scientificNameAuthorship: (Boulenger, 1893); country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 1214 m; verbatimLatitude: 18°14'36.82"N; verbatimLongitude: 105°25'56.42"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: JCLR 00043; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: BF6DE9C4-2C11-5A70-A930-B1BAF9009DB3

### Description

Morphological characters of the specimen from Ha Tinh Province relatively matched descriptions of Boulenger (1893), Bourret (1942), Fei et al. (2009), Pham et al. (2019a)

and Lyu et al. (2023): SVL 41.8 mm in male ( $n = 1$ ); head longer than wide (HL 15.4, HW 15.0 mm, HL/SVL 36.8%, HW/SVL 35.9%); snout obliquely truncate, projecting beyond upper jaw (RL 5.5 mm), longer than horizontal diameter of eye (ED 4.8 mm); loreal region concave; canthus rostralis well developed; nostril lateral, slightly closer to eye than to tip of snout (NS 2.8 mm, EN 2.5 mm); interorbital space flat, broader than upper eyelid and narrower than internarial distance (IOD 4.3 mm, UEW 3.9 mm, IN 4.8 mm); tympanum distinct (TD 3.0 mm); vomerine teeth present; tongue heart-shaped, round posteriorly; vocal in male present.

Fore-limb slender (HAL 15.7 mm); relative finger lengths  $I \leq II < IV < III$ ; fingers without dermal fringe, free of webbing; tips of fingers round, not swollen; subarticular tubercles indistinct; palmar tubercles indistinct; nuptial pads absent in males. Hind-limb slender, tibia longer than thigh (TbL 19.8 mm, FL 18.5 mm); relative toe lengths  $I < II < V < III < IV$ ; tips of toes round, slightly swollen; webbing rudiment; toes dermal fringe absent; metatarsal tubercle indistinct; subarticular tubercles indistinct; tibiotarsal articulation reaching to anterior corner of eye when limb adpressed along body.

Skin: Dorsal surface smooth, with sparse small granules; dorsum with a weak discontinuous V-shaped ridge; dorsolateral ridges discontinuous; flanks with small glandular warts; supratympanic fold present, from posterior corner of eye to axilla; outer margin of upper eyelid with weak medial bumped appendage; around cloaca with small tubercles; ventral surface smooth.

Colouration in life: Dorsal surface light yellowish-brown; a dark brown triangular marking between eyes; upper lip with vertical dark bars; ventral surface white, a round white spot present on each side of the chest; ventral surface of limbs reddish (Fig. 3).



Figure 3. [doi](#)

*Xenophrys cf. parva* (adult male) from Ha Tinh Province, Vietnam. Photo by V. Q. Dau.



## Distribution

In Vietnam, this species has been recorded from Dien Bien, Lai Chau, Lao Cai, Ha Giang, Son La and Thanh Hoa Provinces (Nguyen et al. 2009, Orlov et al. 2015, Luong et al. 2019, Pham et al. 2019a and Frost 2024). Elsewhere, this species is known from China, Myanmar, Laos and Thailand (Lyu et al. 2023) (see discussion).

## Ecology

The specimen was found on a tree near a small stream at 22:00 h pm. The surrounding habitat was evergreen forest of medium hardwoods and shrubs.

## Notes

The specimen from Ha Tinh differs from the description of Lyu et al. (2023) in having relative finger lengths  $I \leq II < IV < III$  (vs.  $II < IV < I < III$ ) and dorsum with a small dark X-shaped marking (vs. larger dark X-shaped marking).

## *Xenophrys lancangica* Lyu, Wang & Wang, 2023

### Material

- a. scientificName: *Xenophrys lancangica*; scientificNameID: *Xenophrys lancangica*; genus: *Xenophrys*; specificEpithet: *lancangica*; scientificNameAuthorship: Lyu, Wang & Wang, 2023; country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 260m; verbatimLatitude: 18°20'29.72"N; verbatimLongitude: 105°14'29.27"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: DQV 0085; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: D973A0ED-6F81-5A0B-A72B-717AF63D19E0

### Description

Morphological characters of the specimen from Ha Tinh Province matched the description of Lyu et al. (2023): SVL 67.6 mm in male ( $n = 1$ ); head wider than long (HL 24.8, HW 25.1 mm, HL/SVL 36.7%, HW/SVL 37.1%); snout obliquely truncate, projecting beyond upper jaw (RL 10.1 mm), as long as horizontal diameter of eye (ED 10.0 mm); canthus rostralis well developed; loreal region concave; nostril lateral, closer to eye than to tip of snout (NS 5.1 mm, EN 4.7 mm); interorbital space flat, broader than upper eyelid and narrower than internarial distance (IOD 10.5 mm, UEW 7.1 mm, IN 4.6 mm); tympanum distinct (TD 5.2 mm); vomerine teeth present; tongue weakly notched posteriorly; vocal sac present in male. Fore-limb slender (HAL 33.2 mm); relative finger lengths  $II < IV < I < III$ ; fingers without dermal fringe, free of webbing; tips of fingers round, slightly swollen; subarticular tubercles indistinct; palmar tubercles indistinct; nuptial pads on dorsal bases of fingers I and II. Hind-limb slender, tibia longer than thigh (TbL 37.0 mm, FL 34.6 mm); relative toe lengths  $I < II < V < III < IV$ ; tips of

toes round, slightly swollen; webbing rudiment; toes dermal narrow lateral fringes; metatarsal tubercle indistinct; subarticular tubercles indistinct; inner and outer metatarsal tubercles indistinct; tibiotarsal articulation reaching to the position between nostril and tip of snout when limb adpressed along body.

Skin: Dorsal surface smooth with sparse small granules; a weak discontinuous X-shaped ridge on centre of dorsum, dorsolateral ridges present; outer margin of upper eyelid with a weak medial bumped appendage; flank with sparse large tubercles; supratympanic fold present, from posterior edge of eye to axilla; ventral surface smooth.

Colouration in life: Dorsal surface brown-yellowish, with a triangular marking between eyes, a Y-shaped marking on centre of dorsum; dorsal limb with dark transverse bands; throat and chest brown, with a round white spot on each side of the chest present; ventral surface whitish, ventral surface of limbs reddish (Fig. 4).



Figure 4. [doi](#)

*Xenophrys lancangica* (adult male) from Ha Tinh Province, Vietnam. Photo by V. Q. Dau.

### Distribution

In Vietnam, this species is known from Dien Bien, Phu Tho, Thanh Hoa, Nghe An, Quang Binh and Quang Tri Provinces (Frost 2024). Elsewhere, this species is known from China, Laos and Thailand (Frost 2024).

### Ecology

The specimen was found on the tree at 20:24 h pm. The surrounding habitat was evergreen forest of medium hardwoods and shrubs.



## Notes

*Xenophrys lancangica* closely resembles *X. maosonensis* and *X. truongsongensis*, but it differs from *X. maosonensis* by having the tibiotarsal articulation reaching to the position between nostril and tip of snout (vs. reaching to centre of eye in *X. maosonensis*) (Lyu et al. 2023) and from *X. truongsongensis* by having flank with large tubercles (vs. with small tubercles in *X. truongsongensis*); toes lateral fringes absent (vs. toes with narrow lateral fringes); and white spots on flank and back of thigh larger than those in *X. truongsongensis* (Luong et al. 2022).

## *Quasipaa* cf. *verrucospinosa* (Bourret, 1937)

### Materials

- a. scientificName: *Quasipaa* cf. *verucospinosa*; scientificNameID: *Quasipaa* cf. *verucospinosa*; genus: *Quasipaa*; specificEpithet: *verucospinosa*; scientificNameAuthorship: (Bourret, 1937); country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 240 m; verbatimLatitude: 18°17'13.3"N; verbatimLongitude: 105°22'08.1"E; verbatimCoordinateSystem: WGS84; eventDate: December; eventTime: 2023; eventRemarks: collected by A.V Pham and A. D Tran; individualCount: 1; sex: female; lifeStage: adult; catalogNumber: HT.2023.27; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: 0A905E19-8D9C-5061-96D1-92FF26553133
- b. scientificName: *Quasipaa* cf. *verucospinosa*; scientificNameID: *Quasipaa* cf. *verucospinosa*; genus: *Quasipaa*; specificEpithet: *verucospinosa*; scientificNameAuthorship: (Bourret, 1937); country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 240 m; verbatimLatitude: 18°17'13.3"N; verbatimLongitude: 105°22'08.1"E; verbatimCoordinateSystem: WGS85; eventDate: January; eventTime: 2024; eventRemarks: collected by A.V Pham and A. D Tran; individualCount: 1; sex: female; lifeStage: adult; catalogNumber: HT.2023.28; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: 50A86991-6867-504B-9D0B-7A5F02CBE588
- c. scientificName: *Quasipaa* cf. *verucospinosa*; scientificNameID: *Quasipaa* cf. *verucospinosa*; genus: *Quasipaa*; specificEpithet: *verucospinosa*; scientificNameAuthorship: (Bourret, 1937); country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 240 m; verbatimLatitude: 18°17'13.3"N; verbatimLongitude: 105°22'08.1"E; verbatimCoordinateSystem: WGS86; eventDate: February; eventTime: 2025; eventRemarks: collected by A.V Pham and A. D Tran; individualCount: 1; sex: female; lifeStage: adult; catalogNumber: HT.2023.29; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: E09D4017-003D-50E5-B90B-98C4FFB9CDE5

### Description

Morphological characters of the specimens from Ha Tinh Province matched the descriptions of Bourret (1937), Bourret (1942) and Fei et al. (2009): Body large, SVL

76.0–95.9 mm in females ( $n = 3$ ). Head wider than long (HL 30.3–37.0 mm, HW 31.9–39.9 mm; HL/SVL 38.6–40.5%; HW/SVL 41.6–43.5%); snout round, barely projecting beyond upper jaw (RL 12.2–15.0 mm), longer than horizontal diameter of eye (ED 10.6–12.9 mm); canthus rostralis indistinct; loreal region oblique, concave; nostril lateral, closer to eye than to tip of snout (NS 7.2–8.4 mm, EN 5.8–6.4 mm); interorbital space flat, narrower than upper eyelid and internarial distance (IOD 6.3–7.8 mm, UEW 7.7–69.0 mm, IN 8.0–8.8 mm); tympanum visible (TD 3.0–3.3 mm), smaller than the distance from tympanum to eye (TYE 4.7–6.5 mm), about 26–28% eye diameter; vomerine teeth present; tongue cordiform, notched posteriorly. Fore-limb short (HAL 31.2–41.8 mm); relative finger lengths  $II < I < IV < III$ ; fingers free of webbing; fingers II and III with narrow lateral dermal fringes; tips of fingers round, slightly swollen; palmar tubercles two, inner metatarsal tubercle oval, outer metatarsal tubercle elongate. Hind-limb: thigh shorter than tibia (FL 39.5–50.5 mm, TbL 45.0–54.5 mm); relative toe lengths  $I < II < V < III < IV$ ; tips of toes swollen; toes fully webbed; outer sides of toes I and V with dermal fringe; tibiotarsal articulation reaching to nostril when limb adpressed along body; inner metatarsal tubercle elongate, outer metatarsal tubercle absent.

Skin: Dorsal surface with oval, round and elongate tubercles; flank covered by oval and round tubercles; supratympanic fold present, from posterior corner of eye to angle of jaw; ventral surface smooth.

Colouration in life: Dorsal surface yellowish-brown; dorsal limbs light brown with indistinct dark crossbars; throat white with brown markings; ventral surface of limbs, chest and belly immaculate white (Fig. 5).



Figure 5. [doi](#)

*Quasipaa* cf. *verrucospinosa* (adult female) from Ha Tinh Province, Vietnam. Photo by A. V. Pham.

## Distribution

In Vietnam, this species was recorded from Dien Bien, Lao Cai, Son La, Vinh Phuc, Nghe An, Thua Thien Hue and Kon Tum Provinces (Frost 2024). Elsewhere, this species is known from Laos (Frost 2024).

## Ecology

All specimens were found near waterfalls in rocky streams between 20:00 and 22:00 h pm. The surrounding habitat was evergreen forest of large and medium hardwoods mixed with shrubs.

## Notes

Our specimens from Ha Tinh Province slightly differ from those in descriptions of Bourret (1942) and Inger et al. (1999) by having inner metatarsal tubercle oval (vs. inner metatarsal tubercle round); different dorsal pattern (yellowish-brown vs. yellowish-grey); different ventral colour pattern (immaculate white vs. light yellow); and iris dark green (vs. pale copper).

## *Amolops compotrix* (Bain, Stuart & Orlov, 2006)

### Materials

- a. scientificName: *Amolops compotrix*; scientificNameID: *Amolops compotrix*; class: Amphibia; order: Anura; family: Ranidae; genus: *Amolops*; specificEpithet: *compotrix*; scientificNameAuthorship: (Bain, Stuart, and Orlov, 2006); country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 1330 m; verbatimLatitude: 18°14'10.68"N; verbatimLongitude: 105°25'54.05"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: female; lifeStage: adult; catalogNumber: HDU 03109; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: CB80C67B-FEA6-560C-81C2-E7ECB9D4B9A2
- b. scientificName: *Amolops compotrix*; scientificNameID: *Amolops compotrix*; class: Amphibia; order: Anura; family: Ranidae; genus: *Amolops*; specificEpithet: *compotrix*; scientificNameAuthorship: (Bain, Stuart, and Orlov, 2006); country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 1330 m; verbatimLatitude: 18°14'10.68"N; verbatimLongitude: 105°25'54.05"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: HDU 03110; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: F3F59600-071E-5026-8033-B9C07BEF908F
- c. scientificName: *Amolops compotrix*; scientificNameID: *Amolops compotrix*; class: Amphibia; order: Anura; family: Ranidae; genus: *Amolops*; specificEpithet: *compotrix*; scientificNameAuthorship: (Bain, Stuart, and Orlov, 2006); country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 1021 m; verbatimLatitude:

18°17'41.03"N; verbatimLongitude: 105°17'34.30"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: HDU 03149; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: 3A2A6808-B947-5573-8445-D79FA028D20C

## Description

Morphological characters of the specimens from Ha Tinh Province matched the description of Bain et al. (2006): Body elongate, SVL 40.6–42.0 mm in males ( $n = 2$ ) and 60.2 mm in the female ( $n = 1$ ). Head longer than wide (HL 15.4–17.3 mm, HW 13.4–15.0 mm in males; HL 22.0 mm, HW 18.6 mm in the female); snout obtusely pointed in dorsal view, projecting beyond lower jaw (RL 6.3–6.8 mm in males and 9.4 mm in the female), longer than horizontal diameter of eye (ED 6.0–6.5 mm in males and 8.4 mm in the female); canthus rostralis distinct, slightly constricted behind nostrils; lores concave and oblique; nostril lateral, midway between tip of snout and eye (NS 2.7–2.8 mm, EN 2.8–3.0 mm in males and NS 4.3 mm, EN 4.1 mm in the female); interorbital distance narrower than upper eyelid and internarial distance (IOD 3.7–3.8 mm, UEW 4.6–4.8 mm, IN 3.9–4.4 mm in males and IOD 4.5 mm, UEW 5.8 mm, IN 5.2 mm in the female); pineal body distinct; tympanum distinct, round, 43% of eye diameter in males or 40% of eye diameter in the female; vomerine teeth present; tongue cordiform, deeply notched posteriorly; vocal openings on floor of mouth at corner. Fore-limb long (HAL 21.3–23.7 mm in males, 30.8 mm in the female); relative finger lengths  $I < II < IV < III$ ; fingers without dermal fringe, free of webbing; tips of fingers expanded with circum-marginal grooves; palmar tubercles two, oval, in contact; nuptial pads present. Hind-limb slender, thigh shorter than tibia (FL 24.3–25.3 mm, TbL 24.7–27.4 mm in males and FL 35.5 mm, TbL 37.9 in the female); relative toe lengths  $I < II < V < III < IV$ ; tips of toes expanded, width of toe IV disc equal to that of finger III disc; webbing formula  $I0-II0-0III0-1IV1-0V$ ; tibiotarsal articulation reaching to beyond tip of snout when limb adpressed along body; inner metatarsal tubercle distinct, elongate, oval; outer metatarsal tubercle round, small.

Skin: Dorsal surface smooth; supratympanic fold present, from posterior edge of eye to axilla; glandular dorsolateral fold present; flank smooth; ventral surface smooth.

Colouration in life: Dorsal surface green, with dark brown speckling diurnally, lateral side of body dark brown below edge of dorsolateral fold; lip stripe white, from tip of snout to posterior of arm insertion; dorsal surface of limbs greyish-brown with dark crossbars; ventral surface of jaw, throat, chest and belly white; ventral surface of limbs cream (Fig. 6).

## Distribution

In Vietnam, this species was recorded from Nghe An, Quang Binh, Thua Tien Hue and Kon Tum Provinces (Pham et al. 2019b, Dau et al. 2020 and Frost 2024). Elsewhere, this species is known from Laos (Frost 2024).



Figure 6. [doi](#)

*Amolops compotrix* (adult male) from Ha Tinh Province, Vietnam. Photo by V. Q. Dau.

### Ecology

All specimens were found on a leaf or tree branches near small streams between 19:30 and 21:00 h pm. The surrounding habitat was evergreen forest of large hardwoods and shrubs.

### *Kurixalus odontotarsus* (Ye & Fei, 1993)

#### Materials

- a. scientificName: *Kurixalus odontotarsus*; scientificNameID: *Kurixalus odontotarsus*; class: Amphibia; order: Anura; family: Rhacophoridae; genus: *Kurixalus*; specificEpithet: *odontotarsus*; scientificNameAuthorship: Nguyen, Duong, Luu, and Poyarkov, 2020; country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 631 m; verbatimLatitude: 18°15'41.09"N; verbatimLongitude: 105°26'38.14"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: HDU 03120; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: DFCD61C5-110D-5E34-AC35-E6109A63B719
- b. scientificName: *Kurixalus odontotarsus*; scientificNameID: *Kurixalus odontotarsus*; class: Amphibia; order: Anura; family: Rhacophoridae; genus: *Kurixalus*; specificEpithet: *odontotarsus*; scientificNameAuthorship: Nguyen, Duong, Luu, and Poyarkov, 2020; country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 290 m; verbatimLatitude: 18°20'43.04"N; verbatimLongitude: 105°17'46.03"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: female; lifeStage: adult;

- catalogNumber: HDU 03144; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: 57636F02-A6E9-505F-99EF-9F9495AD607C
- c. scientificName: *Kurixalus odontotarsus*; scientificNameID: *Kurixalus odontotarsus*; class: Amphibia; order: Anura; family: Rhacophoridae; genus: *Kurixalus*; specificEpithet: *odontotarsus*; scientificNameAuthorship: Nguyen, Duong, Luu, and Poyarkov, 2020; country: Vietnam; countryCode: VN; stateProvince: Ha Tinh; county: Ha Tinh; municipality: Vu Quang; locality: near Huong Quang Commune; verbatimElevation: 290 m; verbatimLatitude: 18°20'43.04"N; verbatimLongitude: 105°17'46.03"E; verbatimCoordinateSystem: WGS84; eventDate: June; eventTime: 2013; eventRemarks: collected by V. Q. Dau and T. C. Thai; individualCount: 1; sex: male; lifeStage: adult; catalogNumber: HDU 03145; language: en; collectionCode: Amphibians; basisOfRecord: PreservedSpecimen; occurrenceID: 5E9B7883-0BE4-55E4-8F5A-CDD022177708

## Description

Morphological characters of specimens from Ha Tinh Province matched the descriptions of Fei et al. (2009) and Wu et al. (2022): SVL 28.7–33.2 mm in males ( $n = 2$ ) and 47.1 mm in the female ( $n = 1$ ). Head wider than long (HL 10.8–11.4 mm, HW 11.0–12.0 mm, HL/SVL 34.3–37.5%, HW/SVL 36.0–38.2% in males; HL 17.1 mm, HW 17.4 mm, HL/SVL 36.2%, HW/SVL 37.0% in the female); snout slightly pointed, projecting beyond upper jaw (RL 4.7–5.3 mm in males and 6.7 mm in the female), longer than horizontal diameter of eye (ED 4.0–4.1 mm in males and 4.9 mm in the female); canthus blunt; loreal region oblique, concave; nostril lateral, closer to eye than to tip of snout (NS 1.7–2.2 mm, EN 2.1–2.5 mm in males and NS 2.2 mm, EN 3.5 mm in the female); interorbital space flat, broader than upper eyelid and internarial distance (IOD 3.2–3.7 mm, UEW 3.0–3.5 mm, IN 2.4–2.8 mm in males and IOD 4.4 mm, UEW 4.0 mm, IN 3.6 mm in the female); tympanum distinct, round (TD 2.0–2.3 mm in males, 3.2 mm in the female); vomerine teeth present; tongue heart-shaped, notched posteriorly; vocal single internal. Fore-limb long (HAL 14.2–16.1 mm in males, 23.3 mm in the female); relative finger lengths  $I < II < IV < III$ ; fingers with lateral dermal fringes, with basal webbing; tips of finger expanded into wide discs with circum-marginal and transverse ventral grooves; palmar tubercles two; finger I of two males with nuptial pads. Hind-limb slender, thigh shorter than tibia (FL 15.1–16.5 mm, TbL 15.6–16.7 mm in males and FL 22.5 mm, TbL 22.6 in the female); relative toe lengths  $I < II < V < III < IV$ ; tips of toes expanded into discs, webbing formula  $I2-2\frac{1}{2}II1\frac{1}{2}-3III1\frac{1}{2}-3\frac{1}{2}IV3-1\frac{1}{2}V$ ; inner metatarsal tubercle distinct, outer metatarsal tubercle absent; tibiotarsal articulation reaching to the position between the eye and snout when limb adpressed along body.

Skin: Dorsal surface with small tubercles; supratympanic fold present, from posterior edge of eye to axilla; along outer edge of forearm with a row of warts, forming a serrated fringe; along outer edge of tarsus and metatarsus with a series of tubercles forming serrated dermal fringe; below vent with a patch of white pustules; ventral surfaces granulated.

Colouration in life: Dorsal surface yellowish-brown or greenish-brown, with a dark-brown interorbital bar in triangle shape and a dark-brown Y-shaped saddle-like marking



on the back; flank light brown with indistinct brownish spots; dorsal limbs brown with dark crossbars; a white conical projection on tibiotarsal articulation; ventral surface with dark-grey mottling, becoming denser on throat, more sparse on chest and belly (Figs 7, 8).



Figure 7. [doi](#)

*Kurixalus odontotarsus* (adult male) from Ha Tinh Province, Vietnam. Photo by V. Q. Dau.



Figure 8. [doi](#)

*Kurixalus odontotarsus* (adult female) from Ha Tinh Province, Vietnam. Photo by V. Q. Dau.

### Distribution

In Vietnam, this species was recorded from Lao Cai and Ha Giang Provinces (Frost 2024). Elsewhere, this species is known from China, Laos and Thailand (Frost 2024).

## Ecology

All specimens were found on leaves near large streams between 21:00 and 22:00 h. The surrounding habitat was evergreen forest of large hardwoods and shrubs.

## Discussion

Our new records bring the total number of amphibian species from Ha Tinh Province to 42, comprising one species of Ichthyophiidae, two species of Bufonidae, seven species of Megophryidae, six species of Microhylidae, seven species of Dicroglossidae, ten species of Ranidae and nine species of Rhacophoridae. Most of records of the amphibian fauna in Ha Tinh Province are known from Ke Go Nature Reserve (30 species) and Vu Quang NP (29 species), of which 22 species occur in both Ho Ke Go Nature Reserve and Vu Quang NP (Suppl. material 1). In terms of conservation status, Vu Quang harbours four species that are listed in the Red Data Book of Vietnam (Dang et al. 2007), with three species categorised as Endangered - EN (*Rhacophorus kio*, *Theloderma corticale*, *Zhangixalus feae*) and one species as Vulnerable - VU (*Ingerophrynus galeatus*); one species (*Microhyla annamensis*) is listed as {Vulnerable} in the IUCN Red List (IUCN 2024).

According to Mahony et al. (2020), *Xenophrys parva* is only known with certainty from eastern Myanmar and the old range statement for nominal *Megophrys parva* (western to eastern Nepal and Bhutan, India through Bangladesh and Myanmar to western Thailand); and records of this species in China (Xizang, Yunnan, and Guangxi), northern Vietnam, and northern Laos can apply to other named and unnamed species. However, Lyu et al. (2023) provided a detailed account of *Xenophrys parva*, placing it in the *Xenophrys lekagulii* group, as well as providing an account and specifically including genetically confirmed records of the species from Chiang Mai (Thailand), Laos and north-western Vietnam.

## Acknowledgements

We are grateful to the directorates of Vu Quang ASEAN Heritage Park for support of our fieldwork and issuing relevant permits. This research has been done under the research project QG.23.18 of Vietnam National University, Hanoi.

## References

- Bain R, Stuart B, Orlov N (2006) Three new Indochinese species of cascade frogs (Amphibia: Ranidae) allied to *Rana archotaphus*. Copeia 2006 (1): 43-59. [https://doi.org/10.1643/0045-8511\(2006\)006\[0043:tnisoc\]2.0.co;2](https://doi.org/10.1643/0045-8511(2006)006[0043:tnisoc]2.0.co;2)
- Boulenger GA (1893) Concluding report on the reptiles and batrachians obtained in Burma by Signor L. Fea dealing with the collection made in Pegu and the Karin Hills in 1887-88. Annali del Museo Civico di Storia Naturale di Genova 2 (13): 304-347.

- Bourret R (1937) Notes herpétologiques sur l'Indochine française. XIV. Les batraciens de la collection du Laboratoire des Sciences Naturelles de l'Université. Descriptions de quinze espèces ou variétés nouvelles. Annexe au Bulletin Général de l'Instruction Publique. Hanoi 1937: 5-56.
- Bourret R (1942) Les Batraciens de l'Indochine, Hanoi. Institut Océanographique de l'Indochine 529.
- Dang NT, Tran K, Dang HH, Nguyen C, Nguyen TN, Nguyen HY, Dang TD (2007) Vietnam Red Data Book. Part I. Animals. Natural Science and Technology Publishing House, Hanoi
- Dau QV, Ong VA, Thai CT, Nguyen KT (2015) First records of *Theloderma corticale* (Boulenger, 1903) in Nghe An and Ha Tinh provinces. Proceedings of the 6th National Scientific Conference on Ecology and Biological Resources Ha Noi, Vietnam 34: 405-408.
- Dau QV, Thieu TH, Le DP, Phung MT, Ong VA (2020) First records two species of Megophryidae Bonaparte, 1850 from North Central Vietnam. Vinh University Scientific Journal 49 (1A): 71-77.
- Do VT, Nguyen QT, Cao TT, Luu TK (2017) New records of two species of the family Megophryidae from Nghe An Province. Proceedings of the 7th National Scientific Conference on Ecology and Biological Resources, Ha Noi Vietnam 413-418.
- Fei L, Hu Sq, Ye Cy, Huang YZ (2009) Fauna Sinica. Amphibia 2: 957.
- Frost D (2024) Amphibian species of the world: An Online Reference, ver. 6.0, American Museum of Natural History, New York. <http://research.amnh.org/herpetology/amphibia/index.php>. Accessed on: 2024-3-03.
- Glaw F, Vences M (2007) A field guide to the amphibians and reptiles of Madagascar. Third Edition, Frosch Verlag, Cologne
- Hoang CV, Jiang Jp, Nguyen HQ, Orlov NL, Le MV, Nguyen LH, Nguyen VH, Nguyen TV, Nguyen TT, Ziegler T (2021) A new species of Vietnamophryne from Vietnam. Revue Suisse de Zoologie 128: 207-219. <https://doi.org/10.35929/RSZ.0046>
- Inger RF, Orlov NL, Darevsky I (1999) Frogs of Vietnam: a report on new collections. Fieldiana Zoology 92: 1-46.
- IUCN (2024) The IUCN Red List of Threatened Species. Version 2023.1. <http://www.iucnredlist.org>. Accessed on: 2024-3-03.
- Luong AM, Nguyen HQ, Le DT, Nguyen SL, Nguyen TQ (2019) New records of amphibians (Anura: Megophryidae, Ranidae) from Dien Bien Province, Vietnam. Herpetology Notes 12: 375-387. <https://doi.org/10.11646/zootaxa.5150.3.2>
- Luong MA, Hoang VC, Pham TC, Nguyen TT, Orlov NL, Ziegler T, Nguyen QT (2022) A new species of *Xenophrys* (Amphibia: Anura: Megophryidae) from Truong Son Range Vietnam. Zootaxa 5150: 333-356. <https://doi.org/10.11646/zootaxa.5150.3.2>
- Lyu ZT, Wang SQ, Zhang SY, Zhao J, Zeng ZC, Wan H, Yang JH, Mo YM, Wang YY (2023) Generic classification of Asian horned toads (Anura: Megophryidae: Megophryinae) and monograph of Chinese species. Zoological Research. Kunming 44: 380-450. <https://doi.org/10.24272/j.issn.2095-8137.2022.372>
- Mahony S, Kamei RG, Teeling EC, Biju SD (2020) Taxonomic review of the Asian horned frogs (Amphibia: *Megophrys* Kuhl & Van Hasselt) of Northeast India and Bangladesh previously misidentified as *M. parva* (Boulenger), with descriptions of three new species. Journal of Natural History. London 54: 119-194. <https://doi.org/10.1080/00222933.2020.1736679>

- Matsui M (2013) A new *Leptobrachium* (Vibrissaphora) from Laos (Anura: Megophryidae). *Current Herpetology*. Kyoto 32: 182-189. <https://doi.org/10.5358/hsj.32.182>
- Nguyen SV, Ho CT, Nguyen TQ (2009) Herpetofauna of Vietnam. Edition Chimaira
- Orlov NL, Poyarkov NA, Nguyen TT (2015) Taxonomic notes on *Megophrys* frogs (Megophryidae: Anura) of Vietnam, with description of a new species. *Russian Journal of Herpetology* 22 (3): 206-218.
- Pham AV, Pham TC, Doan LD, Ziegler T, Nguyen QT (2019a) New records of megophryids (Amphibia: Anura: Megophryidae) from Son La Province. Vietnam. *Biodiversity Data Journal* 7 (e39140): 1-18. <https://doi.org/10.3897/BDJ.7.e39140>
- Pham AV, Sung NB, Pham CT, Le MD, Ziegler T, Nguyen QT (2019b) A new species of *Amolops* (Anura: Ranidae) from Vietnam. *Raffles Bulletin of Zoology* 67: 363-377. <https://doi.org/10.26107/RBZ-2019-0027>
- Pham VA, Le TD, Pham TC, Nguyen LH, Ziegler T, Nguyen QT (2016) Two additional records of megophryid species, *Leptolalax minimus* (Taylor, 1962) and *Leptobrachium masatakasatoi* Matsui, 2013, for the herpetofauna of Vietnam. *Revue suisse de Zoologie* 123 (1): 35-43.
- Sheridan J, Phimmachak S, Sivongxay N, Stuart B (2023) Systematics of the Lao torrent frog, *Amolops cremnobatus* Inger & Kottelat, 1998 (Anura: Ranidae), with descriptions of four new species. *Vertebrate Zoology Senckenberg* (73): 931-956. <https://doi.org/10.3897/vz.73.e102475>
- Simmons JE (2002) Herpetological collecting and collections management. Revised edition. Society for the Study of Amphibians and Reptiles. *Herpetological Circular* 31: 1-153.
- Thai CT (2017) Diversity of the family Rhacophoridae (Amphibia: Anura) in Vu Quang National Park, Vietnam. Master thesis of Vinh University
- Tordoff AW, Tran BQ, Nguyen TD, Le HM (2004) Sourcebook of existing and proposed protected areas in Vietnam. Birdlife International in Indochina and Ministry of Agriculture and Rural Development. 2. Hanoi
- Wu Y, Suwannapoom C, Poyarkov N, Gao W, Karuno A, Yuan Z (2022) First record of *Kurixalus odontotarsus* (Ye et Fei, 1993) and *Raorchestes longchuanensis* (Yang et Li, 1978) (Anura: Rhacophoridae) in Thailand. *Russian Journal of Herpetology* 29: 1-18. <https://doi.org/10.30906/1026-2296-2022-29-1-1-18>
- Ziegler T, Ong VA, Hoang XQ, Pham TC, Nguyen QT (2022) A taxonomic update 20 years after the book release "the amphibians and reptiles of a lowland forest reserve In vietnam"- Ho Ke Go: implications for conservatio. *Academia Journal of Biology* 44 (3): 111-132. <https://doi.org/10.15625/2615-9023/17102>

## Supplementary material

### Suppl. material 1: List of amphibians species

**Authors:** Vinh Quang Dau, Cuong The Pham, Truong Quang Nguyen, Toan Canh Thai, Anh Dinh Tran, Anh Van Pham

**Data type:** Table

**Brief description:** List of amphibians species recorded in Vu Quang NP and Ha Tinh Province, Vietnam.

[Download file](#) (74.50 kb)