



Research Article

Reference DNA barcodes and other mitochondrial markers for identifying Caribbean Octocorals

Jaime G. Morín[‡], Dagoberto E. Venera-Pontón[§], Amy C. Driskell[|], Juan A. Sánchez[¶], Howard R. Lasker[#], Rachel Collin[□]

[‡] Laboratorio de Sistemática Molecular y Filogeografía, Facultad de Ciencias Biológicas, Universidad Nacional Mayor de San Marcos, Lima, Peru

[§] Smithsonian Tropical Research Institute, Panama City, Panama

[|] Laboratories of Analytical Biology, National Museum of Natural History, Smithsonian Institution, Washington, D.C., United States of America

[¶] Laboratorio de Biología Molecular Marina – BIOMMAR, Bogotá, Colombia

[#] Department of Geology, University at Buffalo, Buffalo, United States of America

[□] Smithsonian Tropical Research Institute, Balboa, Panama

Corresponding author: Rachel Collin (collinr@si.edu)

Academic editor: Danwei Huang

Received: 31 Oct 2018 | Accepted: 04 Feb 2019 | Published: 20 Feb 2019

Citation: Morín J, Venera-Pontón D, Driskell A, Sánchez J, Lasker H, Collin R (2019) Reference DNA barcodes and other mitochondrial markers for identifying Caribbean Octocorals. Biodiversity Data Journal 7: e30970. <https://doi.org/10.3897/BDJ.7.e30970>

Abstract

DNA barcoding is a useful tool for documenting the diversity of metazoans. The most commonly used barcode markers, 16S and COI, are not considered suitable for species identification within some "basal" phyla of metazoans. Nevertheless metabarcoding studies of bulk mixed samples commonly use these markers and may obtain sequences for "basal" phyla. We sequenced mitochondrial DNA fragments of cytochrome oxidase c subunit I (COI), 16S ribosomal RNA (16S), NADH dehydrogenase subunits 2 (16S-ND2), 6 (ND6-ND3) and 4L (ND4L-MSH) for 27 species of Caribbean octocorals to create a reference barcode dataset and to compare the utility of COI and 16S to other markers more typically used for octocorals. The most common genera (*Erythropodium*, *Ellisella*, *Briareum*, *Plexaurella*, *Muriceopsis* and *Pterogorgia*) were effectively distinguished by small differences (5 or more substitutions or indels) in COI and 16S sequences. *Gorgonia* and *Antillogorgia* were effectively distinguished from each other by unique haplotypes, but the small genetic differences make distance approaches ineffective for these taxa. *Plexaura*,

Pseudoplexaura and *Eunicea* were indistinguishable from each other but were generally effectively distinguished from other genera, further supporting the idea that these genera have undergone a rapid endemic radiation in the Caribbean.

Keywords

cytochrome oxidase I, 16S, Panama, Bocas del Toro, gorgonian, DNA barcoding

Introduction

DNA barcoding is a useful tool for documenting the diversity of most metazoan groups (Bucklin et al. 2011). Short, easily amplifiable fragments that vary amongst closely related species are sequenced from specimens identified by experts and then used as a tool to identify sequences from unidentified samples (Hebert and Gregory 2005, Bucklin et al. 2011, Hajibabaei et al. 2007). This approach is useful in a variety of contexts (Bucklin et al. 2011, Bucklin et al. 2016, Fonseca et al. 2010, Hajibabaei et al. 2007); however the most commonly used barcode marker, a fragment of cytochrome c oxidase subunit I (COI hereafter), is not suitable for species identification within some metazoan groups such as Porifera and Anthozoa (Shearer et al. 2002, Huang et al. 2008). Consequently, less effort has been invested in building reference databases of COI and 16S barcodes within these taxa, as opposed to those groups where these markers are perceived as useful for species identification.

DNA barcoding has been used extensively to detect cryptic diversity within clades, with specific studies generally focusing on diversity within a single family, class or order. When COI is problematic, alternate barcodes are used, for example, 16S is commonly used as the molecular barcode in hydrozoans (Moura et al. 2011, Zheng et al. 2014). For some clades, significant effort has been spent on identifying reliable, divergent markers and building extensive databases of these sequences for identification purposes (e.g. van der Ham et al. 2009, McFadden et al. 2014).

Recent assessments of biodiversity are increasingly focusing on metabarcoding analyses of bulk mixed samples, such as gut contents (Leray et al. 2013, Oliverio et al. 2009), settlement plates (Zaiko et al. 2016), plankton samples (Bucklin et al. 2016), sediment samples (Fonseca et al. 2010) or environmental DNA from water samples (Stat et al. 2017, Thomsen et al. 2012). These samples are typically analysed using markers that amplify across the most diverse set of taxa; most commonly COI and sometimes 16S for studies of species diversity or the nuclear SSU 18S gene in studies aimed at higher taxonomic levels. Due to the non-selective nature of metabarcoding, this approach generates sequences for taxa where the marker is not useful for species-level identification as well as for those where it is useful. Therefore, a reference set of COI and 16S sequences for these taxa could at least provide the lowest possible identification when these problematic taxa occur in mixed samples.

Most Caribbean gorgonian octocorals are endemic and closely related (Sánchez et al. 2003, Sánchez 2016, Wirshing et al. 2005). This results in many taxa with morphologically similar sister species making it extremely difficult to differentiate a number of the species with morphological characters (Sánchez and Wishing 2005). In addition, octocorals are notoriously difficult to distinguish with molecular data (France and Hoover 2002, McFadden et al. 2010, McFadden et al. 2014, McFadden et al. 2004, van der Ham et al. 2009) since their mitochondria have a unique DNA repair mechanism that is thought to be responsible for the slow evolution of mitochondrial genes (Bilewitch and Degnan 2011). Nevertheless, gorgonians are an abundant and diverse part of the macro-fauna of Caribbean reefs, with as many as 40 species occurring on a single Caribbean reef (Lasker and Coffroth 1983, Sánchez et al. 1998, Sánchez et al. 1997, Etnoyer et al. 2010, Tsounis et al. 2018). In some places, they contribute significantly to three dimensional reef structures and provide microhabitats for macro-invertebrates and fishes (Sánchez 2016). Furthermore, octocoral abundances have increased at some locations in contrast to the overall decline in scleractinians (Edmunds and Lasker 2016, Ruzicka et al. 2013). Therefore, tools that facilitate their inclusion in biodiversity assays would be a useful addition to researchers' toolkits.

Here we had two main objectives: (1) to generate a reference set of DNA barcode sequences for common octocorals from the Southern Caribbean and (2) determine how the 16S and COI barcode fragments compare with other mitochondrial markers in their ability to distinguish genera and, in some cases, species in the fauna of Bocas del Toro, Panama.

Materials and Methods

A total of 180 octocoral tissue samples were collected by SCUBA-diving during the summer of 2007 from the shallow waters around the Bocas del Toro Archipelago on the Caribbean coast of Panama. Tissue samples consisted of 10-20 cm sections clipped from a distal branch for each colony. Samples were identified to species using a combination of visual identification in the field (colony morphology) and microscopic examination of spicule preparations. In some cases, individuals were intermediate in morphology or could not be identified to species and were only identified to genus. Dry tissue vouchers were deposited at the USNM and the University of Panama. Details about individual samples are provided in the BoLD project workbench (Ratnasingham and Herbert 2007).

Genomic DNA was extracted from small samples of tissue (0.5 cm³) from each specimen using the mouse tail kit on a Biosprint 96 (Qiagen). The resuspension volume was 200 µl. We used PCR to amplify fragments of the mitochondrial genes COI, 16S, ND6-ND3, 16S-ND2 and ND4L-MSH with the primers listed in Table 1. When the barcode ("Folmer") fragment of COI could not be amplified with the primers derived from Folmer et al. 1994, we amplified a larger fragment that includes part of the COII gene. We then extracted the barcode sequence from this longer sequence. When neither of these amplifications was successful, we amplified a shorter COI fragment with gorgonian-specific primers. The 10 µl PCR cocktail included 0.5 µl 50 mM MgCl₂, 0.1 µl 20 µg/µl BSA, 0.5 µl dNTPs (2.5 mM

each), 0.3 µl each 10 mM primer and Biolase taq polymerase (Bioline). Annealing temperatures were 46°C for the barcode fragment of COI and ND4L-MSH, 48°C for 16S and the gorgonian-specific COI fragment and 52°C for ND6-ND3, 16S-ND2 and the long fragment of COI. Amplified PCR products were cleaned using the ExoSAP-IT protocol (Affymetrix) and sequenced using the Big Dye protocol (ThermoFisher).

Table 1.

Primers used for amplification and sequencing of loci in this study and size of amplified fragments. *Lengths of fragments that include non-coding regions as well as ribosomal sequences can vary. ¹McFadden et al. 2004; ²France and Hoover 2001; ³Meyer 2003; ⁴this study; ⁵Palumbi et al. 1991; ⁶Brugler and France 2008; ⁷Sánchez et al. 2003.

Locus	Primers	Expected Size*
COI long	<i>COII-8068F</i> ¹ CCATAACAGGACTAGCAGCATC <i>COIOCTR</i> ² ATCATAGCATAGACCATAACC	940
COI Folmer	<i>dgLCO1490</i> ³ GGTC AACAAATCATAAAGAYATYGG <i>dgHCO2198</i> ³ TAAACTTCAGGGTGACCAAARAYCA	655
COI short	<i>GorgM13_F</i> ⁴ CACGACGTTGTAAAACGACGTATGTTAGGAGATGATCATCTATAT <i>GorgM13_R</i> ⁴ GGATAACAATTTACACAGGGAATGTTGTATTAATAATTYCTRTCTGT	468
16S	<i>16Sar</i> ⁵ CGCCTGTTTATCAAAAACAT <i>16Sbr</i> ⁵ CCGGTCTGAACTCAGATCACGT	668
ND6-ND3	<i>ND6-1487F</i> ¹ TTTGGTTAGTTATTGCCTTT <i>ND3-2126R</i> ¹ CACATTCATAGACCGACACTT	611
16S-ND2	<i>16S-647F</i> ¹ TTTGGTTAGTTATTGCCTTT <i>ND2-1418R</i> ¹ ACATCGGGAGCCACATA	758
ND4L-MSH	<i>ND4L-2475F</i> ⁶ TAGTTTTACTGGCCTCTAC <i>MUT-3458R</i> ⁷ TSGAGCAAAGCCACTCC	870

Sequences were screened for quality and contigs of forward and reverse sequences were produced using Sequencher 5.4.6 (Gene Codes). Only sequences with a Phred quality score of at least 30 for more than 85% of the bases were combined into contigs and used for analysis. Sequences were compared internally across our dataset and BLASTned against GenBank sequences to check for contamination or mislabelling. After this step, through an abundance of caution, all 27 ND4L-MSH sequences from a single plate were excluded from the analyses because a subset of them showed unexplained divergences from previously published GenBank sequences. No other plate of sequences in our analysis showed similar problems. Sequences of each marker were aligned with ClustalX (gap-opening penalty: 15, gap-extension penalty: 6.66, transition weight: 0.5, delay divergent cutoff: 30%) and used to generate a matrix of pairwise differences [including the number of substitutions and indels (insertions/deletions)] between all the sequenced specimens. This matrix was then used to build heatmaps displaying the average pairwise differences between species. The patterns observed in heatmaps were contrasted with those observed in maximum parsimony and neighbour-joining trees (BIONJ, Gascuel 1997) built with the same matrix of pairwise differences.

Most markers showed two groups within which the pairwise differences amongst species (and genera) were consistently small. For these groups, we constructed haplotype networks for all the markers using Haplotype Viewer (Center for Integrative Bioinformatics Vienna; <http://www.cibiv.at/>) to determine if haplotypes were unique or related to particular species or genera.

Data Resources

The data underpinning the analysis reported in this paper are deposited in the Barcode of Life Database (dx.doi.org/10.5883/DS-OCTOCORA) (Ratnasingham and Herbert 2007) and the barcode markers have the following GenBank numbers COI:MK153303-MK153482 and 16S:MK153483-MK153626.

Results and Discussion

We were able to collect and successfully sequence 28 species represented by 180 individuals of octocorals from Bocas del Toro (Table 2). This represents 45% of the 61 species and includes 12 of the sixteen genera and all seven families reported for the region (Sánchez and Wishing 2005). Most markers were sequenced successfully for most species but 16S-ND2 could not be amplified for the genus *Erythropodium*, whereas the ND4L-MSH fragment failed to amplify in 5 species. Likewise, samples of *Pterogorgia anceps* almost entirely failed to amplify. Since all the specimens were processed with the same methods, differences in sequencing success may be due to taxon-specific secondary metabolites interfering with PCR.

Table 2.

Number of individuals collected and successfully sequenced for each genetic marker for samples identified to species. The colour of the cells indicates the availability of sequences for the same marker and species in GenBank. Dark grey: A new contribution; no available sequences overlap our fragment by ≥ 100 bp. Light grey: Partial sequence is already available; our sequences overlap existing sequences by >350 bp; our data extends this by >50 bp. White: Sequence already available; sequences available in GenBank overlap ours completely with <50 bp additional data. X mark indicates species that failed to amplify or resulted in unusable or suspicious sequences. *Sequences were also generated for an additional 4 *Antillologorgia*, 3 *Erythropodium*, 44 *Eunicea*, 3 *Muricea*, 1 *Plexaura*, 5 *Plexaurella*, 16 *Pseudoplexaura* and 9 *Pterogorgia* that were not identified to species.

Species	Individuals collected*	COI	16S	16S-ND2	ND6-ND3	ND4L-MSH
<i>Antillologorgia acerosa</i>	4	3	3	3	3	3
<i>Antillologorgia americana</i>	4	4	4	4	4	3
<i>Antillologorgia bipinnata</i>	8	5	5	5	5	4
<i>Antillologorgia rigida</i>	2	1	1	1	1	X

Species	Individuals collected*	COI	16S	16S-ND2	ND6-ND3	ND4L-MSH
<i>Briareum asbestinum</i>	9	8	7	7	3	7
<i>Ellisella schmitti</i>	2	2	2	1	2	2
<i>Erythropodium caribaeorum</i>	3	3	1	X	3	X
<i>Eunicea calyculata</i>	3	3	2	3	3	3
<i>Eunicea clavigera</i>	1	1	1	1	1	1
<i>Eunicea flexuosa</i>	9	8	8	8	8	8
<i>Eunicea fusca</i>	5	4	4	4	4	4
<i>Eunicea laxispica</i>	1	1	1	1	1	1
<i>Eunicea succinea</i>	1	1	1	1	1	X
<i>Eunicea tourneforti</i>	6	6	4	6	6	3
<i>Gorgonia mariae</i>	8	8	8	8	8	X
<i>Gorgonia ventalina</i>	4	4	4	4	4	X
<i>Muricea laxa</i>	7	7	3	6	7	6
<i>Muricea muricata</i>	3	3	3	3	3	3
<i>Muricea pinnata</i>	5	5	3	5	5	5
<i>Muriceopsis bayeriana</i>	6	6	6	6	6	X
<i>Muriceopsis flavida</i>	4	3	4	3	3	X
<i>Plexaura homomalla</i>	11	8	6	8	8	7
<i>Plexaura kuna</i>	4	4	4	4	4	2
<i>Plexaurella dichotoma</i>	3	3	3	3	3	3
<i>Plexaurella nutans</i>	4	2	2	2	2	2
<i>Pseudoplexaura porosa</i>	2	2	1	2	2	2
<i>Pseudoplexaura wagnaari</i>	1	1	X	1	X	1
<i>Pterogorgia anceps</i>	4	X	X	1	X	X
<i>Pterogorgia citrina</i>	2	2	2	2	2	X

New additions to GenBank from this study include COI for 19 species, 16S for 20 species, 16S-ND2 and ND4L-MSH for 3 species and 16S-ND2 and ND6-ND3 for 13 species (Table 2). Of those markers that were already in GenBank for a particular species, >40 of our sequences extended significantly beyond the previously reported fragment. Our samples also increased the geographic coverage of these species. For example, the majority of 16S-ND2 and ND4L-MSH sequences already in GenBank were collected from the Bahamas (Sánchez et al. 2003), so the addition of sequences from Panama provides new references for other areas of the species ranges.

As expected for octocorals, none of our 5 mitochondrial markers was suitable for distinguishing amongst congeneric species across the entire dataset, but they were generally useful to distinguish amongst most genera except those in the *Plexaura*-

Pseudoplexaura-Eunicea group and the *Antillologorgia-Gorgonia* group (Figs 1, 2, 3, 4Fig. 5). The studied markers were able to distinguish species, but their patterns of pairwise differences varied amongst the different markers. COI, 16S-ND2 and ND4L-MSH displayed higher divergences overall than did the other two markers. The genera *Briareum*, *Plexaurella*, *Erythropodium*, *Muriceopsis* and *Pterogorgia* were clearly distinguishable from each other. Other taxa were partially distinguishable. For example, species of *Plexaura*, *Pseudoplexaura* and *Eunicea* were often indistinguishable from each other but could generally be distinguished from species in other genera across the dataset. The same situation occurred with *Antillologorgia* and *Gorgonia* which had very similar haplotypes. *Ellisella* and *Erythropodium* were clearly distinguishable with COI, 16S and 16S-ND2 but surprisingly appeared to fall within the *Plexaura-Pseudoplexaura-Eunicea* clade with ND4L-MSH (*Ellisella*) and ND6-ND3 (both). We could find no problems with our sequences or data flow which would account for this, but this result should be independently confirmed for ND4L-MSH and ND6-ND3 in these two genera.




Figure 1. doi

Heatmap of the mean pairwise differences between species of Caribbean octocorals, based on nucleotide substitutions in their COI sequences. Colours on the diagonal indicate the mean intra-species differences. Species with a dash in the diagonal were represented by only one individual and thus their intra-species differences could not be calculated.

Haplotype network analyses of the *Plexaura-Pseudoplexaura-Eunicea* group and the *Antillologorgia* and *Gorgonia* group, were able to distinguish some genera and species which were not easily distinguished by distance analysis as visualised in the heatmaps. All of the markers except for 16S show that *Gorgonia* and *Antillologorgia* can be separated, as each is comprised of unique haplotypes Fig. 6, although the haplotypes generally differ by only one or a few base pairs. In addition, COI appeared to distinguish the different species within each genus. For example, the COI network clearly separated *G. mariae* from *G. ventalina* but did not clearly separate the species of *Antillologorgia* from each other. However, our small sample sizes are not a strong test of the efficacy of each marker to uniquely identify




Figure 6. [doi](#)

Haplotype networks involving *Gorgonia* and *Antillogorgia* DNA sequences. Each haplotype is represented by a circle. The number of individuals within each haplotype is indicated within the corresponding circle. The circle's sizes are proportional to the number of individuals. The white areas within some circles represent individuals identified only to the genus level (species unknown), whereas other colours are species-specific. The length between haplotypes represents the number of nucleotide differences between them: 1 length unit = 1 substitution or indel (insertion/deletion).

In contrast, network analysis of the *Plexaura-Pseudoplexaura-Eunicea* group further reinforces the results of the heatmaps, indicating that these genera cannot be distinguished using any of the markers Fig. 7. For all of the markers, there were single haplotypes that were shared by *Eunicea*, *Plexaura* and *Pseudoplexaura*. *Muricea* was not clearly distinguished from the other genera using COI and 16S with the pairwise differences, but species in this genus were consistently distinguished with the network approach. These results further support the idea that the *Plexaura-Pseudoplexaura-Eunicea* group has undergone a rapid endemic radiation in the Caribbean making them particularly difficult to distinguish (Sánchez et al. 2003, Sánchez and Wishing 2005, Sánchez 2016). Likewise, both *Antillogorgia* and *Eunicea* include exemplary cases of depth-induced ecological speciation and recent divergence (Prada and Hellberg 2013, Calixto-Botía and Sánchez 2017).

As has been previously demonstrated for octocorals from other regions (Baco and Cairns 2012, France and Hoover 2002, France and Hoover 2001, McFadden et al. 2014, McFadden et al. 2004, McFadden et al. 2010), we found limited ability of the 5 markers to distinguish species based on distance methods typically employed by barcode studies Figs 1, 2, 3, 4 Fig. 5. Nevertheless, metabarcoding studies are liable to obtain COI and 16S sequences from octocorals and researchers are likely to search for species identifications in BoLD or GenBank. Our sequences are the first COI and 16S sequences for many Caribbean octocoral species in GenBank and our results demonstrate that the COI barcode sequence can distinguish Caribbean species as effectively as other mitochondrial

sequences more commonly used in octocoral-focused studies. Despite this, the BoLD database assigned our 28 species to only 9 Barcode Index Numbers (BINs; Ratnasingham and Hebert 2013) highlighting that alternate approaches, such as haplotype network analysis or character-based DNA barcoding (Rach et al. 2008), designed for small divergences, are more appropriate for attempts to use DNA sequences to automatically identify octocorals (McFadden et al. 2010; Baco and Cairns 2012).




Figure 7. [doi](#)

Haplotype networks of DNA sequences belonging to genera that could not be clearly separated by their nucleotide differences (heatmaps). For each gene, only the genera that were not clearly different in the heatmaps were included. Each haplotype is represented by a circle and each genus is represented by a unique colour. The number of individuals belonging to each haplotype are indicated within the corresponding circle. The circle's sizes are proportional to the number of individuals. The length between haplotypes represents the number of nucleotide differences (substitutions or indels) between them. Haplotypes present in multiple species, all belonging to the same genus, are indicated with a black star and haplotypes with all individuals belonging to the same species are indicated with underlined numbers. To be conservative, where samples only identified to genus were included in a circle, we did not consider them conspecific with samples identified to species, although they could have been.

Acknowledgements

We thank the Autoridad de los Recursos Acuáticos de Panamá for issuing permits for this work, the participants in the 2007 workshop on Gorgonian Biology and Systematics in Bocas del Toro and C. McFadden whose suggestions significantly improved this manuscript. This research was supported by a Grand Challenges Level 1 grant from the Smithsonian Institution. All or portions of the laboratory and computer work were conducted in and with the support of the L.A.B. facilities of the National Museum of Natural History.

Author contributions

JM– contributed to the analysis and manuscript preparation; **RC**– contributed to the analysis and manuscript preparation; **DVP**– contributed to the analysis and manuscript preparation; **ACD**– generated the sequences and contributed to the analysis; **JAS**– Collected and identified the samples; **HRL**– Collected and identified the samples

References

- Baco AR, Cairns SD (2012) Comparing molecular variation to morphological species designations in the deep-sea coral *Narella* reveals new insights into seamount coral ranges. *PLoS One* 7 (9): 1-15. <https://doi.org/10.1371/journal.pone.0045555>
- Bilewitch JP, Degnan SM (2011) A unique horizontal gene transfer event has provided the octocoral mitochondrial genome with an active mismatch repair gene that has potential for an unusual self-contained function. *BMC Evolutionary Biology* 11 (1): 228. <https://doi.org/10.1186/1471-2148-11-228>
- Brugler MR, France SC (2008) The mitochondrial genome of a deep-sea bamboo coral (Cnidaria, Anthozoa, Octocorallia, Isididae): genome structure and putative origins of replication are not conserved among octocorals. *Journal of Molecular Evolution* 67 (2): 125-136. <https://doi.org/10.1007/s00239-008-9116-2>
- Bucklin A, Steinke D, Blanco-Bercial L (2011) DNA barcoding of marine Metazoa. *Annual Review of Marine Science* 3 (1): 471-508. <https://doi.org/10.1146/annurev-marine-120308-080950>
- Bucklin A, Lindeque PK, Rodriguez-Ezpeleta N, Albaina A, Lehtiniemi M (2016) Metabarcoding of marine zooplankton: prospects, progress and pitfalls. *Journal of Plankton Research* 38 (3): 393-400. <https://doi.org/10.1093/plankt/fbw023>
- Calixto-Botía I, Sánchez JA (2017) A case of modular phenotypic plasticity in the depth gradient for the gorgonian coral *Antilloorgia bipinnata* (Cnidaria: Octocorallia). *BMC Evolutionary Biology* 17 (1): 55. <https://doi.org/10.1186/s12862-017-0900-8>
- Edmunds PJ, Lasker HR (2016) Cryptic regime shift in benthic community structure on shallow reefs in St. John, US Virgin Islands. *Marine Ecology Progress Series* 559: 1-12. <https://doi.org/10.3354/meps11900>
- Etnoyer P, Wirshing H, Sánchez J (2010) Rapid assessment of octocoral diversity and habitat on Saba Bank, Netherlands Antilles. *PLoS One* 5 (5): e10668. <https://doi.org/10.1371/journal.pone.0010668>
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular marine biology and biotechnology* 3 (5): 294-299.
- Fonseca V, Carvalho G, Sung W, Johnson H, Power D, Neill S, Packer M, Blaxter M, D. Lamshead PJ, Thomas WK, Creer S (2010) Second-generation environmental sequencing unmasks marine metazoan biodiversity. *Nature Communications* 1 (7): 98. <https://doi.org/10.1038/ncomms1095>
- France S, Hoover L (2002) DNA sequences of the mitochondrial COI gene have low levels of divergence among deep-sea octocorals (Cnidaria: Anthozoa). *Hydrobiologia* 471: 149-155. <https://doi.org/10.1023/A:1016517724749>

- France SC, Hoover LL (2001) Analysis of variation in mitochondrial DNA sequences (ND3, ND4L, MSH) among Octocorallia (=Alcyonaria) (Cnidaria: Anthozoa). *Bulletin of the Biological Society of Washington* 10: 110-118.
- Gascuel O (1997) BIONJ: an improved version of the NJ algorithm based on a simple model of sequence data. *Molecular Biology and Evolution* 14 (7): 685-695. <https://doi.org/10.1093/oxfordjournals.molbev.a025808>
- Hajibabaei M, Singer GA, Hebert PD, Hickey DA (2007) DNA barcoding: how it complements taxonomy, molecular phylogenetics and population genetics. *Trends in Genetics* 23 (4): 167-172. <https://doi.org/10.1016/j.tig.2007.02.001>
- Hebert PN, Gregory TR (2005) The promise of DNA barcoding for taxonomy. *Systematic Biology* 54 (5): 852-859. <https://doi.org/10.1080/10635150500354886>
- Huang D, Meier R, Todd PA, Chou LM (2008) Slow mitochondrial COI sequence evolution at the base of the metazoan tree and its implications for DNA barcoding. *Journal of Molecular Evolution* 66 (2): 167-174. <https://doi.org/10.1007/s00239-008-9069-5>
- Lasker HR, Coffroth MA (1983) Octocoral distributions at Carrie Bow Cay, Belize. *Marine Ecology Progress Series* 13: 21-28. <https://doi.org/10.3354/meps013021>
- Leray M, Yang J, Meyer C, Mills S, Agudelo N, Vincent R, Boehm J, Machida R (2013) Data from: A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. *Frontiers in Zoology* 10: 34. <https://doi.org/10.5061/DRYAD.6GD51>
- McFadden CS, Tullis ID, Hutchinson MB, Winner K, Sohm JA (2004) Variation in coding (NADH dehydrogenase subunits 2, 3, and 6) and noncoding intergenic spacer regions of the mitochondrial genome in Octocorallia (Cnidaria: Anthozoa). *Marine Biotechnology* 6 (6): 516-526. <https://doi.org/10.1007/s10126-002-0102-1>
- McFadden CS, Benayahu Y, Pante E, Thoma JN, Nevarez PA, France SC (2010) Limitations of mitochondrial gene barcoding in Octocorallia. *Molecular Ecology Resources* 11 (1): 19-31. <https://doi.org/10.1111/j.1755-0998.2010.02875.x>
- McFadden CS, Brown AS, Brayton C, Hunt CB, van Ofwegen LP (2014) Application of DNA barcoding in biodiversity studies of shallow-water octocorals: molecular proxies agree with morphological estimates of species richness in Palau. *Coral Reefs* 33 (2): 275-286. <https://doi.org/10.1007/s00338-013-1123-0>
- Meyer CP (2003) Molecular systematics of cowries (Gastropoda: Cypraeidae) and diversification patterns in the tropics. *Biological Journal of the Linnean Society* 79 (3): 401-459. <https://doi.org/10.1046/j.1095-8312.2003.00197.x>
- Moura CJ, Cunha MR, Porteiro FM, Rogers AD (2011) The use of the DNA barcode gene 16S mRNA for the clarification of taxonomic problems within the family Sertulariidae (Cnidaria, Hydrozoa). *Zoologica Scripta* 40 (5): 520-537. <https://doi.org/10.1111/j.1463-6409.2011.00489.x>
- Oliverio M, Barco A, Modica MV, Richter A, Mariottini P (2009) Ecological barcoding of corallivory by second internal transcribed spacer sequences: hosts of coralliophiline gastropods detected by the cnidarian DNA in their stomach. *Molecular Ecology Resources* 9 (1): 94-103. <https://doi.org/10.1111/j.1755-0998.2008.02388.x>
- Palumbi SR, Martin A, Romano S, McMillan WO, Stice L, Grabowski G (1991) *The Simple Fool's Guide to PCR, Version 2*. University of Hawaii Press, Honolulu

- Prada C, Hellberg ME (2013) Long prereproductive selection and divergence by depth in a Caribbean candelabrum coral. *Proceedings of the National Academy of Sciences* 110 (10): 3961-3966. <https://doi.org/10.1073/pnas.1208931110>
- Rach J, DeSalle R, Sarkar IN, Schierwater B, Hadrys H (2008) Character-based DNA barcoding allows discrimination of genera, species and populations in Odonata . *Proceedings of the Royal Society B: Biological Sciences* 275 (1632): 237-247. <https://doi.org/10.1098/rspb.2007.1290>
- Ratnasingham S, Herbert PDN (2007) BOLD: The Barcode of Life Data system (<http://www.barcodinglife.org>). *Molecular Ecology Notes* 7 (3): 355-364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Ratnasingham S, Hebert PN (2013) A DNA-Based Registry for All Animal Species: The Barcode Index Number (BIN) System. *PLoS ONE* 8 (7): e66213. <https://doi.org/10.1371/journal.pone.0066213>
- Ruzicka RR, Colella MA, Porter JW, Morrison JM, Kidney JA, Brinkhuis V, Lunz KS, Macaulay KA, Bartlett LA, Meyers MK, Colee J (2013) Temporal changes in benthic assemblages on Florida Keys reefs 11 years after the 1997/1998 El Niño. *Marine Ecology Progress Series* 489: 125-141. <https://doi.org/10.3354/meps10427>
- Sánchez JA, Zea S, Diaz JM (1997) Gorgonian communities of two contrasting environments from oceanic Caribbean atolls. *Bulletin of Marine Science* 61: 61-72.
- Sánchez JA, Diaz JM, Zea S (1998) Octocoral and black coral distribution patterns on the barrier reef-complex of Providencia Island, southwestern Caribbean. *Caribbean Journal of Science* 34: 250-264.
- Sánchez JA, McFadden CS, France SC, Lasker HR (2003) Molecular phylogenetic analyses of shallow-water Caribbean octocorals. *Marine Biology* 142 (5): 975-987. <https://doi.org/10.1007/s00227-003-1018-7>
- Sánchez JA, Wishing HH (2005) A Field key to the identification of tropical western Atlantic zooxanthellate octocorals (Octocorallia: Cnidaria). *Caribbean Journal of Science* 41 (3): 508-522.
- Sánchez JA (2016) Diversity and evolution of octocoral animal forests at both sides of tropical America. *Marine Animal Forests*. https://doi.org/10.1007/978-3-319-17001-5_39-1
- Shearer TL, van Oppen MJH, Romano SL, Wörheide G (2002) Slow mitochondrial DNA sequence evolution in the Anthozoa (Cnidaria). *Molecular Ecology* 11 (12): 2475-2487. <https://doi.org/10.1046/j.1365-294x.2002.01652.x>
- Stat M, Huggett MJ, Bernasconi R, DiBattista JD, Berry TE, Newman SJ, Harvey ES, Bunce M (2017) Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in a tropical marine environment. *Scientific Reports* 7 (1): 12240. <https://doi.org/10.1038/s41598-017-12501-5>
- Thomsen PF, Kielgast J, Iversen LL, Møller PR, Rasmussen M, Willerslev E (2012) Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLoS One* 7 (8): e41732. <https://doi.org/10.1371/journal.pone.0041732>
- Tsounis G, Edmunds PJ, Bramanti L, Gambrel B, Lasker HR (2018) Variability of size structure and species composition in Caribbean octocoral communities under contrasting environmental conditions. *Marine Biology* 165 (2): 1-14. <https://doi.org/10.1007/s00227-018-3286-2>

- van der Ham JL, Brugler MR, France SC (2009) Exploring the utility of an indel-rich, mitochondrial intergenic region as a molecular barcode for bamboo corals (Octocorallia: Isididae). *Marine Genomics* 2: 183-192. <https://doi.org/10.1016/j.margen.2009.10.002>
- Wirshing HH, Messing CG, Douady CJ, Reed J, Stanhope MJ, Shivji MS (2005) Molecular evidence for multiple lineages in the gorgonian family Plexauridae (Anthozoa: Octocorallia). *Marine Biology* 147 (2): 497-508. <https://doi.org/10.1007/s00227-005-1592-y>
- Zaiko A, Schimanski K, Pochon X, Hopkins GA, Goldstien S, Floerl O, Wood SA (2016) Metabarcoding improves detection of eukaryotes from early biofouling communities: implications for pest monitoring and pathway management. *Biofouling* 32 (6): 671-684. <https://doi.org/10.1080/08927014.2016.1186165>
- Zheng L, He J, Lin Y, Cao W, Zhang W (2014) 16S rRNA is a better choice than COI for DNA barcoding hydrozoans in the coastal waters of China. *Acta Oceanologica Sinica* 33 (4): 55-76. <https://doi.org/10.1007/s13131-014-0415-8>