



The first record of the genus *Belenois* (Lepidoptera: Pieridae) from China

Si-Xun Ge[‡], Shao-Ji Hu[§], Hong-Liang Shi[‡], Feng-Ying Han[‡], Ming-Jun Li[¶], Li-Li Ren[‡]

[‡] College of Forestry, Beijing Forestry University, Beijing, China

[§] Institute of International Rivers and Eco-security, Yunnan University, Kunming, China

[‡] Tongliao forest pest control and quarantine station, Tongliao, China

[¶] Kuntouhe forestry centre, Chifeng, China

Corresponding author: Si-Xun Ge (gsx_pieris@qq.com), Li-Li Ren (lily_ren@bjfu.edu.cn)

Academic editor: Martin Wiemers

Received: 24 Nov 2020 | Accepted: 08 Jan 2021 | Published: 18 Jan 2021

Citation: Ge S-X, Hu S-J, Shi H-L, Han F-Y, Li M-J, Ren L-L (2021) The first record of the genus *Belenois* (Lepidoptera: Pieridae) from China. Biodiversity Data Journal 9: e61332. <https://doi.org/10.3897/BDJ.9.e61332>

Abstract

Background

The family Pieridae is a large group of butterflies which plays an important role in evolutionary biology and contains many potential pests (Courtney 1986). Pieridae is a cosmopolitan family while the tropics harbour higher species richness. In a very recent expedition to the Chinese-Indian border area in Tibet Autonomous Region, a migratory species, *Belenois aurota* (Fabricius, 1793), was discovered for the first time, which comprises the first record of the genus *Belenois* in China and the highest altitude record of this species.

New information

The species *B. aurota* (Fabricius, 1793) is the first record of the genus *Belenois* from China. The specimen was collected at an altitude of about 3,000 m in Tibet Autonomous Region. Relevant details are presented for the species.

Keywords

Belenois aurota, Tibet, new record, high altitude distribution, migratory species

Introduction

The family Pieridae, which is a cosmopolitan family, includes over 1,000 species in 83 genera (Ackery et al. 1999, Braby 2005). The adults are of medium size and always sombre colour, typically white or yellow and with black or red patterns. Some species exhibit seasonal phenotypic variation (Braby 2005, Shou et al. 2006).

The genus *Belenois* Hübner, [1819] includes 29 species which are mainly found in the Ethiopian Region and there is only one species (*Belenois java*) that is distributed in the Oriental and Australian Realms. China is a country with mega-biodiversity, with new taxa and new records of butterflies discovered nearly every year in recent times (Hu and Zhang 2009, Hu and Zhang 2010, Hu et al. 2012a, Hu et al. 2012b, Hu et al. 2014, Hu et al. 2018, Zhang and Hu 2018, Zhang et al. 2020a, Zhang et al. 2020b). Even so, a large proportion of Chinese territory is still insufficiently surveyed.

Belenois and *Dixeia* are sister groups and belong to the subtribe Aporiina of the tribe Pierini (Wahlberg et al. 2014). The genus *Belenois* was established by Hübner with *Belenois calypso* as its type-species. This genus has 29 known species which are mainly distributed in Africa and south-west Asia and also recorded in Oceania (Shou et al. 2006). However, there is no report of this genus occurring in China.

In this contribution, a well-known migratory species *B. aurota* (Fabricius, 1793) has been collected from Tibet Autonomous Region. The migration of the African colonies of this species almost every year has received lots of attention. The biology of *B. aurota* is also well studied. *B. aurota* larvae feed on plants of the family Capparidaceae, in particular the genus *Capparis*. Under some circumstances, the population erupts at local scale and defoliates the entire bush of *Capparis*; for example, in the Rwenzori National Park in Uganda, it has been a regular pest on *C. decidua* (Chandra 1985). The specimens collected in this contribution are the first record of the genus *Belenois* from China.

Materials and methods

Photographs of the adult were taken with an interchangeable lens digital camera Olympus E-M1 with the lens M. ZUIKO DIGITAL ED 60 mm F2.8 Macro. After removal, the abdomen was soaked in 10% potassium hydroxide solution at room temperature for about 24 hours and was dissected under a Nikon SMZ18 microscope. The genitalia preparation was photographed by a Nikon D7500 digital camera attached to the microscope. Final plates were prepared in Adobe Photoshop CC (Adobe Systems Inc., San Jose, CA, USA). The specimen examined is deposited in the insect collection, Department of Forest Protection, Beijing Forestry University (BFU), Beijing.

Taxon treatment

Belenois (Anaphaeis) aurota (Fabricius, 1793)

Material

- a. scientificName: *Belenois aurota* (Fabricius, 1793); taxonomicStatus: accepted; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Lepidoptera; family: Pieridae; taxonRank: species; verbatimTaxonRank: sp.; vernacularName: Pioneer White; genus: *Belenois*; subgenus: *Anaphaeis*; specificEpithet: *aurota*; scientificNameAuthorship: (Fabricius, 1793); country: China; stateProvince: Tibet Autonomous Region; county: Zanda County; locality: East of Diya township; verbatimElevation: 2983 m; verbatimCoordinates: 31°7883'N, 78°8788'E; samplingProtocol: sweep net; year: 2019; month: 7; day: 17; habitat: Desolate valley beside the road; individualCount: 1; sex: male; lifeStage: adult; recordedBy: Sixun Ge; disposition: in collection; identifiedBy: Sixun Ge; dateIdentified: 2019; basisOfRecord: PreservedSpecimen

Description

Male (Fig. 1). Forewing length 44 mm. Body blackish dorsally covered with grey hair and whitish ventrally. Antennae blackish, in typical club shape. Both wings white on the upperside, but with a creamy yellow hue on the underside. Wings with the following markings. Forewing upperside: costa grey, a short black band at the end of the discocell, a curved black subapical band running from costa to vein CuA₁ and outwardly extending along veins to termen. Forewing underside similar to upperside, but all markings, especially the ones in the apical area, tinged with a golden-brown hue. Hindwing upperside: a black spot at the end of vein Sc+R₁, termen black fused with a series of subterminal black crescents before vein CuA₂. Hindwing underside: the previously-mentioned markings, all veins and a short discal band before vein Rs golden-brown.




Figure 1. [doi](#)

Male *Belenois aurota* (Fabricius, 1793) collected in Diya Township. **A.** upperside; **B.** underside.

Male genitalia (Fig. 2). Highly sclerotised. Ring straight of moderate width; tegumen narrow in dorsal view; uncus broad at the base, but abruptly narrowed into the same width towards the tip in the dorsal view, while strongly curved ventrally in the lateral view. Saccus slender with a flattened tip. Valve triangulate, slightly elongated distally with an acutely pointed tip, with a long harpe running through the median part. Aedeagus slender and straight, with a pointed tip, Juxta fan-shaped in posterior view.




Figure 2. [doi](#)

Male genitalia of *B. aurota*. **A.** entire genitalia; **B.** lateral view of ring; **C.** left valve; **D.** lateral view of aedeagus; **E.** dorsal view of tegumen and uncus; **F.** ventral view of juxta.

Distribution

South, Southeast and Central Asia; Tropical Africa; China (New record)

Discussion

Belenois aurota is found both in Asia and Africa, this species is migratory, the migration of the species in South Africa has received considerable attention in previous studies (Dickson and Schofield 1958, Taylor 1959), while, except for the populations of the Arabian Peninsula, few records of migration of the Asian populations have been observed; phylogenetic and phylogeographic patterns show that populations of Africa and Asia represent different clusters (Irungbam et al. 2019).

Interestingly, although there has been no official publication of this species in China, we found a specimen labelled “China, Sichuan, Mt. Kintushan, [probably Jinfo Shan in Chongqing] (ohne Datum) [no date], ex coll. A. SCHULTE. Männchen [male].” on a European butterfly website (http://www.euroleps.ch/seiten/s_art.php?art=pier_aurota). We are more inclined to recognise this specimen as a stray butterfly based on its location — far from the known distribution area and a single record for several decades.

In previous studies, the Himalayan populations of *B. aurota* were mainly distributed in areas below 1,800 m, the highest altitude recorded for the distribution of this species being 2,400 m in Nilgiris, India (Bingham 1907, Cotton et al. 2015). However, in our field investigation, an individual of *B. aurota* was collected at an altitude of about 3,000 m which is the highest altitude record of this species. Colonies of an undetermined *Capparis* species were also found around Diya Township (Fig. 3); we speculate that this species might be *Capparis himalayensis*, which could be a suitable host plant for *B. aurota* larvae. Due to the existence of the host plant, we believe that our *B. aurota* specimen, collected in the Diya Township, was highly likely from a local population rather than a stray butterfly.



Figure 3. [doi](#)

Possible larval food plant, *Capparis himalayensis*, found in Diya Township.

Acknowledgements

We thank Wei-Feng Yan (Beijing Forestry University, Beijing, China) and Ping-Zhou Zhu (Institute of Zoology, Chinese Academy of Sciences, Beijing, China) for their kind assistance in specimen collection, Hua-Zhao Li (Huazhong Agricultural University, Wuhan, China) for image processing and Adam M. Cotton (Chiang Mai, Thailand) for improving the earlier drafts of this article. We also thank the staff of General Station of Forest Pest Management in Tibet Autonomous Region for their support in our fieldwork.

References

- Ackery PR, De Jong R, Vane-Wright RI (1999) The butterflies: Hedyloidea, Hesperioidea and Papilionoidea. In: Kristensen NP (Ed.) *Lepidoptera, moths and butterflies*. 1. de Gruyter, Berlin, 263-300 pp.
- Bingham CT (1907) *The Fauna of British India, Including Ceylon and Burma*. 2. Nabu Press, Charleston, 155–158 pp.

- Braby MF (2005) Provisional checklist of genera of the Pieridae (Lepidoptera: Papilionoidea). *Zootaxa* 832 (1): 1-16. <https://doi.org/10.11646/zootaxa.832.1.1>
- Chandra H (1985) *Belenois aurota*, Fabricius (Lepidoptera: Pieridae) a serious pest of kair (*Capparis decidua*) plant. Indian Forester.
- Cotton AM, Fric ZF, Gupta IJ, Gasse PV, Smetacek P (2015) A synoptic catalogue of the butterflies of India. Butterfly Research Centre, Bhimtal & Indinov Publishing, New Delhi, 79 pp.
- Courtney SP (1986) The ecology of pierid butterflies: dynamics and interactions. *Advances in Ecological Research* 15: 51-131. [https://doi.org/10.1016/S0065-2504\(08\)60120-8](https://doi.org/10.1016/S0065-2504(08)60120-8)
- Dickson CGC, Schofield TW (1958) Observations on the migration of *Belenois aurota* (F.) (Lepidoptera: Pieridae). *Journal of the Entomological Society of south Africa* 21 (1): 427-428.
- Hu SJ, Zhang X (2009) New records and corrections of butterflies in Yunnan Province I. *Sichuan Journal of Zoology* 28 (4): 596-597.
- Hu SJ, Zhang X (2010) New records and corrections of butterflies in Yunnan Province II. *Sichuan Journal of Zoology* 29 (5): 574-577.
- Hu SJ, Zhu JQ, Zhang X (2012a) Five new records of butterflies (Lepidoptera: Lycaenidae & Hesperidae) from tropical Yunnan, China. *The Journal of Tropical Asian Entomology* 1: 32-36.
- Hu SJ, Zhu JQ, Zhang X (2012b) New records of butterflies from China (Lepidoptera: Rhopalocera): one genus and four species. *Sichuan Journal of Zoology* 31 (1): 98-101.
- Hu SJ, Zhang X, Cotton AM, Ye H (2014) Discovery of a third species of *Lamproptera* Gray, 1832 (Lepidoptera: Papilionidae). *Zootaxa* 3786 (4): 469-482. <https://doi.org/10.11646/zootaxa.3786.4.5>
- Hu SJ, Cotton AM, Condamine FL, Kuang D, Rong-Jiang W, Yu-Feng H, et al. (2018) Revision of *Pazala* Moore, 1888: the *Graphium* (*Pazala*) *mandarinus* (Oberthür, 1879) group, with treatments of known taxa and descriptions of new species and new subspecies (Lepidoptera: Papilionidae). *Zootaxa* 4441 (3): 401-440. <https://doi.org/10.11646/zootaxa.4441.3.1>
- Irungbam M, Irungbam J, Fric ZF (2019) Phylogeographic patterns of *Belenois aurota* (Fabricius, 1793) (Pieridae) on a transcontinental scale. *Asian Lepidoptera Conservation Symposium, Kolkata, West Bengal, India, 2019.9.21*.
- Shou JX, Zhou Y, Li YF (2006) *Systematic butterfly names of the World*. Shaanxi Science and Technology Press, Xi'an, 36-37 pp.
- Taylor JS (1959) A migratory flight of *Belenois aurala* (F.) at Port Elizabeth. *Journal of the Entomological Society of south Africa* 22 (1): 420-422.
- Wahlberg N, Rota J, Braby MF, Pierce NE, Wheat CW (2014) Revised systematics and higher classification of pierid butterflies (Lepidoptera: Pieridae) based on molecular data. *Zoologica Scripta* 43 (6): 641-650. <https://doi.org/10.1111/zsc.12075>
- Zhang HH, Hu SJ (2018) A new subspecies of the little known *Lethe tengchongensis* Lang, 2016 from Central Yunnan, West China. *Atalanta* 49 (1-4): 139-142.
- Zhang HH, Cotton AM, Condamine FL, Wang RJ, Hu SJ (2020a) Revision of *Pazala* Moore, 1888: the *Graphium* (*Pazala*) *alebion* and *G. (P.) tamerlanus* groups, with notes on taxonomic and distribution confusions (Lepidoptera: Papilionidae). *Zootaxa* 4759 (1): 77-97. <https://doi.org/10.11646/zootaxa.4759.1.5>

- Zhang HH, Zhu JQ, Dong RH, Yue YS, Hu SJ (2020b) New records of butterflies from China (Lepidoptera: Rhopalocera): two genera and seven species. *Sichuan Journal of Zoology* 39 (2): 214-220.