



Database of summer fish fauna sampled in river estuaries in the southern part of the Boso Peninsula, Japan

Rei Itsukushima[‡], Yuichi Kano[§]

[‡] Tokyo Institute of Technology, Yokohama, Japan

[§] Kyushu University, Fukuoka, Japan

Corresponding author: Rei Itsukushima (itsukushima.r.aa@m.titech.ac.jp)

Academic editor: Yahui Zhao

Received: 09 Apr 2021 | Accepted: 14 May 2021 | Published: 29 Jun 2021

Citation: Itsukushima R, Kano Y (2021) Database of summer fish fauna sampled in river estuaries in the southern part of the Boso Peninsula, Japan. Biodiversity Data Journal 9: e67168.

<https://doi.org/10.3897/BDJ.9.e67168>

Abstract

Background

River estuaries provide various ecosystem services, such as nutrient circulation, climate change mitigation, habitats and coastal defence. Information on the various taxonomic groups is collected from large-scale estuaries; however, few studies have focused on river estuaries of small and medium-sized rivers. In particular, information on river estuaries in peninsulas and islands with complex marine environments is lacking.

New information

This paper provides basic information on summer fish fauna in the southern part of the Boso Peninsula, Japan. The Boso Peninsula is located at the northernmost point of where the warm current (Kuroshio) reaches and is considered to have highly endemic fish fauna. In total, 28 families, 51 species and 2,908 individuals were collected from the 27 river estuaries. The data are all accessible from the document “database_fish_estuary_boso (http://ipt.pensoft.net/manage/resource.do?r=database_fish_estuary_boso)”. Further, *Sicy-*

opterus japonicus and *Microphis brachyurus*, which appear in estuaries that are influenced by the Kuroshio, were confirmed. However, these species were confirmed in few of the rivers studied, highlighting the importance of habitat conservation.

Introduction

River estuaries have complex and dynamic environments due to the influence of waves, periodic tides and mixing of freshwater and saltwater (Dyer 1997, Schroder-Adams et al. 2014). The river estuarine biological community is comprised of marine and freshwater organisms, in addition to the endemic species of brackish water (Sousa et al. 2007, Sheaves and Johnston 2008, Whitfield et al. 2012). Furthermore, the intertidal environment provides important nursery habitats for marine larvae and juvenile marine fishes (Bozeman and Dean 1980, Winkler et al. 2003, Vanalderweireldt et al. 2019). Therefore, river estuaries are especially important targets for management and conservation.

In addition, river estuaries provide various ecosystem services, such as nutrient circulation, climate change mitigation, habitats and coastal defence (Boorman 1999), estimated at \$22,832/hectare/year amongst 21 biomes (Costanza et al. 1997). Humans actively use river estuaries, which results in anthropogenic effects, such as river improvement (Cohen and Carlton 1998, Edgar et al. 2000). Studies have estimated that 61% of the world's population live in coastal areas (Alongi 1998, Bianchi 2007) and serious environmental problems, such as water degradation, disappearance of wildlife habitats and natural resource depletion, are increasing (Clark 1992, McIntyre 1995, Brown and McLachlan 2002, Kennish 2002, Howarth 2008). Information on the distribution and abundance of species is essential to conserve river estuaries where anthropogenic impacts are strong. Data regarding fish fauna in large estuaries or in the estuaries of large rivers have been collected by the National Census on River Environments (conducted by the Ministry of Land, Infrastructure, Transport and Tourism) and the National Survey on the Natural Environment (conducted by the Ministry of Environment). On the other hand, data for small and medium rivers belonging to the peninsula or islands are managed by local governments and have rarely been investigated in Japan. The lack of data has led to a shortage of knowledge about sites relevant to conservation, resulting in difficulty in determining the importance of many river estuaries.

The ocean in the southern part of the Boso Peninsula, which is the subject of this study, consists of various environments with contrasting elements, such as inner bay and open ocean, shallow and deep ocean and warm and cold currents. In particular, the Peninsula is located at the northern limit of where the Kuroshio flows along the coast of the Japanese Archipelago. As the biotas of river estuaries are strongly influenced by complex marine environments, the biota of each river is assumed to be different, although located in the same Peninsula. In this paper, we report data on fish fauna collected from 27 rivers in the southern part of the Boso Peninsula, Japan, with the aim of providing information for the conservation of the estuaries of rivers that have diverse marine environments.

Sampling methods

Sampling description: Habitats in half tide and spring tide belonging to one reach section (approximately 10 times the width of the river mouth) were selected as investigation sites. As it is known that the fish biomass and number of species in estuarine areas increases in summer (Shimamura and Nakamura 2001, Selleslagh et al. 2012), this study focused on the summer season or, more precisely, from 20 August to 3 October 2020. Surveys conducted throughout the year in the surf zones of outer Tokyo Bay (close to the study site) have shown that the highest number of fish species occurs in the summer, with more than 70% of the year's total species (Arayama et al. 2002). The fishes were collected by hand nets and throwing nets at each habitat (rapid, riffle, run and pool). For each habitat, approximately 20 net casts (half mesh 5.0 mm, 14.0 m in circumference) and 30 min of sampling with a hand net (500 mm in diameter, 6 mm mesh) were conducted. The survey was conducted from 20 August 2020 to 3 October 2020. In this research, we recorded 279 occurrence data and they were identified on site according to Okamura and Amaoka (1997), Kawanabe and Mizuno (1989), Seno (2007), Toyota and Seki (2019) and Miura (2008). Of 279 occurrence data, the tissue sections (e.g. fins of the fish) of 77 specimens were preserved for future DNA analyses in absolute ethanol (-30°C) in addition to three formalin specimens. The specimens were temporarily numbered by Y. Kano's personal acronym (QUYK) and they will be deposited in official institutes (such as The Kyushu University Museum) in the future. The dataset of this paper was registered as <https://ffish.asia/BosoBrackish>.




Figure 1. [doi](#)

Location of the study site.

Geographic coverage

Description: We surveyed 27 river estuaries in the southern part of the Boso Peninsula in Japan (Fig. 1). Watershed areas of investigated rivers ranged from 1.8 km² to 82.0 km².

Coordinates: 34.888 and 35.284 Latitude; 139.730 and 140.416 Longitude.

Taxonomic coverage

Description: Of the fish fauna, 28 families, 51 species and 2,908 individuals were collected from the 27 river estuaries (Suppl. material 1). The Nagao River had the highest number of species (13 species) and the Oobizo River had the highest number of individuals (235 individuals). By contrast, the Kawaguchi River and the Soro River presented the lowest number of species (three species each) and the Motona River the lowest number of individuals (10 individuals). The highest number of individuals found was 1,492 of *Mugil cephalus*, which appeared in all target rivers. We recorded species within the following order: Perciformes (30 species), Cypriniformes (4 species), Pleuronectiformes (3 species), Tetraodontiformes (3 species), Mugiliformes (2 species), Anguilliformes (1 species), Beloniformes (1 species), Clupeiformes (1 species), Gasterosteiformes (1 species), Gonorynchiformes (1 species), Myliobatiformes (1 species), Osmeriformes (1 species), Scorpaeniformes (1 species) and Siluriformes (1 species) (Fig. 2). We recorded species from the following families: Gobiidae (13 species), Cyprinidae (4 species), Carangidae (3 species), Sparidae (3 species), Lutjanidae (3 species), Mugilidae (2 species), Terapontidae (2 species), Adrianichthyidae (1 species), Anguillidae (1 species), Chanidae (1 species), Clupeidae (1 species), Cynoglossidae (1 species), Dasyatidae (1 species), Eleotridae (1 species), Gerreidae (1 species), Haemulidae (1 species), Lateolabracidae (1 species), Leiognathidae (1 species), Monacanthidae (1 species), Osmeridae (1 species), Paralichthyidae (1 species), Platycephalidae (1 species), Pleuronectidae (1 species), Plotosidae (1 species), Sillaginidae (1 species), Syngnathidae (1 species), Tetraodontidae (1 species) and Triacanthidae (1 species) (Fig. 3).





Figure 2. [doi](#)

Taxonomic coverage of fish fauna (by order).

Figure 3. [doi](#)

Taxonomic coverage of fish fauna (by family).

Taxa included:

Rank	Scientific Name
species	<i>Microphis brachyurus</i> (Bleeker, 1854)
species	<i>Anguilla japonica</i> Temminck & Schlegel, 1846
species	<i>Platichthys bicoloratus</i> (Basilewsky, 1855)
species	<i>Sillago japonica</i> Temminck & Schlegel, 1843
species	<i>Mugil cephalus</i> Linnaeus, 1758
species	<i>Planiliza macrolepis</i> (Smith, 1846)
species	<i>Nuclepoma nuchalis</i> (Temminck & Schlegel, 1845)
species	<i>Terapon jarbua</i> (Forsskål, 1775)
species	<i>Rhyncopelates oxyrhynchus</i> (Temminck & Schlegel, 1842)
species	<i>Gerres equulus</i> Temminck & Schlegel, 1844
species	<i>Plotosus japonicus</i> Yoshino & Kishimoto, 2008
species	<i>Paraplagusia japonica</i> Temminck & Schlegel, 1846
species	<i>Hemitygon akajei</i> (Müller & Henle, 1841)
species	<i>Plectorhynchus cinctus</i> (Temminck & Schlegel, 1843)
species	<i>Platycephalus</i> sp. <i>sensu</i> Nakabo & Kai, 2013
species	<i>Takifugu alboplumbeus</i> (Richardson, 1845)
species	<i>Scomberoides lysan</i> (Forsskål, 1775)

species	<i>Caranx sexfasciatus</i> Quoy & Gaimard, 1825
species	<i>Caranx ignobilis</i> (Forsskål, 1775)
species	<i>Favonigobius gymnauchen</i> (Bleeker, 1860)
species	<i>Acanthogobius flavimanus</i> (Temminck & Schlegel, 1845)
species	<i>Acanthogobius lactipes</i> (Hilgendorf, 1879)
species	<i>Tridentiger obscurus</i> (Temminck & Schlegel, 1845)
species	<i>Chaenogobius annularis</i> Gill, 1859
species	<i>Tridentiger trigonocephalus</i> (Gill, 1859)
species	<i>Bathygobius</i> sp. (unidentified) Bleeker, 1878
species	<i>Luciogobius</i> sp. (unidentified) Gill, 1859
species	<i>Luciogobius guttatus</i> Gill, 1859
species	<i>Rhinogobius nagoyae</i> Jordan & Seale, 1906
species	<i>Sicyopterus japonicus</i> (Tanaka, 1909)
species	<i>Tridentiger brevispinis</i> Katsuyama Arai & Nakamura, 1972
species	<i>Rhinogobius similis</i> Gill, 1859
species	<i>Eleotris oxycephala</i> Temminck & Schlegel, 1845
species	<i>Lateolabrax japonicus</i> (Cuvier, 1828)
species	<i>Stephanolepis cirrhifer</i> (Temminck & Schlegel, 1850)
species	<i>Triacanthus biaculeatus</i> (Bloch, 1786)
species	<i>Konosirus punctatus</i> (Temminck & Schlegel, 1846)
species	<i>Acanthopagrus schlegelii</i> (Bleeker, 1854)
species	<i>Rhabdosargus sarba</i> (Forsskål, 1775)
species	<i>Acanthopagrus latus</i> (Houttuyn, 1782)
species	<i>Lutjanus fulvus</i> (Forster, 1801)
species	<i>Lutjanus russellii</i> (Bleeker, 1849)
species	<i>Lutjanus argentimaculatus</i> (Forsskål, 1775)
species	<i>Paralichthys olivaceus</i> (Temminck & Schlegel, 1846)
species	<i>Pseudaspis hakonensis</i> (Günther, 1877)
species	<i>Cyprinus carpio</i> Linnaeus, 1758
species	<i>Zacco platypus</i> (Temminck & Schlegel, 1846)
species	<i>Carassius auratus</i> (Linnaeus, 1758)

species	<i>Chanos chanos</i> (Forsskål, 1775)
species	<i>Oryzias latipes</i> (Temminck & Schlegel, 1846)
species	<i>Plecoglossus altivelis</i> (Temminck & Schlegel, 1846)

Usage licence

Usage licence: Creative Commons Public Domain Waiver (CC-Zero)

Data resources

Data package title: database_fish_estuary_boso

Resource link: <https://www.gbif.org/dataset/2baad33a-e52e-4789-95ad-b288607673f8>

Alternative identifiers: http://ipt.pensoft.net/resource?r=database_fish_estuary_boso

Number of data sets: 1

Data set name: database_fish_estuary_boso

Column label	Column description
occurrenceID	An identifier for the Occurrence.
basisOfRecord	The specific nature of the data record.
eventDate	The date-time or interval during which an Event occurred.
scientificName	The full scientific name.
kingdom	The full scientific name of the kingdom in which the taxon is classified.
phylum	The full scientific name of the phylum or division in which the taxon is classified.
class	The full scientific name of the class in which the taxon is classified.
order	The full scientific name of the order in which the taxon is classified.
family	The full scientific name of the family in which the taxon is classified.
taxonRank	The taxonomic rank of the most specific name in the scientificName as it appears in the original record.
identifiedBy	A list (concatenated and separated) of names of people, groups or organisations who assigned the Taxon to the subject.
decimalLatitude	The geographic latitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location.
decimalLongitude	The geographic longitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic centre of a Location.

geodeticDatum	The ellipsoid, geodetic datum or spatial reference system (SRS) upon which the geographic coordinates given in decimalLatitude and decimalLongitude are based.
countryCode	The standard code for the country in which the Location occurs. Recommended best practice is to use ISO 3166-1-alpha-2 country codes.
individualCount	The number of individuals represented present at the time of the Occurrence.
organismQuantity	A number or enumeration value for the quantity of organisms.
organismQuantityType	The type of quantification system used for the quantity of organisms.
habitat	A category or description of the habitat in which the Event occurred.
catalogNumber	A list (concatenated and separated) of previous or alternative fully qualified catalogue numbers or other human-used identifiers for the same Occurrence, whether in the current or any other data set or collection.
language	A language of the resource. Recommended best practice is to use a controlled vocabulary, such as RFC 4646 [RFC4646]
country	The name of the country or major administrative unit in which the Location occurs. Recommended best practice is to use a controlled vocabulary such as the Getty Thesaurus of Geographic Names.
stateProvince	The name of the next smallest administrative region than country (state, province, canton, department, region etc.) in which the Location occurs.
municipality	The full, unabbreviated name of the next smallest administrative region than county (city, municipality etc.) in which the Location occurs. Do not use this term for a nearby named place that does not contain the actual location.
locality	The specific description of the place. Less specific geographic information can be provided in other geographic terms (higherGeography, continent, country, stateProvince, county, municipality, waterBody, island, islandGroup). This term may contain information modified from the original to correct perceived errors or standardise the description.
modified	The most recent date-time on which the resource was changed. For Darwin Core, recommended best practice is to use an encoding scheme, such as ISO 8601:2004(E).
year	The four-digit year in which the Event occurred, according to the Common Era Calendar.
month	The ordinal month in which the Event occurred.
day	The integer day of the month on which the Event occurred.
locationID	An identifier for the set of location information (data associated with dcterms:Location). May be a global unique identifier or an identifier specific to the dataset.

Additional information

Fish fauna of the Pacific Ocean side of the Japanese Archipelago has been strongly influenced by the dispersal and vicariance of the Kuroshio (Senou et al. 2006). Itsukushima

(2019) classified the fish fauna of the large rivers belonging to the Japanese Archipelago and showed that rivers flowing into the Pacific Ocean had different migratory fish that appeared depending on the presence or absence of the influence of the Kuroshio and that the boundary of the classification of fish fauna is near the Boso Peninsula. The Boso Peninsula is located at the northernmost point of the Kuroshio and is considered to have highly endemic fish fauna due to its influence. As a result of this survey, appearance of *Sicyopterus japonicus* and *Micropis brachyurus*, which appear in estuaries that are influenced by the Kuroshio (Dotu and Mito 1955, Nakazato and Fujita 1986), were confirmed in the Nagao and Sugai Rivers, respectively. These two species are known to be warm-water species dependent on the Kuroshio Current, although they have been confirmed in the north of the Boso Peninsula (Hata 2020). Both rivers are located on the Pacific side of the Archipelago, near the southern tip of the Boso Peninsula and the strong influence of the Kuroshio may be the reason for the appearance of these species. These species are widely distributed in rivers influenced by the Kuroshio; however, there are few confirmed in the rivers of the Boso Peninsula and they are important as a local population. Furthermore, the distribution of fish species that are thought to be dispersed by the Kuroshio is assumed to have moved northwards due to the rise in seawater temperature caused by climate change (Yamakawa et al. 2020). In addition, the velocity of the Kuroshio is reported to increase by 30% over 100 years (Sakamoto et al. 2005), which may lead to changes in the distribution area of dispersed species and fish fauna in the Boso Peninsula. Therefore, the fish fauna data, obtained in this study, are crucial because they provide the basis for climate change impact assessments in each river.

Of the rivers surveyed, the mouths of the Soro and Shinmei Rivers were the only ones completely closed and the number of fish species were only three and five, respectively. In these two rivers, unlike the others, river mouth closure had occurred, blocking the movement between the river and the ocean. Although there is a variety of factors that degrade estuarine biota (McKinley et al. 2011, Itsukushima et al. 2019, Schulz et al. 2020), no significant differences in habitat or water quality were identified between these two rivers and the other rivers surveyed. Therefore, we concluded that mouth closure had influenced the decline of fish species. In addition to water quality degradation and salinity reduction in brackish waters (Uno et al. 2014, Watanabe et al. 2015), river mouth closure leads to fish migration impediments from marine to river habitats and disruption of spawning accretion (McDowall 1995, Kanda et al. 2009). Furthermore, the degree of river mouth closure influences the biodiversity of the system, with the number of species at its lowest in case of complete closure (Torii et al. 2011). The results of this study also indicate that the number of species was significantly reduced in completely closed river mouths, suggesting that dredging to maintain openings of river channels and other habitat protection measures are needed to improve the habitats for fishes.

This study was conducted during the summer season when the species diversity and biomass were the highest. However, several migratory species—which seasonally utilised estuarine habitats during this survey period—have not been identified and some species that utilise estuarine habitats only during winter may not have been sampled. For example, species such as *Ophieleotris* sp.1 of Akihito et al., 2013 and *Oxyurichthys lonchotus*

(Lenkins, 1903) have been reported to be present in the target area (Yamakawa et al. 2018). Therefore, it is necessary to conduct surveys in each season over multiple years. This survey, however, was conducted over a wide area at the boundary of biogeography—where no data had been previously obtained—and thus presents valuable data.

Enter subsection title

Enter subsection text

Acknowledgements

This work was supported by JSPS KAKENHI Grant Number JP19H02250 and JP20HP8020.

References

- Alongi DM (1998) Coastal Ecosystem Processes. CRC Press, New York.
- Arayama K, Imai H, Kanou K, Kohno H (2002) Ichthyofauna of surf zones in the outer Tokyo Bay. *La mer* 40: 59-70. [In Japanese].
- Bianchi TS (2007) Biogeochemistry of Estuaries. Oxford University Press
- Boorman LA (1999) Salt marshes - present functioning and future change. *Mangroves and Salt Marshes* 3: 227-24. <https://doi.org/10.1023/A:1009998812838>
- Bozeman EL, Dean JM (1980) The abundance of estuarine larval and juvenile fish in a South Carolina intertidal creek. *Estuaries* 3 (2): 89-97. <https://doi.org/10.2307/1351552>
- Brown AC, McLachlan A (2002) Sandy shore ecosystems and the threats facing them: Some predictions for the year 2025. *Environmental Conservation* 29 (1): 62-77.
- Clark RB (1992) Marine Pollution. 3rd edition. Clarendon Press
- Cohen AN, Carlton JT (1998) Accelerating invasion rate in a highly invaded estuary. *Science* 279 (5350): 555-558. <https://doi.org/10.1126/science.279.5350.555>
- Costanza R, D'Arge R, De Groot R, Farber S, Grasso M, Hannon B, Limburg K, Naeem S, O'Neill RV, Paruelo J, Raskin RG, Sutton P, Van Den Belt M (1997) The value of the world's ecosystem services and natural capital. *Nature* 387 (253): 260.
- Dotu Y, Mito S (1955) Life history of a Gobioid fish, *Sicyopterus japonicum* Tanaka. *Science Bulletin of the Faculty of Agriculture, Kyushu University* 15: 213-221.
- Dyer KR (1997) Estuaries, a physical introduction. 2nd edition. John Wiley & Sons, Chichester.
- Edgar GJ, Barrett NS, Graddon DJ, Last PR (2000) The conservation significance of estuaries: A classification of Tasmanian estuaries using ecological, physical and demographic attributes as a case study. *Biological Conservation* 92 (3): 383-397. [https://doi.org/10.1016/S0006-3207\(99\)00111-1](https://doi.org/10.1016/S0006-3207(99)00111-1)
- Hata K (2020) The first recorded of warm water fishes collected from rivers in Miyagi Prefecture, Japan. *Izunuma-Uchinuma Wetland Researches* 14: 69-80. [In Japanese]. https://doi.org/10.20745/izu.14.0_69

- Howarth RW (2008) Coastal nitrogen pollution: A review of sources and trends globally and regionally. *Harmful Algae* 8 (1): 14-20. <https://doi.org/10.1016/j.hal.2008.08.015>
- Itsukushima R (2019) Study of aquatic ecological regions using fish fauna and geographic archipelago factors. *Ecological Indicators* 96: 69-80. <https://doi.org/10.1016/j.ecolind.2018.08.057>
- Itsukushima R, Yoshikawa Y, Morita K (2019) Relationship between physical environmental factors and presence of molluscan species in medium and small river estuaries. *Estuarine, Coastal and Shelf Science* 218 (5): 300-309. <https://doi.org/10.1016/j.ecss.2019.01.001>
- Kanda T, Uehara S, Shibuno T (2009) Fish fauna in inland water of Ishigaki island, Yaeyama archipelago, Japan. *Bulletin of the Faculty of Agriculture, University of Miyazaki* 55: 13-24.
- Kawanabe H, Mizuno N (Eds) (1989) *Freshwater fishes of Japan*. YAMA-KEI Publishers Co., Ltd, Tokyo. [In Japanese]. [ISBN 4635056066]
- Kennish MJ (2002) Environmental threats and environmental future of estuaries. *Environmental Conservation* 29 (1): 78-107. <https://doi.org/10.1017/S0376892902000061>
- McDowall RM (1995) Seasonal pulses in migrations of New Zealand diadromous fish and the potential impacts of river mouth closure. *New Zealand Journal of Marine and Freshwater Research* 29: 517-526. <https://doi.org/10.1080/00288330.1995.9516684>
- McIntyre AD (1995) Human impact on the oceans: The 1990s and beyond. *Marine Pollution Bulletin* 31 (4-12): 147-151. [https://doi.org/10.1016/0025-326X\(95\)00099-9](https://doi.org/10.1016/0025-326X(95)00099-9)
- McKinley AC, Miskiewicz A, Taylor MD, Johnston EL (2011) Strong links between metal contamination, habitat modification and estuarine larval fish distributions. *Environmental Pollution* 159 (6): 1499-1509. <https://doi.org/10.1016/j.envpol.2011.03.008>
- Miura T (2008) *A Picture book of tidal flat organisms*. Nanpo-Shinsha, Kagoshima. [In Japanese]. [ISBN 4861241391]
- Nakazato Y, Fujita S (1986) Distribution, spawning habit, development of eggs and prelarvae of the pipefish, *Oostethus brachyurus brachyurus*. *The Aquiculture* 33 (4): 230-239.
- Okamura S, Amaoka K (Eds) (1997) *Sea fishes of Japan*. YAMA-KEI Publishers Co., Ltd, Tokyo. [In Japanese]. [ISBN 4-635-09027-2]
- Sakamoto TT, Hasumi H, Ishii M, Emori S, Suzuki T, Nishimura T, Sumi A (2005) Responses of the Kuroshio and the Kuroshio Extension to global warming in a high-resolution climate model. *Geophysical Research Letters* 32: L14617.
- Schroder-Adams CJ, Boyd RL, Tran T (2014) Estuarine foraminiferal biofacies pattern compared to the brackish ichnofacies model: Port Stephens, southeast Australia. *Estuarine, Coastal and Shelf Science* 139: 78-87. <https://doi.org/10.1016/j.ecss.2013.12.006>
- Schulz K, Stevens PW, Hill JE, Trotter AA, Ritch JL, Tuckett QM, Patterson JT (2020) Coastal restoration evaluated using dominant habitat characteristics and associated fish communities. *PLoS One* 22: e0240623. <https://doi.org/10.1371/journal.pone.0240623>
- Selleslagh J, Lobry J, N'Zigou AR, Bachelet G, Blanchet H, Chaalali A, Sautour B, Boëta P (2012) Seasonal succession of estuarine fish, shrimps, macrozoobenthos and plankton: Physico-chemical and trophic influence. The Gironde estuary as a case study. *Estuarine, Coastal and Shelf Science* 112: 243-254. <https://doi.org/10.1016/j.ecss.2012.07.030>

- Seno H (Ed.) (2007) A Photographic Guide to the Gobioid Fishes of Japan. 2nd. Heibonsha, Tokyo. [In Japanese]. [ISBN 4582542360]
- Senou H, Matsuura K, Shinohara G (2006) Checklist of fishes in the Sagami Sea with zoogeographical comments on shallow water fishes occurring along the coastlines under the influence of the Kuroshio Current. *Memoirs of the National Museum of Nature and Science* 41: 389-54.
- Sheaves M, Johnston R (2008) Influence of marine and freshwater connectivity on the dynamics of subtropical estuarine wetland fish metapopulations. *Marine Ecology Progress Series* 357: 225-243. <https://doi.org/10.3354/meps07292>
- Shimamura K, Nakamura M (2001) Succession of biotic community on an artificial reef in the coast of Lake Nakaumi, brackish water, Shimane, Japan. *Aquaculture Science* 49 (3): 299-304. [In Japanese]. <https://doi.org/10.11233/aquaculturesci1953.49.299>
- Sousa R, Antunes C, Guilhermino L (2007) Species composition and monthly variation of the molluscan fauna in the freshwater subtidal area of the River Minho estuary. *Estuarine, Coastal and Shelf Science* 75 (1-2): 90-100. <https://doi.org/10.1016/j.ecss.2007.02.020>
- Torii T, Shione H, Kato K, Sugiura Y, Kurokawa T, Oono M, Oshiro T, Arakaki T (2011) Effects of river mouth closure on fish fauna in tidal river areas. *Ecology and Civil Engineering* 13 (2): 123-139. <https://doi.org/10.3825/ece.13.123>
- Toyota K, Seki S (2019) Fresh and brackish water shrimps and crabs of Japan. Midori-Shobo, Tokyo. [In Japanese]. [ISBN 4895313913]
- Uno K, Kishimoto S, Kinoshita A, Tsujimoto G, Kakinoki T (2014) Annual change of water environment and topographic feature at urban river mouth. *Journal of Japan Society of Civil Engineers, Series B3, Ocean Development* 70: I_558-I_563.
- Vanalderweireldt L, Winkler G, Mingelbier M, Sirois P, Hidalgo M (2019) Early growth, mortality, and partial migration of striped bass (*Morone saxatilis*) larvae and juveniles in the St. Lawrence estuary, Canada. *ICES Journal of Marine Science* 76 (7): 2235-2246. <https://doi.org/10.1093/icesjms/fsz116>
- Watanabe K, Okonobi T, Imai Y (2015) Investigating of saline water intrusion and tidal discharge for different type of river mouths at the Sea of Japan side. *Journal of Japan Society of Civil Engineers, Series B2, Coastal Engineering* 71: I_409-I_414. https://doi.org/10.2208/kaigan.71.I_409
- Whitfield AK, Elliott M, Basset A, Blaber SJ, West RJ (2012) Paradigms in estuarine ecology - A review of the Remane diagram with a suggested revised model for estuaries. *Estuarine, Coastal and Shelf Science* 97: 78-90. <https://doi.org/10.1016/j.ecss.2011.11.026>
- Winkler G, Dodson JJ, Bertrand N, Thivierge D, Vincent WF (2003) Trophic coupling across the St. Lawrence river estuarine transition zone. *Marine Ecology Progress Series* 251: 59-71. <https://doi.org/10.3354/meps251059>
- Yamakawa U, Mitsui S, Maruyama T, Kato S, Sakai S, Senou H (2018) Notes on Eighteen Fish Species Recorded from the Rivers and Coastal Areas of Sagami Bay and Adjacent Waters, Japan: Northward Range Extension of Warm Water Fishes in Recent Years. *Bull. Kanagawa prefect. Mus. (Nat. Sci)* 47: 35-57. https://doi.org/10.32225/bkpmnh.2018.47_35
- Yamakawa U, Senou H, Mitsui S, Oda T, Morita Y, Taketo A, Maruyama T, Tanaka S, Saito H, Tsuda Y (2020) Records of seven fish species showing a northward shift in

distribution pattern in the Sagami Bay region, Japan. Natural History Report of Kanagawa 41: 71-8.

Supplementary material

Suppl. material 1: List of populations of fish species from 27 rivers in the southern part of the Boso Peninsula, Japan [doi](#)

Authors: Itsukushima R and Kano Y

Data type: occurrences

Brief description: List of populations of fish species from 27 rivers in the southern part of the Boso Peninsula, Japan

[Download file](#) (15.92 kb)